

Ohio Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	8.0	6.3	12.7	4.3 *
Neonatal Mortality Rate	5.3	4.2	8.6	2.8 *
Post-Neonatal Mortality Rate	2.7	2.1	4.1	**
Fetal Mortality Rate	7.1	6.4	8.9	**
Perinatal Mortality Rate	7.6	6.5	10.7	2.2 *
SUIDS Mortality Rate	1.35	1.04	2.10	**
Preterm birth percent	11.9	11.0	14.2	10.7
Late preterm birth percent	8.2	7.9	9.2	7.9
Early preterm birth percent	3.7	3.2	5.0	2.9
Extreme preterm birth percent	2.2	1.8	3.3	1.5 *
Early term birth percent	27.3	26.3	29.1	28.5
Low birthweight birth percent	10.8	9.3	14.5	8.7
Moderately low birthweight birth percent	8.9	7.8	11.5	7.3
Very low birthweight birth percent	2.0	1.5	3.0	1.4 *
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	34.1	34.4	35.5	28.9 *
Prenatal care received in first trimester percent	64.3	67.6	59.2	60.4
Adequacy of prenatal care percent	69.4	73.4	62.0	65.3
Mother's smoking abstinence third trimester percent	80.3	74.1	90.4 *	93.3 *
Mother's alcohol abstinence third trimester percent	99.7 *	99.7 *	99.5 *	**
Mother had breast fed baby at discharge percent	61.4	60.2	61.1	69.7
Normal weight pre-pregnancy percent	31.9	32.5	29.7	29.2
Obese weight pre-pregnancy percent	34.7	33.8	37.8	37.0
Appropriate weight gain during pregnancy percent	25.3	25.4	24.5	26.4

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

OEI Counties Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	8.7	6.3	12.4	4.6 *
Neonatal Mortality Rate	5.9	4.4	8.2	2.7 *
Post-Neonatal Mortality Rate	2.8	1.9 *	4.1	**
Fetal Mortality Rate	7.4	6.7	8.5	**
Perinatal Mortality Rate	8.4	7.2	10.4	2.7 *
SUIDS Mortality Rate	1.49	1.03	2.04	**
Preterm birth percent	12.5	11.3	14.2	10.6
Late preterm birth percent	8.4	7.9	9.2	7.8
Early preterm birth percent	4.0	3.4	5.0	2.8
Extreme preterm birth percent	2.4	1.9	3.2	1.4 *
Early term birth percent	27.8	26.3	29.2	29.4
Low birthweight birth percent	11.9	9.8	14.6	9.0
Moderately low birthweight birth percent	9.7	8.2	11.6	7.6
Very low birthweight birth percent	2.2	1.6	3.0	1.4 *
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	33.2	33.3	35.2	27.4 *
Prenatal care received in first trimester percent	61.3	64.7	58.9	60.0
Adequacy of prenatal care percent	65.1	69.0	61.4	64.9
Mother's smoking abstinence third trimester percent	84.6	75.8	91.1 *	94.6 *
Mother's alcohol abstinence third trimester percent	99.6 *	99.6 *	99.5 *	**
Mother had breast fed baby at discharge percent	63.7	62.7	62.2	72.7
Normal weight pre-pregnancy percent	31.7	32.8	29.7	29.3
Obese weight pre-pregnancy percent	34.6	32.7	37.8	35.9
Appropriate weight gain during pregnancy percent	25.0	25.1	24.4	26.1

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Non-OEI Counties Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	6.9	6.3	14.8	**
Neonatal Mortality Rate	4.5	4.0 *	11.0	**
Post-Neonatal Mortality Rate	2.4	2.3	**	**
Fetal Mortality Rate	6.6	6.2	11.7	**
Perinatal Mortality Rate	6.4	6.0	12.6	**
SUIDS Mortality Rate	1.16	1.04	**	**
Preterm birth percent	11.1	10.8	14.0	11.2
Late preterm birth percent	7.9	7.8	9.0	8.1
Early preterm birth percent	3.2	3.0	5.0	3.0
Extreme preterm birth percent	1.9	1.7	3.5	1.8
Early term birth percent	26.6	26.3	28.5	26.7
Low birthweight birth percent	9.4	8.9	14.2	8.3
Moderately low birthweight birth percent	7.8	7.4	11.2	6.7
Very low birthweight birth percent	1.6	1.5	3.0	1.5
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	35.2	35.2	37.6	32.4
Prenatal care received in first trimester percent	68.6	69.6	61.6	61.5
Adequacy of prenatal care percent	75.2	76.4	66.9	66.3
Mother's smoking abstinence third trimester percent	74.6	73.0	84.7	90.6 *
Mother's alcohol abstinence third trimester percent	99.8 *	99.8 *	**	**
Mother had breast fed baby at discharge percent	58.2	58.4	52.8	63.3
Normal weight pre-pregnancy percent	32.1	32.2	29.8	28.9
Obese weight pre-pregnancy percent	34.8	34.6	38.0	39.2
Appropriate weight gain during pregnancy percent	25.6	25.5	25.2	27.1

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Butler County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	7.1	**	**	**
Neonatal Mortality Rate	4.9	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**
Fetal Mortality Rate	7.5	7.9	**	**
Perinatal Mortality Rate	9.3	9.3	**	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	10.8	10.9	12.0	10.2
Late preterm birth percent	8.0	7.8	9.7	7.4
Early preterm birth percent	2.8	3.1	**	**
Extreme preterm birth percent	1.7	1.8	**	**
Early term birth percent	26.6	25.4	31.7	26.3
Low birthweight birth percent	8.8	8.9	10.8	7.7 *
Moderately low birthweight birth percent	7.4	7.6	9.2	6.5
Very low birthweight birth percent	1.3 *	1.3 *	**	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	32.6	35.1	33.3	23.5 *
Prenatal care received in first trimester percent	61.4	61.2	64.8	58.6
Adequacy of prenatal care percent	71.2	71.7	72.0	67.9
Mother's smoking abstinence third trimester percent	81.2	74.0	92.7 *	**
Mother's alcohol abstinence third trimester percent	**	**	**	**
Mother had breast fed baby at discharge percent	54.7	50.6	57.8	69.2
Normal weight pre-pregnancy percent	31.5	31.8	27.6	26.5
Obese weight pre-pregnancy percent	34.5	35.1	38.9	32.9
Appropriate weight gain during pregnancy percent	25.5	25.8	22.4	28.1

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Cuyahoga County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	10.6	6.0 *	13.8	**
Neonatal Mortality Rate	7.4	5.0	9.0	**
Post-Neonatal Mortality Rate	3.2	**	4.7	**
Fetal Mortality Rate	8.1	6.3	9.4	**
Perinatal Mortality Rate	9.3	6.0	11.5	**
SUIDS Mortality Rate	1.56	**	2.26	**
Preterm birth percent	14.2	11.2	16.5	12.4
Late preterm birth percent	9.6	7.9	11.0	9.4
Early preterm birth percent	4.6	3.3	5.6	2.9
Extreme preterm birth percent	3.0	1.8	3.9	1.6
Early term birth percent	28.6	26.2	30.1	29.0
Low birthweight birth percent	13.7	10.0	16.5	10.5
Moderately low birthweight birth percent	10.9	8.4	12.8	9.0
Very low birthweight birth percent	2.8	1.6	3.7	1.5
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	29.3 *	31.0	29.4 *	25.1 *
Prenatal care received in first trimester percent	60.9	67.1	57.0	70.4
Adequacy of prenatal care percent	62.9	69.1	58.6	69.4
Mother's smoking abstinence third trimester percent	89.2 *	81.7	92.9 *	93.3 *
Mother's alcohol abstinence third trimester percent	99.6 *	99.6 *	99.6 *	**
Mother had breast fed baby at discharge percent	65.9	69.9	62.1	74.4
Normal weight pre-pregnancy percent	29.7	32.6	27.3	25.6
Obese weight pre-pregnancy percent	37.6	32.7	41.3	42.1
Appropriate weight gain during pregnancy percent	25.9	26.2	25.6	25.0

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Franklin County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	7.5	6.4	9.9	**
Neonatal Mortality Rate	4.9	4.1 *	6.5	**
Post-Neonatal Mortality Rate	2.5	**	3.4	**
Fetal Mortality Rate	6.6	7.2	6.9	**
Perinatal Mortality Rate	7.1	8.2	7.6	**
SUIDS Mortality Rate	1.61	**	**	**
Preterm birth percent	11.6	11.8	11.8	10.1
Late preterm birth percent	7.6	8.0	7.3	7.3
Early preterm birth percent	4.0	3.8	4.5	2.8
Extreme preterm birth percent	2.4	2.3	2.8	1.7
Early term birth percent	26.2	26.2	26.0	26.7
Low birthweight birth percent	11.3	10.5	12.3	7.5 *
Moderately low birthweight birth percent	9.1	8.5	9.7	5.8 *
Very low birthweight birth percent	2.2	2.1	2.6	1.7
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	35.9	32.8	41.6	27.3 *
Prenatal care received in first trimester percent	59.2	62.9	57.2	57.3
Adequacy of prenatal care percent	59.5	62.9	57.2	59.0
Mother's smoking abstinence third trimester percent	87.0	76.5	92.8 *	97.3 *
Mother's alcohol abstinence third trimester percent	99.7 *	99.5 *	99.7 *	**
Mother had breast fed baby at discharge percent	68.9	64.3	69.8	80.9
Normal weight pre-pregnancy percent	33.7	32.4	34.1	28.2
Obese weight pre-pregnancy percent	31.6	32.1	32.3	35.7
Appropriate weight gain during pregnancy percent	26.1	26.7	24.1	29.8

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Hamilton County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	9.8	**	14.3	**
Neonatal Mortality Rate	7.1	**	10.6	**
Post-Neonatal Mortality Rate	2.7	**	3.7	**
Fetal Mortality Rate	6.0	6.4	6.3	**
Perinatal Mortality Rate	8.6	4.9 *	12.0	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	12.6	10.9	14.2	9.0 *
Late preterm birth percent	8.1	7.1	8.7	6.5 *
Early preterm birth percent	4.5	3.8	5.5	2.4
Extreme preterm birth percent	2.5	1.9	3.2	**
Early term birth percent	29.7	26.1	31.2	34.0
Low birthweight birth percent	12.7	9.5	15.3	8.1
Moderately low birthweight birth percent	10.4	8.0	12.2	7.3
Very low birthweight birth percent	2.3	1.5	3.1	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	32.4	31.5	35.2	22.3 *
Prenatal care received in first trimester percent	62.6	62.9	64.5	53.3
Adequacy of prenatal care percent	68.2	69.5	68.3	63.5
Mother's smoking abstinence third trimester percent	84.6	71.3	91.2 *	96.6 *
Mother's alcohol abstinence third trimester percent	99.6 *	**	99.5 *	**
Mother had breast fed baby at discharge percent	59.1	57.7	57.3	75.8
Normal weight pre-pregnancy percent	31.8	33.9	28.6	37.2
Obese weight pre-pregnancy percent	33.5	29.3 *	38.1	25.3 *
Appropriate weight gain during pregnancy percent	23.3	22.4	23.1	26.1

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Lucas County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	9.0	6.2	14.1	**
Neonatal Mortality Rate	6.7	5.1	9.7	**
Post-Neonatal Mortality Rate	**	**	**	**
Fetal Mortality Rate	6.4	6.1	7.4	**
Perinatal Mortality Rate	9.0	8.7	10.4	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	11.6	10.4	13.4	9.6
Late preterm birth percent	8.2	7.6	9.0	6.3 *
Early preterm birth percent	3.4	2.8	4.5	3.3
Extreme preterm birth percent	1.9	1.7	2.4	**
Early term birth percent	29.0	28.5	30.1	31.7
Low birthweight birth percent	11.5	9.9	14.1	10.7
Moderately low birthweight birth percent	9.7	8.2	11.9	9.1
Very low birthweight birth percent	1.9	1.7	2.2	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	38.4	38.1	38.9	36.5
Prenatal care received in first trimester percent	59.9	64.6	52.8	60.1
Adequacy of prenatal care percent	64.1	70.0	54.7	65.4
Mother's smoking abstinence third trimester percent	83.8	78.5	90.4 *	89.9 *
Mother's alcohol abstinence third trimester percent	99.5 *	**	**	**
Mother had breast fed baby at discharge percent	55.9	58.9	50.0	57.4
Normal weight pre-pregnancy percent	29.9	30.6	28.5	28.6
Obese weight pre-pregnancy percent	37.8	35.9	41.0	42.2
Appropriate weight gain during pregnancy percent	23.8	24.7	22.3	21.6

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Mahoning County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	6.8	**	**	**
Neonatal Mortality Rate	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**
Fetal Mortality Rate	7.4	**	**	**
Perinatal Mortality Rate	7.4	**	**	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	13.3	12.4	14.9	14.5
Late preterm birth percent	9.9	9.6	10.6	10.4
Early preterm birth percent	3.5	2.8	4.3	**
Extreme preterm birth percent	1.7	1.2 *	2.6	**
Early term birth percent	33.2	31.5	36.3	36.4
Low birthweight birth percent	11.8	9.9	15.7	13.3
Moderately low birthweight birth percent	10.4	8.9	13.6	12.1
Very low birthweight birth percent	1.4 *	**	2.1	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	36.2	35.6	38.4	33.0
Prenatal care received in first trimester percent	71.1	76.3	64.1	70.6
Adequacy of prenatal care percent	75.0	81.9 *	65.3	76.5
Mother's smoking abstinence third trimester percent	74.5	71.0	76.3	81.9
Mother's alcohol abstinence third trimester percent	99.3 *	**	**	**
Mother had breast fed baby at discharge percent	55.5	59.0	48.3	58.4
Normal weight pre-pregnancy percent	33.4	35.0	33.5	29.8
Obese weight pre-pregnancy percent	33.9	31.0	36.1	41.5
Appropriate weight gain during pregnancy percent	25.3	27.0	25.1	20.5

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Montgomery County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	6.5	5.8 *	8.6	**
Neonatal Mortality Rate	4.3	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**
Fetal Mortality Rate	8.0	**	13.0	**
Perinatal Mortality Rate	7.8	**	12.4	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	13.7	12.9	15.1	9.3 *
Late preterm birth percent	9.4	9.3	9.6	7.0
Early preterm birth percent	4.3	3.6	5.5	**
Extreme preterm birth percent	2.6	2.1	3.4	**
Early term birth percent	27.0	25.3	28.2	29.1
Low birthweight birth percent	11.8	9.8	14.6	9.3
Moderately low birthweight birth percent	9.7	8.1	11.9	8.7
Very low birthweight birth percent	2.1	1.7	2.7	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	32.6	31.3	35.1	28.0 *
Prenatal care received in first trimester percent	61.5	63.6	59.6	55.0
Adequacy of prenatal care percent	68.9	70.2	67.1	72.6
Mother's smoking abstinence third trimester percent	85.5	79.0	91.7 *	**
Mother's alcohol abstinence third trimester percent	99.5 *	**	**	**
Mother had breast fed baby at discharge percent	64.5	64.5	62.2	73.1
Normal weight pre-pregnancy percent	29.7	32.2	26.4	22.8
Obese weight pre-pregnancy percent	36.5	34.3	41.0	34.2
Appropriate weight gain during pregnancy percent	24.9	23.1	26.4	24.8

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Stark County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	9.5	10.0	**	**
Neonatal Mortality Rate	5.4	5.9	**	**
Post-Neonatal Mortality Rate	**	**	**	**
Fetal Mortality Rate	9.4	9.9	**	**
Perinatal Mortality Rate	9.9	11.1	**	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	11.1	10.9	11.7	12.6
Late preterm birth percent	7.5	7.5	6.9	10.8
Early preterm birth percent	3.6	3.4	4.8	**
Extreme preterm birth percent	2.0	1.5 *	4.1	**
Early term birth percent	24.8	24.7	26.0	28.8
Low birthweight birth percent	10.0	9.4	12.0	**
Moderately low birthweight birth percent	8.3	8.2	8.5	**
Very low birthweight birth percent	1.6	1.2 *	3.4	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	35.4	35.6	35.2	37.5
Prenatal care received in first trimester percent	61.2	62.0	59.9	36.2
Adequacy of prenatal care percent	66.6	68.0	63.8	43.5
Mother's smoking abstinence third trimester percent	74.1	71.5	82.5	90.1 *
Mother's alcohol abstinence third trimester percent	**	**	**	**
Mother had breast fed baby at discharge percent	58.5	59.0	57.9	40.2
Normal weight pre-pregnancy percent	32.4	32.9	31.4	41.3
Obese weight pre-pregnancy percent	33.2	32.6	34.8	23.9 *
Appropriate weight gain during pregnancy percent	23.8	23.6	24.6	24.8

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Summit County Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	9.9	7.0	15.6	**
Neonatal Mortality Rate	5.6	**	9.5	**
Post-Neonatal Mortality Rate	4.3	**	**	**
Fetal Mortality Rate	9.1	**	13.7	**
Perinatal Mortality Rate	8.5	**	13.8	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	12.1	10.4	15.6	9.3 *
Late preterm birth percent	8.0	7.1	10.0	**
Early preterm birth percent	4.1	3.3	5.5	**
Extreme preterm birth percent	2.5	1.7	3.9	**
Early term birth percent	26.9	25.0	29.5	26.2
Low birthweight birth percent	11.9	9.2	16.1	**
Moderately low birthweight birth percent	9.5	7.8	12.3	**
Very low birthweight birth percent	2.4	1.4 *	3.8	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	30.0	33.1	29.9	27.4 *
Prenatal care received in first trimester percent	61.5	67.5	57.9	55.7
Adequacy of prenatal care percent	67.9	71.4	64.0	68.0
Mother's smoking abstinence third trimester percent	79.7	70.5	87.5	**
Mother's alcohol abstinence third trimester percent	99.6 *	**	**	**
Mother had breast fed baby at discharge percent	71.9	71.4	68.6	88.8 *
Normal weight pre-pregnancy percent	33.4	35.5	29.2	34.7
Obese weight pre-pregnancy percent	33.5	31.9	38.0	39.6
Appropriate weight gain during pregnancy percent	25.0	25.8	24.5	23.0

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Rural Counties Scorecard for Medicaid by Race or Ethnicity
April 2016 to March 2017

	Race or Ethnicity			
	All Races	White Race	Black Race	Hispanic Ethnicity
Medicaid-Covered Infants				
Infant Mortality Rate	6.5	6.6	**	**
Neonatal Mortality Rate	3.7 *	3.8 *	**	**
Post-Neonatal Mortality Rate	2.8	2.8	**	**
Fetal Mortality Rate	7.9	7.7	**	**
Perinatal Mortality Rate	6.5	6.7	**	**
SUIDS Mortality Rate	**	**	**	**
Preterm birth percent	10.7	10.7	11.6	9.8
Late preterm birth percent	7.6	7.6	8.7	6.8 *
Early preterm birth percent	3.1	3.0	**	**
Extreme preterm birth percent	1.8	1.7	**	**
Early term birth percent	25.9	25.8	26.0	27.0
Low birthweight birth percent	8.9	8.8	10.4	8.1
Moderately low birthweight birth percent	7.3	7.2	9.2	6.8
Very low birthweight birth percent	1.6	1.6	**	**
Medicaid-Covered Mothers				
Interpregnancy Interval less than 18 months percent	35.7	35.6	47.1	35.4
Prenatal care received in first trimester percent	73.0	73.2	71.5	72.3
Adequacy of prenatal care percent	79.3 *	79.6 *	77.6 *	77.5
Mother's smoking abstinence third trimester percent	72.0	71.6	74.3	89.7 *
Mother's alcohol abstinence third trimester percent	99.8 *	99.9 *	**	**
Mother had breast fed baby at discharge percent	58.6	58.6	56.1	69.3
Normal weight pre-pregnancy percent	31.7	31.9	24.7	25.5
Obese weight pre-pregnancy percent	36.0	36.1	35.9	39.5
Appropriate weight gain during pregnancy percent	25.2	25.2	24.4	25.7

* Meets target

** data are suppressed due to low numbers

Data were updated on 06/18/2018

Ohio Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	8.4	7.7	7.7	9.7	8.7	6.9
Neonatal Mortality Rate	**	5.1	5.2	5.8	5.6	6.2	4.0 *
Post-Neonatal Mortality Rate	**	3.3	2.5	1.9 *	4.1	2.5	2.9
Fetal Mortality Rate	**	8.0	6.0	9.4	7.8	7.3	6.7
Perinatal Mortality Rate	**	7.7	7.0	8.7	8.0	9.0	6.7
SUIDS Mortality Rate	**	1.84	1.50	**	2.11	**	**
Preterm birth percent	**	11.8	11.7	14.0	11.4	11.2	10.6
Late preterm birth percent	**	8.2	8.3	8.7	7.8	8.1	7.3
Early preterm birth percent	**	3.5	3.4	5.3	3.5	3.2	3.3
Extreme preterm birth percent	**	2.0	2.0	3.3	2.0	1.8	2.0
Early term birth percent	30.6	27.3	27.2	26.6	27.5	27.8	28.2
Low birthweight birth percent	27.8	10.4	10.7	12.0	10.5	10.8	10.5
Moderately low birthweight birth percent	**	8.6	8.9	9.0	8.5	9.1	8.8
Very low birthweight birth percent	**	1.8	1.8	2.9	2.0	1.7	1.7
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	36.7	33.6	29.4 *	35.5	36.6	35.3
Prenatal care received in first trimester percent	64.5	65.4	66.3	55.6	63.8	65.7	65.8
Adequacy of prenatal care percent	64.5	71.4	71.3	59.6	69.1	71.0	70.7
Mother's smoking abstinence third trimester percent	63.9	77.9	80.2	87.5	76.3	80.4	77.3
Mother's alcohol abstinence third trimester percent	**	99.7 *	99.7 *	99.6 *	99.8 *	99.5 *	99.6 *
Mother had breast fed baby at discharge percent	44.1	61.7	60.4	68.1	58.7	61.4	58.4
Normal weight pre-pregnancy percent	**	32.7	31.1	34.5	32.2	30.6	31.6
Obese weight pre-pregnancy percent	54.5	33.9	36.1	30.3 *	33.5	36.3	34.4
Appropriate weight gain during pregnancy percent	**	24.7	25.2	26.7	25.8	24.0	25.1

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

OEI Counties Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	9.7	8.3	8.3	10.7	10.1	7.7
Neonatal Mortality Rate	**	6.3	5.6	6.0	6.5	7.5	4.7
Post-Neonatal Mortality Rate	**	3.4	2.7	2.4	4.2	**	**
Fetal Mortality Rate	**	8.7	6.3	8.7	9.0	7.2	7.9
Perinatal Mortality Rate	**	8.7	7.9	8.3	9.6	10.1	8.3
SUIDS Mortality Rate	**	**	1.78	**	**	**	**
Preterm birth percent	**	12.1	12.3	14.3	12.5	11.4	11.8
Late preterm birth percent	**	8.2	8.4	9.1	8.4	8.0	8.1
Early preterm birth percent	**	3.9	3.8	5.2	4.2	3.4	3.8
Extreme preterm birth percent	**	2.4	2.3	3.2	2.4	2.2	2.4
Early term birth percent	32.3	27.3	27.9	27.0	27.3	29.2	28.5
Low birthweight birth percent	**	11.6	11.6	12.6	12.1	12.0	12.2
Moderately low birthweight birth percent	**	9.5	9.6	9.8	9.6	10.2	10.2
Very low birthweight birth percent	**	2.1	2.1	2.8	2.5	1.9	2.0
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	35.8	32.7	27.9 *	35.0	37.6	36.2
Prenatal care received in first trimester percent	**	60.5	64.5	53.2	59.2	60.3	61.3
Adequacy of prenatal care percent	**	65.2	68.5	56.7	62.4	64.3	64.5
Mother's smoking abstinence third trimester percent	67.7	80.5	84.6	90.0 *	82.4	84.1	82.3
Mother's alcohol abstinence third trimester percent	**	99.6 *	99.7 *	99.5 *	99.7 *	99.4 *	99.4 *
Mother had breast fed baby at discharge percent	48.3	61.6	62.7	70.9	63.3	61.4	61.4
Normal weight pre-pregnancy percent	**	33.5	30.5	35.0	32.7	29.5	31.0
Obese weight pre-pregnancy percent	53.6	33.3	36.7	29.6 *	32.1	36.6	33.7
Appropriate weight gain during pregnancy percent	**	24.4	25.1	26.8	26.0	22.8	23.3

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Non-OEI Counties Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	6.9	6.8	6.7	8.7	6.8	6.1
Neonatal Mortality Rate	**	3.7 *	4.6	5.6	4.8	4.5	3.4 *
Post-Neonatal Mortality Rate	**	3.2	2.2	**	3.9	**	**
Fetal Mortality Rate	**	7.1	5.5 *	10.5	6.7	7.5	5.4 *
Perinatal Mortality Rate	**	6.6	5.6 *	9.4	6.4	7.5	5.1 *
SUIDS Mortality Rate	**	**	1.06	**	**	**	**
Preterm birth percent	**	11.4	11.0	13.6	10.2	10.9	9.3 *
Late preterm birth percent	**	8.2	8.2	8.1	7.3	8.1	6.6 *
Early preterm birth percent	**	3.1	2.8	5.4	2.9	2.8	2.7
Extreme preterm birth percent	**	1.6	1.7	3.5	1.7	1.2 *	1.7
Early term birth percent	**	27.2	26.2	25.9	27.6	25.9	28.0
Low birthweight birth percent	**	9.0	9.4	10.8	8.8	9.1	8.8
Moderately low birthweight birth percent	**	7.6	8.0	7.7	7.4	7.7	7.5
Very low birthweight birth percent	**	1.4 *	1.4 *	3.1	1.5	1.4 *	1.4 *
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	37.9	35.1	32.2	36.0	35.5	34.4
Prenatal care received in first trimester percent	**	71.4	69.1	59.8	68.4	72.6	70.3
Adequacy of prenatal care percent	**	79.0 *	75.6	64.7	75.6	79.5 *	77.1
Mother's smoking abstinence third trimester percent	**	74.7	73.7	83.1	70.2	75.5	72.2
Mother's alcohol abstinence third trimester percent	**	**	99.8 *	99.6 *	**	**	**
Mother had breast fed baby at discharge percent	**	61.7	57.0	63.0	54.1	61.5	55.3
Normal weight pre-pregnancy percent	**	31.7	31.9	33.6	31.7	32.1	32.2
Obese weight pre-pregnancy percent	**	34.6	35.3	31.5	34.9	36.0	35.2
Appropriate weight gain during pregnancy percent	**	25.0	25.3	26.4	25.5	25.5	26.9

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Butler County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	**	8.0	**	**	**	**
Neonatal Mortality Rate	**	**	**	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**	**
Fetal Mortality Rate	**	**	**	**	**	**	**
Perinatal Mortality Rate	**	**	8.0	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**	**
Preterm birth percent	**	**	9.9	12.7	17.6	**	10.7
Late preterm birth percent	**	**	7.4	8.5	15.3	**	9.8
Early preterm birth percent	**	**	2.5	4.2	**	**	**
Extreme preterm birth percent	**	**	1.5 *	2.4	**	**	**
Early term birth percent	**	31.5	27.2	24.3	25.2	24.7	27.7
Low birthweight birth percent	**	**	9.0	8.5	9.2	**	8.9
Moderately low birthweight birth percent	**	**	7.7	6.9	7.6	**	**
Very low birthweight birth percent	**	**	1.3 *	**	**	**	**
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	44.0	32.6	27.2 *	31.9	40.4	39.0
Prenatal care received in first trimester percent	**	58.3	63.8	57.3	49.2	62.7	64.2
Adequacy of prenatal care percent	**	65.5	74.1	64.4	66.4	72.3	72.6
Mother's smoking abstinence third trimester percent	**	79.3	78.1	92.7 *	76.7	82.8	78.2
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	**	54.3	51.3	67.3	51.1	55.1	50.9
Normal weight pre-pregnancy percent	**	35.2	30.6	34.2	36.5	23.5	30.6
Obese weight pre-pregnancy percent	**	25.3 *	36.5	30.3 *	27.0 *	42.4	35.2
Appropriate weight gain during pregnancy percent	**	17.8	25.8	27.1	25.8	20.0	26.4

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Cuyahoga County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan					
	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants						
Infant Mortality Rate	**	10.1	**	**	**	**
Neonatal Mortality Rate	**	7.1	**	**	**	**
Post-Neonatal Mortality Rate	**	3.0	**	**	**	**
Fetal Mortality Rate	**	7.6	**	**	**	**
Perinatal Mortality Rate	**	9.3	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**
Preterm birth percent	18.5	13.3	15.3	16.7	16.1	13.7
Late preterm birth percent	10.8	9.3	10.6	10.4	11.1	8.3
Early preterm birth percent	7.6	4.0	4.7	6.4	5.1	5.4
Extreme preterm birth percent	5.9	2.6	3.0	4.2	3.7	3.6
Early term birth percent	26.6	28.5	29.6	30.7	28.6	26.7
Low birthweight birth percent	17.4	13.1	13.2	16.0	15.6	14.5
Moderately low birthweight birth percent	12.8	10.6	10.4	11.8	12.6	11.1
Very low birthweight birth percent	4.7	2.5	2.8	4.2	3.0	3.4
Medicaid-Covered Mothers						
Interpregnancy Interval less than 18 months percent	28.9 *	28.7 *	25.2 *	32.5	34.2	36.1
Prenatal care received in first trimester percent	60.2	63.6	53.7	56.9	55.0	58.0
Adequacy of prenatal care percent	61.9	65.8	55.6	56.3	59.1	59.5
Mother's smoking abstinence third trimester percent	85.2	88.9 *	91.6 *	87.3	90.2 *	89.8 *
Mother's alcohol abstinence third trimester percent	**	99.6 *	**	**	**	**
Mother had breast fed baby at discharge percent	63.3	65.0	72.3	62.1	66.1	66.3
Normal weight pre-pregnancy percent	35.1	28.3	32.5	33.5	30.4	30.2
Obese weight pre-pregnancy percent	31.6	40.1	32.7	34.7	31.5	35.8
Appropriate weight gain during pregnancy percent	28.8	25.9	26.4	30.8	21.9	22.5

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Franklin County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	**	6.6	7.8	7.9	**	**
Neonatal Mortality Rate	**	**	4.8	5.9	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**	**
Fetal Mortality Rate	**	**	5.0 *	9.2	6.9	**	**
Perinatal Mortality Rate	**	**	6.3	8.3	8.4	**	**
SUIDS Mortality Rate	**	**	**	**	**	**	**
Preterm birth percent	**	10.2	11.2	13.0	11.9	11.2	9.5
Late preterm birth percent	**	8.1	7.3	8.3	7.6	8.3	6.6 *
Early preterm birth percent	**	2.1	3.9	4.7	4.3	2.9	3.0
Extreme preterm birth percent	**	**	2.3	3.1	2.4	**	2.3
Early term birth percent	**	27.4	26.1	25.7	26.2	26.1	27.8
Low birthweight birth percent	**	9.5	10.8	11.6	12.2	14.1	10.4
Moderately low birthweight birth percent	**	8.5	8.6	8.8	9.9	12.8	8.5
Very low birthweight birth percent	**	**	2.2	2.8	2.3	**	1.9
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	44.2	36.5	29.2 *	36.1	42.7	42.8
Prenatal care received in first trimester percent	**	58.5	62.7	51.6	59.7	55.6	59.3
Adequacy of prenatal care percent	**	56.8	63.4	51.1	60.9	56.8	58.0
Mother's smoking abstinence third trimester percent	**	88.9 *	86.8	90.7 *	84.0	81.4	87.9 *
Mother's alcohol abstinence third trimester percent	**	**	99.7 *	**	**	**	**
Mother had breast fed baby at discharge percent	**	69.2	67.1	73.4	67.9	66.8	71.5
Normal weight pre-pregnancy percent	**	38.9	32.6	35.0	32.7	36.8	34.8
Obese weight pre-pregnancy percent	**	29.0 *	33.8	29.0 *	32.3	27.2 *	25.7 *
Appropriate weight gain during pregnancy percent	**	23.5	25.1	29.0	26.1	23.7	27.6

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Hamilton County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	**	7.6	14.2	18.0	**	**
Neonatal Mortality Rate	**	**	5.1	11.4	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**	**
Fetal Mortality Rate	**	**	6.0	**	**	**	**
Perinatal Mortality Rate	**	**	7.3	11.4	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**	**
Preterm birth percent	**	10.5	12.5	14.6	11.3	12.9	11.0
Late preterm birth percent	**	8.0	8.1	8.8	7.7	7.7	7.1
Early preterm birth percent	**	**	4.3	5.8	3.6	5.2	3.9
Extreme preterm birth percent	**	**	2.4	3.7	**	**	**
Early term birth percent	**	30.7	29.2	31.5	30.3	28.2	27.3
Low birthweight birth percent	**	13.0	12.3	14.1	11.5	13.6	13.6
Moderately low birthweight birth percent	**	12.1	10.1	11.2	9.2	11.1	11.7
Very low birthweight birth percent	**	**	2.2	2.9	2.3	**	**
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	39.2	33.4	23.4 *	31.6	38.5	42.8
Prenatal care received in first trimester percent	**	65.9	66.9	53.5	62.4	55.3	54.1
Adequacy of prenatal care percent	**	71.1	72.3	58.8	68.6	59.3	63.7
Mother's smoking abstinence third trimester percent	**	80.9	84.8	88.4 *	81.0	82.6	82.3
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	**	59.7	58.2	66.9	53.1	56.8	54.4
Normal weight pre-pregnancy percent	**	35.9	30.0	37.1	32.4	25.4	33.3
Obese weight pre-pregnancy percent	**	27.4 *	36.0	27.3 *	31.5	38.6	32.0
Appropriate weight gain during pregnancy percent	**	23.2	23.1	24.9	22.6	23.3	20.5

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Lucas County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	13.7	**	**	**	8.8	**
Neonatal Mortality Rate	**	**	**	**	**	6.2	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**	**
Fetal Mortality Rate	**	**	**	**	**	6.2	**
Perinatal Mortality Rate	**	**	**	**	**	8.8	**
SUIDS Mortality Rate	**	**	**	**	**	**	**
Preterm birth percent	**	12.4	14.5	14.4	**	10.3	13.4
Late preterm birth percent	**	9.3	9.8	8.0	**	7.5	10.6
Early preterm birth percent	**	3.0	**	6.4	**	2.8	**
Extreme preterm birth percent	**	1.8	**	3.7	**	1.6	**
Early term birth percent	**	28.2	28.9	22.4	29.0	30.8	26.1
Low birthweight birth percent	**	12.1	12.7	13.3	17.7	10.7	10.6
Moderately low birthweight birth percent	**	10.6	11.0	9.3	**	9.2	9.2
Very low birthweight birth percent	**	1.5	**	4.0	**	1.6	**
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	43.5	27.1 *	37.3	44.8	36.8	43.2
Prenatal care received in first trimester percent	**	56.0	50.8	54.2	51.0	62.9	67.0
Adequacy of prenatal care percent	**	61.0	57.3	61.6	56.0	66.9	62.1
Mother's smoking abstinence third trimester percent	**	80.6	79.1	91.5 *	67.2	84.4	87.1
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	**	46.7	50.6	67.3	40.7	58.4	53.6
Normal weight pre-pregnancy percent	**	29.8	34.1	36.8	27.1	28.8	24.8
Obese weight pre-pregnancy percent	**	37.5	34.1	30.6	42.4	38.9	42.3
Appropriate weight gain during pregnancy percent	**	25.2	24.7	26.1	20.3	22.9	21.9

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Mahoning County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan					
	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants						
Infant Mortality Rate	**	**	**	**	**	**
Neonatal Mortality Rate	**	**	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**
Fetal Mortality Rate	**	**	**	**	**	**
Perinatal Mortality Rate	**	**	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**
Preterm birth percent	**	12.8	12.9	**	**	14.2
Late preterm birth percent	**	9.9	11.4	**	**	8.9
Early preterm birth percent	**	2.9	**	**	**	5.3
Extreme preterm birth percent	**	1.4 *	**	**	**	2.7
Early term birth percent	30.0	32.3	27.3	40.5	34.0	36.5
Low birthweight birth percent	**	11.5	9.1	**	**	13.5
Moderately low birthweight birth percent	**	10.1	9.1	**	**	11.6
Very low birthweight birth percent	**	1.3 *	**	**	**	**
Medicaid-Covered Mothers						
Interpregnancy Interval less than 18 months percent	34.4	36.2	35.5	**	50.0	36.2
Prenatal care received in first trimester percent	70.0	73.4	67.7	64.9	73.9	67.8
Adequacy of prenatal care percent	70.0	77.1	70.9	**	78.3 *	72.0
Mother's smoking abstinence third trimester percent	73.3	74.5	80.8	70.3	76.1	72.8
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	58.3	55.7	72.0	59.5	72.3	47.4
Normal weight pre-pregnancy percent	35.0	34.4	32.3	40.5	34.8	30.8
Obese weight pre-pregnancy percent	23.3 *	35.1	27.7 *	32.4	30.4 *	35.4
Appropriate weight gain during pregnancy percent	26.7	25.3	30.0	**	30.4	23.3

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Montgomery County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan					
	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants						
Infant Mortality Rate	**	6.3	**	**	**	**
Neonatal Mortality Rate	**	3.7 *	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**
Fetal Mortality Rate	**	7.0	**	**	**	**
Perinatal Mortality Rate	**	7.0	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**
Preterm birth percent	12.5	13.1	17.1	9.1 *	11.5	17.6
Late preterm birth percent	**	9.3	10.2	**	**	11.8
Early preterm birth percent	**	3.7	6.9	**	**	**
Extreme preterm birth percent	**	2.2	4.4	**	**	**
Early term birth percent	28.6	27.4	26.1	20.9	30.2	24.3
Low birthweight birth percent	14.3	10.8	15.9	**	**	19.9
Moderately low birthweight birth percent	12.5	9.1	12.3	**	**	16.2
Very low birthweight birth percent	**	1.7	3.5	**	**	**
Medicaid-Covered Mothers						
Interpregnancy Interval less than 18 months percent	24.6 *	33.3	31.1	32.3	27.8 *	32.4
Prenatal care received in first trimester percent	60.2	64.2	49.2	63.2	62.8	58.8
Adequacy of prenatal care percent	71.3	71.7	57.3	70.8	63.8	62.6
Mother's smoking abstinence third trimester percent	83.8	85.4	88.4 *	76.4	89.6 *	80.6
Mother's alcohol abstinence third trimester percent	**	99.4 *	**	**	**	**
Mother had breast fed baby at discharge percent	66.7	64.2	68.8	57.3	62.5	59.3
Normal weight pre-pregnancy percent	28.7	29.4	34.8	27.5	19.8	24.4
Obese weight pre-pregnancy percent	30.6	37.7	30.3 *	28.4 *	43.8	44.3
Appropriate weight gain during pregnancy percent	17.0	25.6	24.7	24.8	19.8	22.1

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Stark County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan					
	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants						
Infant Mortality Rate	**	**	**	**	**	**
Neonatal Mortality Rate	**	**	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**
Fetal Mortality Rate	12.9	**	**	**	**	**
Perinatal Mortality Rate	**	**	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**
Preterm birth percent	11.0	10.0	17.0	**	**	10.3
Late preterm birth percent	7.7	6.4 *	11.5	**	**	8.6
Early preterm birth percent	3.3	3.7	5.5	**	**	**
Extreme preterm birth percent	2.5	1.2 *	4.3	**	**	**
Early term birth percent	22.9	26.0	23.3	25.8	23.8	28.6
Low birthweight birth percent	9.4	10.0	12.9	**	**	9.1
Moderately low birthweight birth percent	7.2	9.2	9.4	**	**	9.1
Very low birthweight birth percent	2.2	**	**	**	**	**
Medicaid-Covered Mothers						
Interpregnancy Interval less than 18 months percent	36.0	35.8	31.1	28.6 *	47.1	34.0
Prenatal care received in first trimester percent	62.0	63.9	48.7	58.3	64.7	59.2
Adequacy of prenatal care percent	66.3	70.6	54.9	63.9	64.7	65.0
Mother's smoking abstinence third trimester percent	74.6	70.8	85.2	72.1	77.8	72.5
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	58.9	59.0	59.8	48.4	69.8	51.1
Normal weight pre-pregnancy percent	31.1	33.0	38.0	28.3	23.8	31.8
Obese weight pre-pregnancy percent	36.2	33.0	25.7 *	28.3 *	41.3	30.6
Appropriate weight gain during pregnancy percent	21.8	24.5	24.3	33.3	22.2	25.3

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Summit County Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan					
	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants						
Infant Mortality Rate	8.1	14.5	**	**	**	**
Neonatal Mortality Rate	**	**	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**
Fetal Mortality Rate	10.4	**	**	**	**	**
Perinatal Mortality Rate	9.6	**	**	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**
Preterm birth percent	11.9	13.3	13.1	10.9	15.7	9.6
Late preterm birth percent	7.5	9.4	7.3	10.1	9.0	7.0
Early preterm birth percent	4.4	3.9	5.8	**	**	2.5
Extreme preterm birth percent	2.4	3.0	**	**	**	**
Early term birth percent	28.6	27.0	24.1	23.2	22.4	27.0
Low birthweight birth percent	11.3	13.6	12.2	8.6	18.7	10.0
Moderately low birthweight birth percent	8.9	10.7	9.8	7.9	14.2	8.4
Very low birthweight birth percent	2.4	2.9	**	**	**	**
Medicaid-Covered Mothers						
Interpregnancy Interval less than 18 months percent	30.2	28.5 *	27.3 *	49.2	39.5	26.4 *
Prenatal care received in first trimester percent	61.5	63.2	57.3	53.6	57.0	64.9
Adequacy of prenatal care percent	68.9	66.9	64.7	64.8	68.8	69.5
Mother's smoking abstinence third trimester percent	79.5	78.1	89.3 *	75.6	74.6	78.8
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	69.2	71.3	82.9 *	76.1	79.1	69.0
Normal weight pre-pregnancy percent	34.2	32.3	35.3	35.3	37.4	30.1
Obese weight pre-pregnancy percent	33.8	34.7	30.7	27.2 *	31.7	35.2
Appropriate weight gain during pregnancy percent	25.8	26.0	27.1	25.9	22.8	20.7

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Rural Counties Scorecard for Medicaid by Plan Coverage
April 2016 to March 2017

	Medicaid Plan						
	Aetna	Buckeye	CareSource	FFS	Molina	Paramount	United
Medicaid-Covered Infants							
Infant Mortality Rate	**	**	5.7 *	12.9	**	**	**
Neonatal Mortality Rate	**	**	3.7 *	**	**	**	**
Post-Neonatal Mortality Rate	**	**	**	**	**	**	**
Fetal Mortality Rate	**	**	4.7 *	12.8	9.0	12.5	**
Perinatal Mortality Rate	**	8.4	4.4 *	15.2	**	**	**
SUIDS Mortality Rate	**	**	**	**	**	**	**
Preterm birth percent	**	10.9	10.5	13.8	8.1 *	11.2	11.0
Late preterm birth percent	**	7.9	7.7	8.7	5.5 *	8.6	7.7
Early preterm birth percent	**	3.1	2.8	5.1	2.6	2.7	3.4
Extreme preterm birth percent	**	1.9	1.6	2.9	1.4 *	1.2 *	2.2
Early term birth percent	**	25.8	25.2	25.7	28.1	24.4	28.1
Low birthweight birth percent	**	8.0	9.2	10.1	7.5 *	8.8	10.1
Moderately low birthweight birth percent	**	6.4 *	7.8	7.8	6.1 *	7.3	8.6
Very low birthweight birth percent	**	1.7	1.4 *	2.4	1.4 *	1.5	**
Medicaid-Covered Mothers							
Interpregnancy Interval less than 18 months percent	**	38.6	35.8	31.8	36.0	34.8	35.1
Prenatal care received in first trimester percent	**	72.7	72.8	67.4	72.7	79.6 *	72.9
Adequacy of prenatal care percent	**	80.4 *	79.4 *	72.0	79.3 *	84.0 *	79.8 *
Mother's smoking abstinence third trimester percent	**	72.5	71.2	80.9	69.1	72.4	67.8
Mother's alcohol abstinence third trimester percent	**	**	**	**	**	**	**
Mother had breast fed baby at discharge percent	**	63.8	56.8	60.8	55.4	63.3	54.4
Normal weight pre-pregnancy percent	**	30.6	32.0	32.7	32.1	32.9	29.1
Obese weight pre-pregnancy percent	**	33.9	36.1	35.4	34.9	37.6	39.9
Appropriate weight gain during pregnancy percent	**	24.3	26.5	25.3	25.6	24.2	22.1

* Meets target

** data are suppressed due to low numbers

Data last updated 06/18/2018

Additional Terms and Definitions for the Medicaid Analyses

Ohio Resident – For the Ohio Department of Health Medicaid analyses, only Ohio resident mothers, or Ohio resident infant or fetal deaths are used, who also were identified as receiving Medicaid coverage.

Medicaid for this delivery or death – This report includes linked Medicaid/Vital Statistics. The Ohio Department of Medicaid (ODM) provided ODH with a list of infant and maternal IDs for Medicaid-covered infants and mothers linked to the Ohio birth certificate, and to an Ohio infant death certificate, if death occurred. Not all mother-infant dyads are able to be matched. A Medicaid ID for either mother or infant is considered a Medicaid covered mother and infant. Please Note: Due to data privacy agreements, ODM and ODH utilize slightly different data sources when reporting Vital Statistics data. ODM utilizes the Vital Statistics occurrence file which contains all births and/or deaths which occurred in Ohio during a given period of time. Conversely, ODH utilizes the Vital Statistics residence file which contains all births and/or deaths of Ohio residents as recorded on the birth or death certificate, respectively. In either case, out of state births to non-Ohio residents and out of state deaths to non-Ohio residents are excluded from these analyses.

Race – The mother's reported race from the birth certificate is used to group births, fetal deaths, the fetal portion of perinatal deaths and calculated fields based on births or births plus fetal deaths. The infant's reported race from the death certificate is used to group the infant portion of perinatal deaths, infant, neonatal, post-neonatal, and SUIDS deaths. This is consistent with our annual infant mortality report.

Mother – An Ohio resident woman physically giving birth to an infant and identified as receiving Medicaid coverage. One birth certificate from each set of plural births (twins, triplets, etc.) was selected to represent the data for that mother. Not all mothers of Medicaid-covered infants were covered by Medicaid.

Infant mortality – Infant mortality is defined as the death of live-born baby before his or her first birthday, who was identified as receiving Medicaid coverage.

Infant mortality rate – Infant mortality rates presented by the Ohio Department of Health are calculated using the number of infant deaths in a specific year as the numerator and the number of live births within that same year as the denominator, using births and deaths with identified Medicaid coverage.

Infant mortality target – Healthy People 2020 Maternal, Infant, and Child Health objective 1.3: Reduce the rate of all infant deaths (within 1 year) to 6.0 or lower, reported per 1,000 live births.

Neonatal mortality – Neonatal mortality is defined as the death of a live-born baby before he or she is 28 days of age, who was identified as receiving Medicaid coverage.

Neonatal mortality rate – Neonatal mortality rates presented by the Ohio Department of Health are calculated using the number of neonatal deaths in a specific year as the numerator and the number of live births within that same year as the denominator, reported per 1,000 live births, using births and deaths with identified Medicaid coverage.

Neonatal mortality target – Healthy People 2020 Maternal, Infant, and Child Health objective 1.4: Reduce the rate of neonatal deaths (within the first 28 days of life) to 4.1 or lower, reported per 1,000 live births.

Post-neonatal mortality – Post-neonatal mortality is defined as the death of a live-born baby when he or she is from 28 days to less than 1 year of age, who was identified as receiving Medicaid coverage.

Post-neonatal mortality rate – Post-neonatal mortality rates presented by the Ohio Department of Health are calculated using the number of post-neonatal deaths in a specific year as the numerator and the number of live births within that same year as the denominator, reported per 1,000 live births, using births and deaths with identified Medicaid coverage.

Post-neonatal mortality target – The Healthy People 2020 Maternal, Infant, and Child Health objective 1.5: Reduce the rate of post-neonatal deaths (between 28 days and 1 year) to 2.0 or lower per 1,000 live births.

Fetal mortality (stillbirth) – The death prior to the complete expulsion or extraction from its mother of a product of human conception of at least 20 weeks of gestation, which after such expulsion or extraction does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. No specific Medicaid plan information was available.

Fetal mortality rate (stillbirth rate) – Fetal mortality rates presented by the Ohio Department of Health are calculated using the number of fetal deaths in a specific year as the numerator, and the number of live births plus fetal deaths within that same year as the denominator, reported per 1,000 live births plus fetal deaths.

Fetal mortality target – The Healthy People 2020 Maternal, Infant, and Child Health objective 1.1: Reduce the rate of Fetal deaths at 20 or more weeks of gestation to 5.6 or fewer per 1,000 live births plus fetal deaths.

Perinatal mortality – Perinatal mortality is defined as the death prior to the complete expulsion or extraction from its mother of a product of human conception of at least 28 or more weeks of gestation or the death of an infant of less than 7 days of life.

Perinatal mortality rate – Perinatal mortality rates presented by the Ohio Department of Health are calculated using the number of perinatal deaths in a specific year as the numerator and the number of live births plus fetal deaths within that same year as the denominator, reported per 1,000 live births plus fetal deaths.

Perinatal mortality target – The Healthy People 2020 Maternal, Infant, and Child Health objective 1.1: Reduce the rate of perinatal deaths to 5.9 or fewer per 1,000 live births plus fetal deaths.

SUIDS – A sudden unexplained infant death is defined as the death of a live-born baby from ICD 10 causes beginning with R95, R99, or W75. For the Medicaid analyses, only those with Medicaid coverage are included.

SUIDS mortality rate – The sum of deceased infants less than one year old from SUIDS, divided by the total number of live births, reported per 1,000 live births, using births and deaths with identified Medicaid coverage.

SUIDS mortality target – Healthy People 2020 Maternal, Infant, and Child Health objective 1.9: Reduce the rate of infant deaths from sudden unexpected infant death syndromes (includes SIDS, Unknown Cause, Accidental Suffocation, and Strangulation in Bed) to 0.84 deaths per 1,000 live births.

Preterm birth – A birth of less than 37 weeks of gestation (as based upon the obstetric estimate of gestation), who was identified as receiving Medicaid coverage.

Preterm percent – The sum of these preterm infants, divided by the sum of total number of live births, reported per 100 live births with known values for weeks of gestation. Unknown term infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Preterm birth target – Healthy People 2020 Maternal, Infant, and Child Health objective 9.1: Reduce total preterm births (less than 37 weeks) to 9.4 percent or less.

Late preterm birth – A birth of 34 to less than 37 weeks of gestation), who was identified as receiving Medicaid coverage.

Late preterm percent – The sum of these late preterm infants, divided by the sum of total number of live births, reported per 100 live births with known values for weeks of gestation. Unknown term infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Late preterm target – Healthy People 2020 Maternal, Infant, and Child Health objective 9.2: Reduce late preterm or live births at 34 to 36 weeks of gestation to 6.8 percent or less.

Early preterm birth – A birth of less than 34 weeks of gestation), who was identified as receiving Medicaid coverage.

Early preterm percent – The sum of these early preterm infants, divided by the sum of total number of live births, reported per 100 live births with known values for weeks of gestation. Unknown term infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Early preterm target – Healthy People 2020 Maternal, Infant, and Child Health objective 9.3: Reduce live births at 32 to 33 weeks of gestation (early preterm) to 1.1 percent or less

Extreme preterm birth – A birth of less than 32 weeks of gestation), who was identified as receiving Medicaid coverage.

Extreme preterm percent – The sum of these extremely preterm infants, divided by the sum of total number of live births, reported per 100 live births with known values for weeks of gestation. Unknown term infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Extreme preterm target – Healthy People 2020 Maternal, Infant, and Child Health objective 9.4: Reduce live births less than or equal to 32 weeks of gestation 1.5 percent or less.

Early term birth – A birth of 37 or 38 weeks gestation), who also were identified as receiving Medicaid coverage.

Early term percent – The sum of these early term infants, divided by the sum of total number of live births, reported per 100 live births with known values for weeks of gestation. Unknown term infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Early term target – National Outcome Measure 6 – Reduce percent of early term births (37, 38 weeks gestation). Following the Healthy People 2020 model, reduce the 2010 value (20.0 percent) by 10 percent to 18 percent or lower.

Low birthweight birth – A birth of less than 2,500 grams of weight), who also were identified as receiving Medicaid coverage.

Low birthweight percent – The sum of these low birth weight infants, divided by the sum of total number of live births, reported per 100 live births with known values for birth weight. Unknown birth weight infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Low birthweight target – Healthy People 2020 Maternal, Infant, and Child Health objective 8.1 - Reduce low birth weight (LBW) of less than 2,500 grams to 7.8 percent or lower.

Moderately low birthweight birth – A birth of 1500 grams of weight to less than 2,500 grams of weight), who also were identified as receiving Medicaid coverage.

Moderately low birthweight percent – The sum of these moderately low birth weight infants, divided by the sum of total number of live births, reported per 100 live births with known values for birth weight. Unknown birth weight infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Moderately low birthweight target – National Outcome Measure 4.3: Reduce moderately low birth weight births to 6.4 percent or lower. The value is determined by subtracting the very low birthweight target from the low birthweight target).

Very low birthweight birth – A birth of less than 1,500 grams of weight), who also were identified as receiving Medicaid coverage.

Very low birthweight percent – The sum of these very low birth weight infants, divided by the sum of total number of live births, reported per 100 live births with known values for birth weight. Unknown birth weight infants are subtracted from the total of live births, using births with identified Medicaid coverage.

Very low birthweight – Healthy People 2020 Maternal, Infant, and Child Health objective 8.2: Reduce very low birth weight births to 1.4 percent or lower.

Interpregnancy interval – For each mother who was identified as receiving Medicaid coverage, having a previous live birth, and no intervening other pregnancy outcomes, this is the number of months between the end of the previous live birth and the estimated conception of the current live birth. The recommended interval is 18 months.

Interpregnancy interval percent – The sum of these mothers defined above with an interval less than 18 months, divided by the sum of the total number of mothers with a live birth, reported per 100 live births with known values for birth interval related fields, using mothers with identified Medicaid coverage.

Interpregnancy interval target – Healthy People 2020 Family Planning 5: Reduce the proportion of pregnancies conceived within 18 months of a previous birth to 29.8 percent or lower.

Prenatal care received in first trimester – A mother who reported receiving prenatal care in the first trimester and identified as receiving Medicaid coverage.

Prenatal care received in first trimester percent – The sum of these mothers defined above receiving prenatal care in the first trimester, divided by the total number of mothers with a known value for month of prenatal care, reported per 100 mothers.

Prenatal care received in first trimester target – Healthy People 2020 Maternal, Infant, and Child Health objective 10.1: Increase the percent of women who receive prenatal care beginning in the first trimester to 77.9 percent or greater.

Adequacy of prenatal care – Each mother identified as receiving Medicaid coverage where the number and timing of prenatal care visits was appropriate as determined by the Kotelchuk Index.

Adequacy of prenatal care percent – The sum of these mothers with an adequate prenatal care divided by the sum of the total number of mothers, reported per 100 mothers with known values for the fields comprising the Kotelchuk Index.

Adequacy of prenatal care target – Healthy People 2020 Maternal, Infant, and Child Health 10.2: Increase the percent of women who receive early and adequate prenatal care per the Kotelchuk Index.

Smoking abstinence in third trimester – A mother identified as receiving Medicaid coverage who did not report smoking during the third trimester.

Smoking abstinence in third trimester percent – The sum of women who reported not smoking in the third trimester, divided by the sum of women who gave live birth, reported per 100 women with known values for smoking in the third trimester. Unknown third trimester smoking women are subtracted from the total of women giving live birth.

Smoking abstinence in third trimester target – State Health Improvement Plan 73: Reduce smoking in late pregnancy to 16.3 percent or less; increase smoking abstinence in the third trimester to 87.8 percent.

Alcohol abstinence in third trimester – A mother identified as receiving Medicaid coverage who did not report drinking during the third trimester.

Alcohol abstinence in third trimester percent – The sum of mothers who did not report drinking in the third trimester, divided by the sum of women who gave live birth, reported per 100 women with known values for drinking in the third trimester. Unknown third trimester drinking women are subtracted from the total of women giving live birth.

Alcohol abstinence in third trimester target – Healthy People 2020 Maternal, Infant, and Child Health objective 11.1: Increase abstinence from alcohol among pregnant women to 98.3 percent or greater or decrease drinking among pregnant women to 1.7 percent or less.

Baby was breastfed at discharge – An infant identified as receiving Medicaid coverage who was reported as being breast fed at the time of discharge.

Baby was breastfed at discharge percent – The sum of infants reported as being breast fed at the time of discharge divided by the total number of live births with known values for being breast fed, reported per 100 live infants.

Baby was breastfed at discharge target – Healthy People 2020 Maternal, Infant, and Child Health objective 21.1: Increase the proportion of infants who are ever breastfed to 81.9 percent or more.

Normal weight pre-pregnancy – A mother identified as receiving Medicaid coverage with a Body mass index (BMI) greater than or equal to 18.5 and less than 25.0.

Normal weight pre-pregnancy percent – The sum of all mothers with a Body mass index (BMI) greater than or equal to 18.5 and less than 25.0 divided by the sum of women who gave live birth, reported per 100 women with known values for pre-pregnancy height and weight.

Normal weight pre-pregnancy target – Healthy People 2020 Maternal, Infant, and Child Health objective 16.5: Increase the percent of mothers delivering a live birth who had a healthy weight prior to pregnancy to 57.8 percent or greater.

Obese weight pre-pregnancy – A mother identified as receiving Medicaid coverage with a Body mass index (BMI) greater than or equal to 30.0.

Obese weight pre- pregnancy percent – The sum of all mothers with a Body mass index (BMI) greater than or equal to 30.0 divided by the sum of women who gave live birth, per 100 mothers with known values for pre-pregnancy height and weight, reported per 100 mothers.

Obese weight pre-pregnancy target – Healthy People 2020 Nutrition and Weight Status 9: Reduce the percent of adults who are obese to 30.5 percent or lower.

Appropriate weight gain during pregnancy – The American College of Obstetricians and Gynecologists has recommended weigh gain amounts based on a woman’s pre-pregnancy weight. A mother identified as receiving Medicaid coverage who has met this standard has had an appropriate weight gain during her pregnancy.

If Mother’s weight is	Appropriate weight gain is between
Underweight	28 – 40 pounds
Normal	25 – 35 pounds
Overweight	15 – 25 pounds
Obese	11 – 20 pounds

Appropriate weight gain during pregnancy percent – The sum of mothers who achieved an appropriate weight gain divided by the sum of women who gave live birth, reported per 100 women with known values for pre-pregnancy height and weight.

Appropriate weight gain during pregnancy target - Healthy People 2020 Maternal, Infant, and Child Health objective (developmental) – Increase the proportion of mothers who achieve a recommended weight gain during their pregnancies. No target was given, however using estimates from the MMWR Weekly / Vol. 64 / No. 43 weight gain in pregnancy, based on analysis of data from 2012 PRAMS for five states and 2013 birth certificates for 41 states, New York City, and District of Columbia, the target of 35.2 percent was used.

OEI – Ohio Equity Institute – The program supports epidemiologists in 9 selected cities or portions of cities within counties with high rates of infant mortality These city and counties are: Hamilton city in Butler county, Columbus in Franklin county, Cincinnati in Hamilton county, Toledo in Lucas county, Youngstown in Mahoning county, Dayton in Montgomery county, Canton in Stark county, and Akron, in Summit county. Selected analyses will be grouped by whether or not a county has one of the OEI programs.

Non-OEI – All other counties of Ohio.

Rural Counties – These are counties which the Centers for Medicare and Medicaid Services have classified as being rural. They include Adams, Auglaize, Brown, Carroll, Champaign, Clinton, Coshocton, Crawford, Darke, Defiance, Fayette, Fulton, Gallia, Guernsey, Hardin, Harrison, Henry, Highland, Hocking, Holmes, Jackson, Logan, Madison, Meigs, Mercer, Monroe, Morgan, Morrow, Noble, Ottawa, Paulding, Perry, Pike, Preble, Putnam, Shelby, Van Wert, Vinton, Washington, Williams, Wyandot.

Data Suppression – When the difference between the number of specific events (e.g., births with low birthweight) and the total number of events (e.g., all births) is less than 10, or the number of events is less than 10, the statistics are suppressed due to instability or confidentiality requirements.