



Ohio Department of Health Seasonal Influenza Activity Summary

MMWR Week 5

January 28th – February 3rd, 2018

Current Influenza Activity:

Current Ohio Activity Level (Geographic Spread) – Widespread

Definition: Increased ILI in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state.

During MMWR Week 5, public health surveillance data sources indicate elevated influenza-like illness (ILI) activity in outpatient settings reported by Ohio’s sentinel ILINet providers. Outpatient medical claims related to influenza-like illness increased during MMWR Week 5. The percentage of emergency department visits with patients exhibiting constitutional symptoms and fever and ILI specified ED visits increased. Reported cases of influenza-associated hospitalizations are above the seasonal threshold*. There were 1126 influenza-associated hospitalizations reported.

Ohio Weekly Influenza-associated Hospitalizations by Ohio Public Health Region

| | |
|---------------------|-------------|
| Central | 170 |
| East Central | 221 |
| Northeast | 263 |
| Northwest | 171 |
| Southeast | 50 |
| Southwest | 124 |
| West Central | 127 |
| Total | 1126 |

Weekly influenza-associated hospitalization totals are provisional and may be subject to change as more information is reported. Combined weekly totals presented in this report may not reflect the seasonal totals reported in subsequent reports.

Ohio Influenza Activity Summary Dashboard:

| Data Source | Current week value | Percent Change from last week ¹ | # of weeks ² | Trend Chart ³ |
|--|--------------------|--|-------------------------|--------------------------|
| Influenza-like Illness (ILI) Outpatient Data (ILINet Sentinel Provider Visits) | 4.62% | 15.50% | ↑ 3 | |
| Thermometer Sales (National Retail Data Monitor) | 3741 | 14.61% | ↑ 8 | |
| Fever and ILI Specified ED Visits (EpiCenter) | 4.08% | 12.71% | ↑ 2 | |
| Constitutional ED Visits (EpiCenter) | 17.27% | 7.07% | ↑ 11 | |
| Confirmed Influenza-associated Hospitalizations (Ohio Disease Reporting System) | 1126 | -10.63% | ↓ 3 | |
| Outpatient Medical Claims Data⁴ | 3.40% | 1.80% | ↑ 2 | |

¹Interpret percent changes with caution. Large variability may be exhibited in data sources with low weekly values.

²Number of weeks that the % change is increasing or decreasing.

³Black lines represent current week's data; red lines represent baseline averages

⁴Medical Claims Data provided by athenahealth®

**The seasonal threshold is 25 cases of influenza-associated hospitalizations; historical data demonstrate that once the weekly count exceeds 25 cases, the number of weekly cases thereafter will likely not decrease until after the peak of influenza activity for the season*

State, Regional, and National Data:

Ohio Surveillance Data:

- **ODH lab** has reported **813** influenza tests from specimens sent from various submitters. 2017-2018 influenza season results: **(571) A/H3N2; (67) A/pdmH1N1; (53) Influenza B;** (through 02/03/2018).
- The **National Respiratory and Enteric Virus Surveillance System (NREVSS)** has reported **43,143** influenza tests performed at participating facilities. 2017-2018 influenza season positive results: **(43) H1N1, (291) A/H3N2, (6657) Flu A Not Subtyped, and (948) Flu B** (through 02/03/2018).
- **3 pediatric influenza-associated mortalities** have been reported during the 2017-2018 season (through 02/03/2018).
- No **novel influenza A virus infections** have been reported during the 2017-2018 season (through 02/03/2018).
- Incidence of confirmed **influenza-associated hospitalizations** in 2017-2018 season = **9,738*** (through 02/03/2018).

HHS Regional Surveillance Data:** During week 4 (January 21st – January 27th, 2018), the proportion of outpatient visits for ILI in Region 5 (Ohio is in Region 5) was 5.08%, which is above the regional baseline of 1.8%. Michigan, West Virginia, Pennsylvania, Ohio, Indiana, and Kentucky reported Widespread Activity.

National Surveillance Data:** During week 4 (January 21st – January 27th), most U.S. states reported Moderate or High influenza activity, though a few are still reporting Low or Minimal activity. The proportion of outpatient visits for ILI was 7.1%, which is **above** the national baseline of 2.2%. All 10 regions reported ILI at or above their region-specific baseline level. The most frequently identified influenza virus type reported by public health laboratories was **influenza A (H3N2)**.

*Weekly influenza-associated hospitalization totals are provisional and may be subject to change as more information is reported. Combined weekly totals presented in this report may not reflect the seasonal totals reported in subsequent reports.


**National-level and regional-level data are reported one week later than Ohio state-level data



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Weekly Influenza Activity Estimates Reported by State and Territorial Epidemiologists*

Week Ending Jan 27, 2018 - Week 4



*This map indicates geographic spread and does not measure the severity of influenza activity.

2017-2018 Influenza Vaccine Components:

| A/B | Virus | Trivalent | Quadrivalent |
|-----|--|-----------|--------------|
| A | Michigan/45/2015 (H1N1)pdm09-like | X | X |
| A | Hong Kong/4801/2014 (H3N2)-like | X | X |
| B | Brisbane/60/2008-like (B/Victoria lineage) | X | X |
| B | Phuket/3073/2013-like (B/Yamagata lineage) | | X |

Influenza Virus Characterization:

CDC has antigenically or genetically characterized 1,289 influenza viruses collected during October 1, 2017 – January 27, 2018, and submitted by U.S. laboratories, including 253 influenza A(H1N1)pdm09 viruses, 668 influenza A(H3N2) viruses, and 368 influenza B viruses.

- **A (H1N1)pdm09:** Phylogenetic analysis of the HA genes from 253 A(H1N1)pdm09 viruses showed that all belonged to clade 6B.1. One hundred thirty six A(H1N1)pdm09 viruses were antigenically characterized, and all were antigenically similar (analyzed using HI with ferret antisera) to the reference 6B.1 virus A/Michigan/45/2015, representing the recommended influenza A(H1N1)pdm09 reference virus for the 2017–18 Northern Hemisphere influenza vaccines.
- **A (H3N2):** Phylogenetic analysis of the HA genes from 668 A(H3N2) viruses revealed extensive genetic diversity with multiple clades/subclades co-circulating. The HA genes of circulating viruses belonged to clade 3C.2a (n=554), subclade 3C.2a1 (n=104) or clade 3C.3a (n=10). Two hundred forty-five influenza A(H3N2) viruses were antigenically characterized, and 242 (98.8%) A(H3N2) viruses tested were well-inhibited (reacting at titers that were within fourfold of the homologous virus titer) by ferret antisera raised against A/Michigan/15/2014 (3C.2a), a cell propagated A/Hong Kong/4801/2014-like reference virus representing the A(H3N2) component of 2017–18 Northern Hemisphere influenza vaccines.


Influenza B Viruses

- **B/Victoria:** Phylogenetic analysis of 51 B/Victoria-lineage viruses indicate that all HA genes belonged to genetic clade V1A, the same genetic clade as the vaccine reference virus, B/Brisbane/60/2008. However, a number of viruses had a 6-nucleotide deletion (encoding amino acids 162 and 163) in the HA (abbreviated as V1A-2Del). Seventeen (58.6%) B/Victoria lineage viruses were well-inhibited by ferret antisera raised against cell -propagated B/Brisbane/60/2008 reference virus, representing a recommended B virus component of 2017–18 Northern Hemisphere influenza vaccines. Twelve (41.4%) B/Victoria lineage viruses reacted poorly (at titers that were 8-fold or greater reduced compared with the homologous virus titer) with ferret antisera raised against cell-propagated B/Brisbane/60/2008, and these viruses had the V1A-2Del HA.
- **B/Yamagata:** Phylogenetic analysis of 317 influenza B/Yamagata-lineage viruses indicate that the HA genes belonged to clade Y3. A total of 202 influenza B/Yamagata-lineage viruses were antigenically characterized, and all were antigenically similar to cell propagated B/Phuket/3073/2013, the reference vaccine virus representing the influenza B/Yamagata-lineage component of the 2017–18 Northern Hemisphere quadrivalent vaccines.

National activity levels and more information can be found at the following CDC pages:

- <http://www.cdc.gov/flu/weekly/usmap.htm>
- <http://www.cdc.gov/flu/>

Positive Influenza Tests (PCR), National Respiratory and Enteric Virus Surveillance System (NREVSS), 2017-2018 Influenza Season, Ohio




| Influenza-Associated Hospitalizations, Ohio 2017-2018 Season* | | | | | | | |
|--|---------------------------------------|--|------------------------------|--------------|---------------------------------------|--|------------------------------|
| County | Influenza-Associated Hospitalizations | Percent of All Influenza-Associated Hospitalizations | Rate per 100,000 Population† | County | Influenza-Associated Hospitalizations | Percent of All Influenza-Associated Hospitalizations | Rate per 100,000 Population† |
| ADAMS | 5 | 0.1% | 17.51 | LOGAN | 8 | 0.1% | 17.45 |
| ALLEN | 133 | 1.4% | 125.08 | LORAIN | 157 | 1.6% | 52.10 |
| ASHLAND | 27 | 0.3% | 50.81 | LUCAS | 378 | 3.9% | 85.56 |
| ASHTABULA | 35 | 0.4% | 34.48 | MADISON | 57 | 0.6% | 131.23 |
| ATHENS | 24 | 0.2% | 37.06 | MAHONING | 209 | 2.1% | 87.51 |
| AUGLAIZE | 31 | 0.3% | 67.47 | MARION | 85 | 0.9% | 127.82 |
| BELMONT | 36 | 0.4% | 51.14 | MEDINA | 151 | 1.6% | 87.62 |
| BROWN | 4 | 0.0% | 8.92 | MEIGS | 9 | 0.1% | 37.86 |
| BUTLER | 325 | 3.3% | 88.28 | MERCER | 29 | 0.3% | 71.05 |
| CARROLL | 43 | 0.4% | 149.12 | MIAMI | 72 | 0.7% | 70.24 |
| CHAMPAIGN | 34 | 0.3% | 84.79 | MONROE | 4 | 0.0% | 27.32 |
| CLARK | 206 | 2.1% | 148.92 | MONTGOMERY | 668 | 6.9% | 124.82 |
| CLERMONT | 180 | 1.8% | 91.20 | MORGAN | 6 | 0.1% | 39.86 |
| CLINTON | 24 | 0.2% | 57.09 | MORROW | 29 | 0.3% | 83.27 |
| COLUMBIANA | 56 | 0.6% | 51.93 | MUSKINGUM | 84 | 0.9% | 97.59 |
| COSHOCTON | 10 | 0.1% | 27.10 | NOBLE | 13 | 0.1% | 88.77 |
| CRAWFORD | 26 | 0.3% | 59.38 | OTTAWA | 31 | 0.3% | 74.83 |
| CUYAHOGA | 1447 | 14.9% | 113.04 | PAULDING | 17 | 0.2% | 86.67 |
| DARKE | 42 | 0.4% | 79.31 | PERRY | 13 | 0.1% | 36.05 |
| DEFIANCE | 23 | 0.2% | 58.92 | PICKAWAY | 38 | 0.4% | 68.23 |
| DELAWARE | 98 | 1.0% | 56.25 | PIKE | 19 | 0.2% | 66.18 |
| ERIE | 62 | 0.6% | 80.44 | PORTAGE | 111 | 1.1% | 68.77 |
| FAIRFIELD | 48 | 0.5% | 32.84 | PREBLE | 35 | 0.4% | 82.80 |
| FAYETTE | 10 | 0.1% | 34.45 | PUTNAM | 22 | 0.2% | 63.77 |
| FRANKLIN | 790 | 8.1% | 67.90 | RICHLAND | 63 | 0.6% | 50.61 |
| FULTON | 30 | 0.3% | 70.26 | ROSS | 48 | 0.5% | 61.49 |
| GALLIA | 26 | 0.3% | 84.05 | SANDUSKY | 36 | 0.4% | 59.07 |
| GEAUGA | 77 | 0.8% | 82.45 | SCIOTO | 74 | 0.8% | 93.08 |
| GREENE | 173 | 1.8% | 107.07 | SENECA | 19 | 0.2% | 33.48 |
| GUERNSEY | 41 | 0.4% | 102.28 | SHELBY | 29 | 0.3% | 58.68 |
| HAMILTON | 597 | 6.1% | 74.40 | STARK | 442 | 4.5% | 117.68 |
| HANCOCK | 42 | 0.4% | 56.16 | SUMMIT | 724 | 7.4% | 133.63 |
| HARDIN | 22 | 0.2% | 68.63 | TRUMBULL | 140 | 1.4% | 66.57 |
| HARRISON | 6 | 0.1% | 37.82 | TUSCARAWAS | 85 | 0.9% | 91.81 |
| HENRY | 25 | 0.3% | 88.61 | UNION | 29 | 0.3% | 55.45 |
| HIGHLAND | 29 | 0.3% | 66.53 | VAN WERT | 10 | 0.1% | 34.79 |
| HOCKING | 15 | 0.2% | 51.06 | VINTON | 12 | 0.1% | 89.32 |
| HOLMES | 13 | 0.1% | 30.68 | WARREN | 165 | 1.7% | 77.58 |
| HURON | 34 | 0.3% | 57.02 | WASHINGTON | 82 | 0.8% | 132.73 |
| JACKSON | 45 | 0.5% | 135.44 | WAYNE | 93 | 1.0% | 81.21 |
| JEFFERSON | 109 | 1.1% | 156.36 | WILLIAMS | 28 | 0.3% | 74.38 |
| KNOX | 21 | 0.2% | 34.47 | WOOD | 99 | 1.0% | 78.89 |
| LAKE | 120 | 1.2% | 52.16 | WYANDOT | 14 | 0.1% | 61.91 |
| LAWRENCE | 28 | 0.3% | 44.84 | UNKNOWN | 0 | 0.0% | * |
| LICKING | 129 | 1.3% | 77.48 | TOTAL | 9738 | 100% | 84.41 |

*2017-2018 Season 10/1/2017 thru 9/30/2018


† Disease rates were calculated by number of cases per 100,000 residents using 2010 census data.

Source: Ohio Disease Reporting System


Influenza-associated Hospitalizations by Public Health Region and MMWR Week, Ohio, 2017-2018 Influenza Season




Ohio Constitutional ED Visits with 5 Year Baseline Average; 2017-2018




Ohio Fever & ILI Specified ED Visits with 5 Year Baseline Average; 2017-2018




Ohio Outpatient Influenza-like Illness Network (ILINet) with 5 Year Baseline Average; 2017-2018




AthenaHealth: Ohio Influenza Related Outpatient Medical Claims 5 Year Baseline Average; 2017-2018



Ohio Thermometer Sales with 5 Year Baseline Average; 2017-2018



Ohio Confirmed Influenza-associated Hospitalizations by MMWR Week; 2017-2018 Season (n=9738)



Sources of Influenza Surveillance Data

- **National Retail Data Monitor (NRDM)-OTC Drug Purchases:** The NRDM collects over-the-counter (OTC) drug sales information from approximately 1,420 Ohio chain drug stores and grocery stores. For influenza surveillance, thermometer and adult cold relief sales are monitored on a weekly basis.
- **Emergency Department Visits (EpiCenter):** EpiCenter collects emergency department chief complaint data from 180 hospitals and urgent care facilities across Ohio in real time and classifies them into symptom and syndrome categories. Chief complaints from the constitutional syndrome category and the fever + ILI symptoms classifier are analyzed for influenza surveillance.
- **Sentinel Providers (ILINet):** Sentinel providers, through the US Influenza-like Illness Surveillance Network (ILINet), collect outpatient influenza-like illness (ILI) data. ILI is defined as a fever (≥ 100 F), **and** cough and/or sore throat without another known cause. Providers report the total number of patients seen and the number of patients with ILI by age group on a weekly basis. Sentinel providers also submit specimens for influenza testing to the ODH laboratory throughout the influenza season. There are 83 sentinel providers enrolled in Ohio for the 2016-2017 season.
- **ODH Laboratory Surveillance:** The Ohio Department of Health Laboratory reports the number of specimens that test positive for influenza each week. Generally, specimens are submitted by sentinel provider participants. A subset of the positive specimens is sent to CDC for further testing during the season.
- **Influenza-associated Hospitalizations (ODRS):** Influenza-associated hospitalizations are reported to ODH from local health departments and hospitals by direct entry into the Ohio Disease Reporting System (ODRS). Hospitalizations can be used as an indicator of the severity of illness during a particular influenza season. This condition became reportable in 2009.
- **Influenza-associated Pediatric Mortality (ODRS):** Influenza-associated pediatric mortalities are reported into ODRS by local health department and hospital staff. Pediatric deaths can be an indicator of the severity of illness during the influenza season. This condition became reportable in 2005.
- **National Respiratory and Enteric Virus Surveillance System (NREVSS):** The National Respiratory and Enteric Virus Surveillance System (NREVSS) is a laboratory-based system that monitors temporal and geographic patterns associated with the detection of respiratory syncytial virus (RSV), human parainfluenza viruses (HPIV), respiratory and enteric adenoviruses and rotavirus. There are 19 facilities in Ohio that submit data to this system.
- **athenahealth®:** athenahealth is a technology and services company for medical billing and electronic health records. Diagnosis and procedure data from primary care visits are automatically queried to produce influenza related statistics.

Ohio Public Health Regions: These counties comprise the Ohio Public Health Regions described in the figures shown on pages 1 and 5.

| Central | | East Central | | Noth East | North West | | South East | | South West | West Central |
|-----------|----------|--------------|------------|-----------|------------|----------|------------|------------|------------|--------------|
| CRAWFORD | LOGAN | ASHLAND | RICHLAND | ASHTABULA | ALLEN | MERCER | ATHENS | MONROE | ADAMS | CHAMPAIGN |
| DELAWARE | MADISON | CARROLL | STARK | CUYAHOGA | AUGLAIZE | OTTAWA | BELMONT | MORGAN | BROWN | CLARK |
| FAIRFIELD | MARION | COLUMBIANA | SUMMIT | GEAUGA | DEFIANCE | PAULDING | COSHOCTON | MUSKINGUM | BUTLER | DARKE |
| FAYETTE | MORROW | HOLMES | TRUMBULL | LAKE | ERIE | PUTNAM | GALLIA | NOBLE | CLERMONT | GREENE |
| FRANKLIN | PICKAWAY | MAHONING | TUSCARAWAS | LORAIN | FULTON | SANDUSKY | GUERNSEY | PERRY | CLINTON | MIAMI |
| HARDIN | UNION | MEDINA | WAYNE | | HANCOCK | SENECA | HARRISON | PIKE | HAMILTON | MONTGOMERY |
| KNOX | WYANDOT | PORTAGE | | | HENRY | VAN WERT | HOCKING | ROSS | HIGHLAND | PREBLE |
| LICKING | | | | | HURON | WILLIAMS | JACKSON | SCIOTO | WARREN | SHELBY |
| | | | | | LUCAS | WOOD | JEFFERSON | VINTON | | |
| | | | | | | | LAWRENCE | WASHINGTON | | |
| | | | | | | | MEIGS | | | |

If you have any further questions or comments about surveillance for seasonal influenza for the State of Ohio, please contact the Infectious Disease Informatics and Vaccine Preventable Disease Epidemiology Unit at SMED@odh.ohio.gov or call (614) 995-5599.