

Section 2: HIV Prevention Planning Regions Summary Data

Section 2 presents summary data for the 11 HIV Prevention Planning Regions in Ohio. Data presented include HIV/AIDS surveillance data on reported persons living with a diagnosis of HIV infection, reported diagnoses of HIV infection and reported cases of AIDS.

Ohio HIV Prevention Planning Groups Breakdown

Table 26. Ohio HIV prevention regional planning groups breakdown

Regional Advisory Groups		
Region 1	Region 7	Region 8C
Defiance	Butler	Adams
Fulton	Darke	Brown
Henry	Miami	Clark
Lucas	Montgomery	Clermont
Ottawa	Preble	Clinton
Sandusky	Warren	Fayette
Williams		Greene
Wood		Highland
	Region 8A	Jackson
Region 2	Allen	Lawrence
Cuyahoga	Auglaize	Madison
	Champaign	Pickaway
Region 3	Delaware	Pike
Ashtabula	Hancock	Ross
Geauga	Hardin	Scioto
Lake	Logan	
Portage	Marion	Region 8D
Summit	Mercer	Licking
Trumbull	Paulding	Muskingum
	Putnam	Guernsey
Region 4	Seneca	Belmont
Carroll	Shelby	Fairfield
Columbiana	Union	Perry
Harrison	Van Wert	Morgan
Jefferson	Wyandot	Noble
Mahoning		Monroe
Stark	Region 8B	Hocking
Tuscarawas	Ashland	Athens
	Coshocton	Washington
Region 5	Crawford	Vinton
Franklin	Erie	Meigs
	Holmes	Gallia
Region 6	Huron	
Hamilton	Knox	
	Lorain	
	Medina	
	Morrow	
	Richland	
	Wayne	

HIV Infection Data by HIV Prevention Planning Region

The following tables present HIV infection data for the nine HIV Prevention Planning Groups in Ohio. There are eight data tables for each HIV prevention planning region:

- Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics
- Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category
- Reported persons living with a diagnosis of HIV infection by year and selected characteristics, 2007-2011
- Reported persons living with a diagnosis of HIV infection by year and transmission category, 2007-2011
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by selected characteristics, 2007-2011
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by selected characteristics, 2007-2011
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, 2007-2011
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, 2007-2011

When viewing these tables, it is important to take into account the limitations of the data. These data represent confidential reports of HIV infection and AIDS diagnoses; not all persons with HIV infection. The HIV diagnosis data represent the earliest date of diagnosis reported to the HIV/AIDS Surveillance Program. The earliest date reported may not be the earliest date an individual became aware of their HIV infection. Individuals may have previously tested anonymously or were diagnosed out-of-state prior to being confidentially tested and reported to Ohio.

HIV Prevention Planning Region 1

Table 27. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 1

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	184.5	734	72%	357	71%	377	73%
Females	67.8	283	28%	147	29%	136	27%
Age at diagnosis (yr)							
<13	5.2	7	1%	3	1%	4	1%
13-14	*	1	<1%	1	<1%	-	-
15-19	13.4	8	1%	6	1%	2	<1%
20-24	53.1	33	3%	28	6%	5	1%
25-29	96.8	49	5%	35	7%	14	3%
30-34	154.2	74	7%	46	9%	28	5%
35-39	266.1	126	12%	66	13%	60	12%
40-44	359.7	182	18%	96	19%	86	17%
45-49	342.4	191	19%	77	15%	114	22%
50-54	272.4	170	17%	70	14%	100	19%
55-64	133.5	144	14%	64	13%	80	16%
65+	27.9	32	3%	12	2%	20	4%
Race/Ethnicity							
White, not Hispanic	80.5	523	51%	257	51%	266	52%
Black/African American, not Hispanic	457.8	409	40%	207	41%	202	39%
Hispanic/Latino	113.4	57	6%	23	5%	34	7%
Asian/Pacific Islander	70.4	7	1%	4	1%	3	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	21	2%	13	3%	8	2%
Race/Sex							
White, not Hispanic Males	133.2	423	42%	202	40%	221	43%
White, not Hispanic Females	30.1	100	10%	55	11%	45	9%
Black/African American, not Hispanic Males	586.3	249	24%	126	25%	123	24%
Black/African American, not Hispanic Females	341.4	160	16%	81	16%	79	15%
Hispanic/Latino Males	155.0	39	4%	15	3%	24	5%
Hispanic/Latina Females	71.8	18	2%	8	2%	10	2%
Asian/Pacific Islander Males	126.5	6	1%	4	1%	2	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	21	2%	13	3%	8	2%
Total	124.7	1,017		504		513	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 28. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 1

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	527	73%	252	72%	275	74%
Injection drug use (IDU)	32	4%	13	4%	19	5%
Male-to-male sexual contact and IDU	46	6%	21	6%	25	7%
Heterosexual contact	72	10%	36	10%	36	10%
Other/unknown	49	7%	30	9%	19	5%
Subtotal	726	100%	352	100%	374	100%
Female adult or adolescent						
Injection drug use	24	9%	10	7%	14	11%
Heterosexual contact	209	76%	107	75%	102	77%
Other/unknown	41	15%	25	18%	16	12%
Subtotal	274	100%	142	100%	132	100%
Child (<13 yrs at diagnosis)						
Perinatal	17	100%	10	100%	7	100%
Other/unknown	-	-	-	-	-	-
Subtotal	17	100%	10	100%	7	100%
Total	1,017		504		513	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 29. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 1, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	582	71%	627	71%	661	71%	690	72%	734	72%
Females	242	29%	259	29%	267	29%	273	28%	283	28%
Age at diagnosis (yr)										
<13	6	1%	7	1%	7	1%	7	1%	7	1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	5	1%	5	1%	5	1%	5	1%	8	1%
20-24	7	1%	12	1%	17	2%	24	2%	33	3%
25-29	24	3%	37	4%	41	4%	44	5%	49	5%
30-34	53	6%	60	7%	65	7%	68	7%	74	7%
35-39	97	12%	107	12%	109	12%	117	12%	126	12%
40-44	147	18%	162	18%	169	18%	174	18%	182	18%
45-49	179	22%	181	20%	185	20%	187	19%	191	19%
50-54	150	18%	154	17%	160	17%	165	17%	170	17%
55-64	124	15%	129	15%	138	15%	139	14%	144	14%
65+	31	4%	31	3%	31	3%	32	3%	32	3%
Race/Ethnicity										
White, not Hispanic	439	53%	462	52%	480	52%	500	52%	523	51%
Black/African American, not Hispanic	318	39%	350	40%	371	40%	384	40%	409	40%
Hispanic/Latino	47	6%	52	6%	54	6%	55	6%	57	6%
Asian/Pacific Islander	5	1%	7	1%	7	1%	7	1%	7	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	15	2%	15	2%	16	2%	17	2%	21	2%
Race/Sex										
White, not Hispanic Males	350	42%	370	42%	385	41%	402	42%	423	42%
White, not Hispanic Females	89	11%	92	10%	95	10%	98	10%	100	10%
Black/African American, not Hispanic Males	184	22%	204	23%	221	24%	231	24%	249	24%
Black/African American, not Hispanic Females	134	16%	146	16%	150	16%	153	16%	160	16%
Hispanic/Latino Males	31	4%	35	4%	36	4%	37	4%	39	4%
Hispanic/Latina Females	16	2%	17	2%	18	2%	18	2%	18	2%
Asian/Pacific Islander Males	5	1%	6	1%	6	1%	6	1%	6	1%
Asian/Pacific Islander Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	15	2%	15	2%	16	2%	17	2%	21	2%
Total	824		886		928		963		1,017	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 30. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 1, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	413	72%	446	72%	474	73%	496	73%	527	73%
Injection drug use (IDU)	30	5%	31	5%	31	5%	31	5%	32	4%
Male-to-male sexual contact and IDU	41	7%	43	7%	44	7%	45	7%	46	6%
Heterosexual contact	53	9%	61	10%	65	10%	68	10%	72	10%
Other/unknown	37	6%	38	6%	39	6%	42	6%	49	7%
Subtotal	574	100%	619	100%	653	100%	682	100%	726	100%
Female adult or adolescent										
Injection drug use	23	10%	24	10%	24	9%	24	9%	24	9%
Heterosexual contact	178	76%	192	77%	198	77%	203	77%	209	76%
Other/unknown	33	14%	34	14%	36	14%	37	14%	41	15%
Subtotal	234	100%	250	100%	258	100%	264	100%	274	100%
Child (<13 yrs at diagnosis)										
Perinatal	16	100%	17	100%	17	100%	17	100%	17	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	16	100%	17	100%	17	100%	17	100%	17	100%
Total	824		886		928		963		1,017	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 31. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 1, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	6.3	25	58%	11.3	45	73%	8.5	34	79%	8.0	32	82%	11.3	45	80%
Females	4.3	18	42%	4.1	17	27%	2.2	9	21%	1.7	7	18%	2.6	11	20%
Age at diagnosis (yr)															
<13	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	5%	*	2	3%	*	1	2%	*	2	5%	*	4	7%
20-24	8.0	5	12%	20.9	13	21%	11.3	7	16%	12.9	8	21%	16.1	10	18%
25-29	9.9	5	12%	19.7	10	16%	*	4	9%	*	1	3%	9.9	5	9%
30-34	20.8	10	23%	18.7	9	15%	*	3	7%	12.5	6	15%	16.7	8	14%
35-39	14.8	7	16%	31.7	15	24%	14.8	7	16%	21.1	10	26%	21.1	10	18%
40-44	15.8	8	19%	*	3	5%	*	4	9%	*	2	5%	9.9	5	9%
45-49	*	4	9%	*	4	6%	10.8	6	14%	*	3	8%	*	4	7%
50-54	*	2	5%	*	3	5%	8.0	5	12%	*	3	8%	9.6	6	11%
55-64	*	-	-	*	2	3%	5.6	6	14%	*	4	10%	*	4	7%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity															
White, not Hispanic	2.8	18	42%	3.5	23	37%	2.8	18	42%	3.5	23	59%	3.5	23	41%
Black/African American, not Hispanic	24.6	22	51%	35.8	32	52%	24.6	22	51%	15.7	14	36%	30.2	27	48%
Hispanic/Latino	*	2	5%	9.9	5	8%	*	2	5%	*	1	3%	*	2	4%
Asian/Pacific Islander	*	-	-	*	2	3%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	-	-	*	1	2%	*	1	3%	*	4	7%
Race/Sex															
White, not Hispanic Males	4.4	14	33%	6.3	20	32%	4.7	15	35%	6.3	20	51%	6.6	21	38%
White, not Hispanic Females	*	4	9%	*	3	5%	*	3	7%	*	3	8%	*	2	4%
Black/African American, not Hispanic Males	21.2	9	21%	47.1	20	32%	40.0	17	40%	23.5	10	26%	44.7	19	34%
Black/African American, not Hispanic Females	27.7	13	30%	25.6	12	19%	10.7	5	12%	*	4	10%	17.1	8	14%
Hispanic/Latino Males	*	1	2%	*	4	6%	*	1	2%	*	1	3%	*	2	4%
Hispanic/Latina Females	*	1	2%	*	1	2%	*	1	2%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	-	-	*	1	2%	*	1	3%	*	4	7%
Total	5.3	43		7.6	62		5.3	43		4.8	39		6.9	56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 32. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 1, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	21	84%	33	73%	28	82%	24	75%	32	71%
Injection drug use (IDU)	-	-	1	2%	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	-	-	2	4%	1	3%	1	3%	1	2%
Heterosexual contact	3	12%	8	18%	4	12%	3	9%	4	9%
Other/unknown	1	4%	1	2%	1	3%	4	13%	7	16%
Subtotal	25	100%	45	100%	34	100%	32	100%	45	100%
Female adult or adolescent										
Injection drug use	-	-	1	6%	-	-	-	-	-	-
Heterosexual contact	17	94%	14	88%	6	67%	6	86%	7	64%
Other/unknown	1	6%	1	6%	3	33%	1	14%	4	36%
Subtotal	18	100%	16	100%	9	100%	7	100%	11	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-
Total	43		62		43		39		56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 33. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 1, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	12	48%	23	64%	27	75%	22	79%	25	83%	854	80%
Females	13	52%	13	36%	9	25%	6	21%	5	17%	214	20%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	15	1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	2	8%	1	3%	-	-	-	-	1	3%	12	1%
20-24	2	8%	2	6%	1	3%	2	7%	1	3%	63	6%
25-29	2	8%	5	14%	3	8%	-	-	3	10%	137	13%
30-34	5	20%	6	17%	5	14%	8	29%	1	3%	220	21%
35-39	4	16%	9	25%	7	19%	5	18%	5	17%	215	20%
40-44	6	24%	5	14%	10	28%	4	14%	4	13%	160	15%
45-49	3	12%	5	14%	3	8%	5	18%	6	20%	119	11%
50-54	1	4%	2	6%	3	8%	3	11%	5	17%	69	6%
55-64	-	-	1	3%	4	11%	1	4%	4	13%	46	4%
65+	-	-	-	-	-	-	-	-	-	-	11	1%
Race/Ethnicity												
White, not Hispanic	10	40%	16	44%	13	36%	18	64%	16	53%	640	60%
Black/African American, not Hispanic	13	52%	18	50%	22	61%	7	25%	12	40%	366	34%
Hispanic/Latino	2	8%	1	3%	1	3%	3	11%	2	7%	49	5%
Asian/Pacific Islander	-	-	1	3%	-	-	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	8	1%
Race/Sex												
White, not Hispanic Males	6	24%	14	39%	10	28%	14	50%	15	50%	559	52%
White, not Hispanic Females	4	16%	2	6%	3	8%	4	14%	1	3%	81	8%
Black/African American, not Hispanic Males	5	20%	9	25%	16	44%	5	18%	8	27%	248	23%
Black/African American, not Hispanic Females	8	32%	9	25%	6	17%	2	7%	4	13%	118	11%
Hispanic/Latino Males	1	4%	-	-	1	3%	3	11%	2	7%	36	3%
Hispanic/Latina Females	1	4%	1	3%	-	-	-	-	-	-	13	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	1	3%	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	-	-	8	1%
Total	25		36		36		28		30		1,068	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 1

Table 34. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 1, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	75%	18	78%	21	78%	18	82%	18	72%	615	73%
Injection drug use (IDU)	-	-	1	4%	-	-	1	5%	1	4%	48	6%
Male-to-male sexual contact and IDU	-	-	1	4%	1	4%	-	-	1	4%	64	8%
Heterosexual contact	3	25%	2	9%	4	15%	1	5%	2	8%	55	7%
Other/unknown	-	-	1	4%	1	4%	2	9%	3	12%	62	7%
Subtotal	12	100%	23	100%	27	100%	22	100%	25	100%	844	100%
Female adult or adolescent												
Injection drug use	-	-	1	8%	-	-	-	-	-	-	31	15%
Heterosexual contact	13	100%	10	83%	6	67%	5	83%	4	80%	147	71%
Other/unknown	-	-	1	8%	3	33%	1	17%	1	20%	28	14%
Subtotal	13	100%	12	100%	9	100%	6	100%	5	100%	206	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	15	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	17%
Subtotal	-	-	1	100%	-	-	-	-	-	-	18	100%
Total	25		36		36		28		30		1,068	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 35. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 2

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	487.1	2,938	78%	1,484	77%	1,454	78%
Females	126.4	843	22%	441	23%	402	22%
Age at diagnosis (yr)							
<13	5.6	11	<1%	7	<1%	4	<1%
13-14	*	4	<1%	3	<1%	1	<1%
15-19	26.3	23	1%	17	1%	6	<1%
20-24	176.7	139	4%	109	6%	30	2%
25-29	283.8	233	6%	177	9%	56	3%
30-34	359.6	278	7%	186	10%	92	5%
35-39	403.1	290	8%	178	9%	112	6%
40-44	700.6	572	15%	297	15%	275	15%
45-49	773.5	699	18%	315	16%	384	21%
50-54	657.5	666	18%	269	14%	397	21%
55-64	415.7	710	19%	302	16%	408	22%
65+	78.7	156	4%	65	3%	91	5%
Race/Ethnicity							
White, not Hispanic	156.3	1,217	32%	606	31%	611	33%
Black/African American, not Hispanic	585.2	2,179	58%	1,121	58%	1,058	57%
Hispanic/Latino	467.2	293	8%	134	7%	159	9%
Asian/Pacific Islander	35.5	12	<1%	10	1%	2	<1%
American Indian/Alaska Native	*	3	<1%	1	<1%	2	<1%
Unknown	*	77	2%	53	3%	24	1%
Race/Sex							
White, not Hispanic Males	288.4	1,084	29%	531	28%	553	30%
White, not Hispanic Females	33.0	133	4%	75	4%	58	3%
Black/African American, not Hispanic Males	937.6	1,584	42%	813	42%	771	42%
Black/African American, not Hispanic Females	292.5	595	16%	308	16%	287	15%
Hispanic/Latino Males	636.4	197	5%	91	5%	106	6%
Hispanic/Latina Females	302.3	96	3%	43	2%	53	3%
Asian/Pacific Islander Males	55.0	9	<1%	7	<1%	2	<1%
Asian/Pacific Islander Females	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	77	2%	53	3%	24	1%
Total	297.6	3,781		1,925		1,856	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 36. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 2

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,937	67%	943	64%	994	69%
Injection drug use (IDU)	160	5%	55	4%	105	7%
Male-to-male sexual contact and IDU	137	5%	42	3%	95	7%
Heterosexual contact	212	7%	104	7%	108	7%
Other/unknown	466	16%	324	22%	142	10%
Subtotal	2,912	100%	1,468	100%	1,444	100%
Female adult or adolescent						
Injection drug use	113	14%	35	8%	78	20%
Heterosexual contact	496	61%	244	57%	252	65%
Other/unknown	210	26%	150	35%	60	15%
Subtotal	819	100%	429	100%	390	100%
Child (<13 yrs at diagnosis)						
Perinatal	44	88%	23	82%	21	95%
Other/unknown	6	12%	5	18%	1	5%
Subtotal	50	100%	28	100%	22	100%
Total	3,781		1,925		1,856	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 37. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 2, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,327	78%	2,492	77%	2,639	77%	2,776	78%	2,938	78%
Females	675	22%	727	23%	774	23%	803	22%	843	22%
Age at diagnosis (yr)										
<13	9	<1%	9	<1%	11	<1%	11	<1%	11	<1%
13-14	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
15-19	16	1%	16	<1%	16	<1%	19	1%	23	1%
20-24	29	1%	43	1%	63	2%	100	3%	139	4%
25-29	123	4%	157	5%	184	5%	206	6%	233	6%
30-34	162	5%	195	6%	225	7%	245	7%	278	7%
35-39	215	7%	242	8%	259	8%	278	8%	290	8%
40-44	463	15%	497	15%	533	16%	548	15%	572	15%
45-49	621	21%	643	20%	665	19%	679	19%	699	18%
50-54	584	19%	613	19%	628	18%	644	18%	666	18%
55-64	631	21%	650	20%	672	20%	691	19%	710	19%
65+	146	5%	150	5%	153	4%	154	4%	156	4%
Race/Ethnicity										
White, not Hispanic	998	33%	1,067	33%	1,126	33%	1,166	33%	1,217	32%
Black/African American, not Hispanic	1,685	56%	1,813	56%	1,934	57%	2,044	57%	2,179	58%
Hispanic/Latino	240	8%	259	8%	268	8%	280	8%	293	8%
Asian/Pacific Islander	10	<1%	10	<1%	11	<1%	12	<1%	12	<1%
American Indian/Alaska Native	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
Unknown	66	2%	67	2%	71	2%	74	2%	77	2%
Race/Sex										
White, not Hispanic Males	886	30%	948	29%	1,002	29%	1,039	29%	1,084	29%
White, not Hispanic Females	112	4%	119	4%	124	4%	127	4%	133	4%
Black/African American, not Hispanic Males	1,217	41%	1,307	41%	1,392	41%	1,479	41%	1,584	42%
Black/African American, not Hispanic Females	468	16%	506	16%	542	16%	565	16%	595	16%
Hispanic/Latino Males	160	5%	172	5%	178	5%	187	5%	197	5%
Hispanic/Latina Females	80	3%	87	3%	90	3%	93	3%	96	3%
Asian/Pacific Islander Males	8	<1%	8	<1%	8	<1%	9	<1%	9	<1%
Asian/Pacific Islander Females	2	<1%	2	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	66	2%	67	2%	71	2%	74	2%	77	2%
Total	3,002		3,219		3,413		3,579		3,781	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 38. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 2, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,518	66%	1,636	66%	1,736	66%	1,826	66%	1,937	67%
Injection drug use (IDU)	148	6%	151	6%	157	6%	157	6%	160	5%
Male-to-male sexual contact and IDU	124	5%	132	5%	136	5%	136	5%	137	5%
Heterosexual contact	173	8%	188	8%	203	8%	206	7%	212	7%
Other/unknown	340	15%	361	15%	381	15%	425	15%	466	16%
Subtotal	2,303	100%	2,468	100%	2,613	100%	2,750	100%	2,912	100%
Female adult or adolescent										
Injection drug use	101	15%	104	15%	107	14%	111	14%	113	14%
Heterosexual contact	410	63%	448	64%	475	63%	481	62%	496	61%
Other/unknown	141	22%	151	21%	168	22%	187	24%	210	26%
Subtotal	652	100%	703	100%	750	100%	779	100%	819	100%
Child (<13 yrs at diagnosis)										
Perinatal	41	87%	42	88%	44	88%	44	88%	44	88%
Other/unknown	6	13%	6	13%	6	12%	6	12%	6	12%
Subtotal	47	100%	48	100%	50	100%	50	100%	50	100%
Total	3,002		3,219		3,413		3,579		3,781	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 39. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 2, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	28.5	172	82%	29.7	179	76%	25.9	156	76%	23.2	140	83%	27.4	165	80%
Females	5.5	37	18%	8.4	56	24%	7.5	50	24%	4.3	29	17%	6.0	40	20%
Age at diagnosis (yr)															
<13	*	-	-	*	1	<1%	*	2	1%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
15-19	13.7	12	6%	12.6	11	5%	9.2	8	4%	21.7	19	11%	13.7	12	6%
20-24	47.0	37	18%	41.9	33	14%	35.6	28	14%	40.7	32	19%	49.6	39	19%
25-29	24.4	20	10%	42.6	35	15%	41.4	34	17%	25.6	21	12%	35.3	29	14%
30-34	27.2	21	10%	36.2	28	12%	20.7	16	8%	29.8	23	14%	36.2	28	14%
35-39	45.9	33	16%	45.9	33	14%	48.7	35	17%	23.6	17	10%	13.9	10	5%
40-44	35.5	29	14%	31.8	26	11%	29.4	24	12%	17.1	14	8%	34.3	28	14%
45-49	25.5	23	11%	32.1	29	12%	27.7	25	12%	18.8	17	10%	28.8	26	13%
50-54	16.8	17	8%	18.8	19	8%	16.8	17	8%	11.8	12	7%	14.8	15	7%
55-64	5.9	10	5%	10.5	18	8%	7.6	13	6%	7.6	13	8%	9.4	16	8%
65+	3.5	7	3%	*	2	1%	*	4	2%	*	-	-	*	2	1%
Race/Ethnicity															
White, not Hispanic	8.1	63	30%	9.5	74	31%	8.2	64	31%	5.4	42	25%	6.7	52	25%
Black/African American, not Hispanic	34.9	130	62%	37.1	138	59%	34.4	128	62%	29.8	111	66%	36.8	137	67%
Hispanic/Latino	22.3	14	7%	33.5	21	9%	14.4	9	4%	19.1	12	7%	20.7	13	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	1	<1%	*	1	1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	2	1%	*	4	2%	*	3	2%	*	3	1%
Race/Sex															
White, not Hispanic Males	15.4	58	28%	17.8	67	29%	15.4	58	28%	10.4	39	23%	12.2	46	22%
White, not Hispanic Females	1.2	5	2%	1.7	7	3%	1.5	6	3%	*	3	2%	1.5	6	3%
Black/African American, not Hispanic Males	60.4	102	49%	58.0	98	42%	53.3	90	44%	52.1	88	52%	63.3	107	52%
Black/African American, not Hispanic Females	13.8	28	13%	19.7	40	17%	18.7	38	18%	11.3	23	14%	14.7	30	15%
Hispanic/Latino Males	32.3	10	5%	42.0	13	6%	19.4	6	3%	29.1	9	5%	32.3	10	5%
Hispanic/Latina Females	*	4	2%	25.2	8	3%	*	3	1%	*	3	2%	*	3	1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	2	1%	*	4	2%	*	3	2%	*	3	1%
Total	16.5	209		18.5	235		16.2	206		13.3	169		16.1	205	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 40. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 2, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	127	74%	127	71%	99	64%	92	66%	113	68%
Injection drug use (IDU)	4	2%	3	2%	7	5%	-	-	3	2%
Male-to-male sexual contact and IDU	4	2%	7	4%	5	3%	-	-	1	1%
Heterosexual contact	14	8%	16	9%	15	10%	3	2%	6	4%
Other/unknown	23	13%	26	15%	28	18%	45	32%	42	25%
Subtotal	172	100%	179	100%	154	100%	140	100%	165	100%
Female adult or adolescent										
Injection drug use	3	8%	4	7%	3	6%	4	14%	2	5%
Heterosexual contact	25	68%	41	75%	27	54%	6	21%	15	38%
Other/unknown	9	24%	10	18%	20	40%	19	66%	23	58%
Subtotal	37	100%	55	100%	50	100%	29	100%	40	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	1	100%	2	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	2	100%	-	-	-	-
Total	209		235		206		169		205	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 41. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 2, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	71	79%	75	76%	76	75%	66	73%	66	80%	3,353	82%
Females	19	21%	24	24%	25	25%	24	27%	17	20%	728	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	41	1%
13-14	1	1%	-	-	-	-	-	-	-	-	3	<1%
15-19	4	4%	-	-	1	1%	3	3%	2	2%	30	1%
20-24	10	11%	9	9%	5	5%	11	12%	6	7%	193	5%
25-29	4	4%	12	12%	6	6%	10	11%	8	10%	512	13%
30-34	7	8%	14	14%	13	13%	9	10%	9	11%	817	20%
35-39	15	17%	14	14%	21	21%	11	12%	5	6%	900	22%
40-44	15	17%	18	18%	15	15%	12	13%	15	18%	655	16%
45-49	15	17%	15	15%	18	18%	17	19%	23	28%	448	11%
50-54	11	12%	6	6%	9	9%	11	12%	5	6%	225	6%
55-64	2	2%	10	10%	12	12%	6	7%	8	10%	203	5%
65+	6	7%	1	1%	1	1%	-	-	2	2%	54	1%
Race/Ethnicity												
White, not Hispanic	33	37%	27	27%	32	32%	24	27%	19	23%	1,499	37%
Black/African American, not Hispanic	50	56%	54	55%	60	59%	58	64%	60	72%	2,241	55%
Hispanic/Latino	7	8%	16	16%	7	7%	7	8%	3	4%	302	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	5	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	5	<1%
Unknown	-	-	2	2%	2	2%	1	1%	1	1%	29	1%
Race/Sex												
White, not Hispanic Males	28	31%	25	25%	29	29%	20	22%	18	22%	1,389	34%
White, not Hispanic Females	5	6%	2	2%	3	3%	4	4%	1	1%	110	3%
Black/African American, not Hispanic Males	38	42%	41	41%	42	42%	41	46%	44	53%	1,720	42%
Black/African American, not Hispanic Females	12	13%	13	13%	18	18%	17	19%	16	19%	521	13%
Hispanic/Latino Males	5	6%	8	8%	4	4%	4	4%	3	4%	214	5%
Hispanic/Latina Females	2	2%	8	8%	3	3%	3	3%	-	-	88	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	5	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	-	-	2	2%	2	2%	1	1%	1	1%	29	1%
Total	90		99		101		90		83		4,081	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 2

Table 42. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 2, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	50	70%	49	65%	45	59%	42	64%	39	60%	2,272	68%
Injection drug use (IDU)	3	4%	2	3%	6	8%	2	3%	1	2%	339	10%
Male-to-male sexual contact and IDU	2	3%	6	8%	6	8%	3	5%	1	2%	256	8%
Heterosexual contact	8	11%	11	15%	11	14%	2	3%	3	5%	183	5%
Other/unknown	8	11%	7	9%	8	11%	17	26%	21	32%	283	8%
Subtotal	71	100%	75	100%	76	100%	66	100%	65	100%	3,333	100%
Female adult or adolescent												
Injection drug use	5	28%	2	8%	2	8%	5	23%	1	6%	179	26%
Heterosexual contact	11	61%	20	83%	19	76%	7	32%	8	47%	406	58%
Other/unknown	2	11%	2	8%	4	16%	10	45%	8	47%	113	16%
Subtotal	18	100%	24	100%	25	100%	22	100%	17	100%	698	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	2	100%	1	100%	45	90%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	10%
Subtotal	1	100%	-	-	-	-	2	100%	1	100%	50	100%
Total	90		99		101		90		83		4,081	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 43. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 3

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	139.4	907	79%	455	77%	452	82%
Females	34.6	237	21%	137	23%	100	18%
Age at diagnosis (yr)							
<13	3.9	8	1%	8	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	9.7	9	1%	8	1%	1	<1%
20-24	37.2	32	3%	24	4%	8	1%
25-29	90.1	68	6%	53	9%	15	3%
30-34	118.3	88	8%	71	12%	17	3%
35-39	144.2	108	9%	59	10%	49	9%
40-44	168.1	150	13%	73	12%	77	14%
45-49	211.0	207	18%	92	16%	115	21%
50-54	193.9	210	18%	98	17%	112	20%
55-64	116.1	218	19%	91	15%	127	23%
65+	22.3	46	4%	15	3%	31	6%
Race/Ethnicity							
White, not Hispanic	58.3	670	59%	339	57%	331	60%
Black/African American, not Hispanic	350.4	401	35%	217	37%	184	33%
Hispanic/Latino	150.8	41	4%	17	3%	24	4%
Asian/Pacific Islander	35.7	7	1%	5	1%	2	<1%
American Indian/Alaska Native	*	4	<1%	3	1%	1	<1%
Unknown	*	21	2%	11	2%	10	2%
Race/Sex							
White, not Hispanic Males	101.4	569	50%	284	48%	285	52%
White, not Hispanic Females	17.1	101	9%	55	9%	46	8%
Black/African American, not Hispanic Males	524.3	285	25%	145	24%	140	25%
Black/African American, not Hispanic Females	193.1	116	10%	72	12%	44	8%
Hispanic/Latino Males	220.5	31	3%	13	2%	18	3%
Hispanic/Latina Females	76.2	10	1%	4	1%	6	1%
Asian/Pacific Islander Males	*	4	<1%	4	1%	-	-
Asian/Pacific Islander Females	*	3	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	2	<1%	2	<1%	-	-
Unknown	*	21	2%	11	2%	10	2%
Total	85.7	1,144		592		552	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 44. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 3

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	584	65%	265	59%	319	71%
Injection drug use (IDU)	42	5%	11	2%	31	7%
Male-to-male sexual contact and IDU	31	3%	10	2%	21	5%
Heterosexual contact	71	8%	31	7%	40	9%
Other/unknown	168	19%	129	29%	39	9%
Subtotal	896	100%	446	100%	450	100%
Female adult or adolescent						
Injection drug use	22	10%	10	8%	12	12%
Heterosexual contact	131	57%	68	52%	63	64%
Other/unknown	77	33%	53	40%	24	24%
Subtotal	230	100%	131	100%	99	100%
Child (<13 yrs at diagnosis)						
Perinatal	16	89%	14	93%	2	67%
Other/unknown	2	11%	1	7%	1	33%
Subtotal	18	100%	15	100%	3	100%
Total	1,144		592		552	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 45. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 3, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	718	79%	768	79%	819	79%	851	79%	907	79%
Females	187	21%	199	21%	212	21%	225	21%	237	21%
Age at diagnosis (yr)										
<13	5	1%	6	1%	6	1%	6	1%	8	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	6	1%	6	1%	7	1%	7	1%	9	1%
20-24	7	1%	12	1%	17	2%	20	2%	32	3%
25-29	24	3%	35	4%	52	5%	56	5%	68	6%
30-34	53	6%	64	7%	73	7%	83	8%	88	8%
35-39	85	9%	93	10%	100	10%	102	9%	108	9%
40-44	127	14%	130	13%	135	13%	140	13%	150	13%
45-49	177	20%	184	19%	194	19%	201	19%	207	18%
50-54	188	21%	193	20%	199	19%	205	19%	210	18%
55-64	193	21%	203	21%	207	20%	212	20%	218	19%
65+	40	4%	41	4%	41	4%	44	4%	46	4%
Race/Ethnicity										
White, not Hispanic	539	60%	569	59%	606	59%	631	59%	670	59%
Black/African American, not Hispanic	311	34%	338	35%	361	35%	376	35%	401	35%
Hispanic/Latino	32	4%	36	4%	38	4%	40	4%	41	4%
Asian/Pacific Islander	6	1%	6	1%	6	1%	7	1%	7	1%
American Indian/Alaska Native	3	<1%	3	<1%	3	<1%	4	<1%	4	<1%
Unknown	14	2%	15	2%	17	2%	18	2%	21	2%
Race/Sex										
White, not Hispanic Males	463	51%	489	51%	517	50%	536	50%	569	50%
White, not Hispanic Females	76	8%	80	8%	89	9%	95	9%	101	9%
Black/African American, not Hispanic Males	215	24%	235	24%	255	25%	266	25%	285	25%
Black/African American, not Hispanic Females	96	11%	103	11%	106	10%	110	10%	116	10%
Hispanic/Latino Males	25	3%	28	3%	29	3%	30	3%	31	3%
Hispanic/Latina Females	7	1%	8	1%	9	1%	10	1%	10	1%
Asian/Pacific Islander Males	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Asian/Pacific Islander Females	2	<1%	2	<1%	2	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
Unknown	14	2%	15	2%	17	2%	18	2%	21	2%
Total	905		967		1,031		1,076		1,144	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 46. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 3, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	464	65%	504	66%	533	66%	551	65%	584	65%
Injection drug use (IDU)	36	5%	38	5%	40	5%	41	5%	42	5%
Male-to-male sexual contact and IDU	29	4%	30	4%	30	4%	31	4%	31	3%
Heterosexual contact	58	8%	61	8%	68	8%	68	8%	71	8%
Other/unknown	122	17%	126	17%	139	17%	151	18%	168	19%
Subtotal	709	100%	759	100%	810	100%	842	100%	896	100%
Female adult or adolescent										
Injection drug use	19	10%	21	11%	22	11%	22	10%	22	10%
Heterosexual contact	113	62%	116	60%	124	60%	129	59%	131	57%
Other/unknown	49	27%	55	29%	59	29%	67	31%	77	33%
Subtotal	181	100%	192	100%	205	100%	218	100%	230	100%
Child (<13 yrs at diagnosis)										
Perinatal	14	93%	14	88%	14	88%	14	88%	16	89%
Other/unknown	1	7%	2	13%	2	13%	2	13%	2	11%
Subtotal	15	100%	16	100%	16	100%	16	100%	18	100%
Total	905		967		1,031		1,076		1,144	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 47. Reported diagnoses of HIV infection, by year of diagnosis and selected characteristics, Region 3, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	8.6	56	77%	7.8	51	81%	8.3	54	79%	5.1	33	70%	9.1	59	83%
Females	2.5	17	23%	1.8	12	19%	2.0	14	21%	2.0	14	30%	1.8	12	17%
Age at diagnosis (yr)															
<13	*	1	1%	*	1	2%	*	-	-	*	-	-	*	2	3%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	3	4%	*	4	6%	*	3	4%	*	1	2%	*	4	6%
20-24	7.0	6	8%	12.8	11	17%	19.7	17	25%	*	4	9%	16.3	14	20%
25-29	15.9	12	16%	14.6	11	17%	13.2	10	15%	6.6	5	11%	15.9	12	17%
30-34	6.7	5	7%	8.1	6	10%	6.7	5	7%	12.1	9	19%	*	4	6%
35-39	12.0	9	12%	*	3	5%	8.0	6	9%	*	3	6%	13.3	10	14%
40-44	19.1	17	23%	*	4	6%	11.2	10	15%	*	4	9%	6.7	6	8%
45-49	9.2	9	12%	9.2	9	14%	7.1	7	10%	9.2	9	19%	5.1	5	7%
50-54	7.4	8	11%	6.5	7	11%	4.6	5	7%	5.5	6	13%	7.4	8	11%
55-64	*	3	4%	3.7	7	11%	2.7	5	7%	*	3	6%	3.2	6	8%
65+	*	-	-	*	-	-	*	-	-	*	3	6%	*	-	-
Race/Ethnicity															
White, not Hispanic	2.7	31	42%	2.9	33	52%	3.4	39	57%	2.3	26	55%	3.6	41	58%
Black/African American, not Hispanic	31.5	36	49%	22.7	26	41%	21.8	25	37%	13.1	15	32%	22.7	26	37%
Hispanic/Latino	*	4	5%	*	3	5%	*	2	3%	*	2	4%	*	1	1%
Asian/Pacific Islander	*	1	1%	*	-	-	*	-	-	*	2	4%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	2%	*	-	-
Unknown	*	1	1%	*	1	2%	*	2	3%	*	1	2%	*	3	4%
Race/Sex															
White, not Hispanic Males	4.6	26	36%	5.2	29	46%	5.3	30	44%	3.4	19	40%	6.2	35	49%
White, not Hispanic Females	0.8	5	7%	*	4	6%	1.5	9	13%	1.2	7	15%	1.0	6	8%
Black/African American, not Hispanic Males	46.0	25	34%	35.0	19	30%	38.6	21	31%	20.2	11	23%	36.8	20	28%
Black/African American, not Hispanic Females	18.3	11	15%	11.7	7	11%	*	4	6%	*	4	9%	10.0	6	8%
Hispanic/Latino Males	*	4	5%	*	2	3%	*	1	1%	*	1	2%	*	1	1%
Hispanic/Latina Females	*	-	-	*	1	2%	*	1	1%	*	1	2%	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	2%	*	-	-
Asian/Pacific Islander Females	*	1	1%	*	-	-	*	-	-	*	1	2%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	1	2%	*	-	-
Unknown	*	1	1%	*	1	2%	*	2	3%	*	1	2%	*	3	4%
Total	5.5	73		4.7	63		5.1	68		3.5	47		5.3	71	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 48. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 3, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	33	60%	44	86%	30	56%	18	55%	33	58%
Injection drug use (IDU)	2	4%	1	2%	2	4%	1	3%	1	2%
Male-to-male sexual contact and IDU	1	2%	-	-	-	-	1	3%	-	-
Heterosexual contact	9	16%	3	6%	8	15%	1	3%	3	5%
Other/unknown	10	18%	3	6%	14	26%	12	36%	20	35%
Subtotal	55	100%	51	100%	54	100%	33	100%	57	100%
Female adult or adolescent										
Injection drug use	1	6%	2	18%	1	7%	-	-	-	-
Heterosexual contact	13	76%	3	27%	9	64%	5	36%	2	17%
Other/unknown	3	18%	6	55%	4	29%	9	64%	10	83%
Subtotal	17	100%	11	100%	14	100%	14	100%	12	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	2	100%
Other/unknown	-	-	1	100%	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-	2	100%
Total	73		63		68		47		71	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 49. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 3, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	26	79%	23	79%	26	84%	14	74%	21	78%	1,040	86%
Females	7	21%	6	21%	5	16%	5	26%	6	22%	164	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	3	<1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	1	3%	-	-	-	-	2	11%	-	-	9	1%
20-24	-	-	2	7%	2	6%	1	5%	3	11%	51	4%
25-29	2	6%	1	3%	4	13%	2	11%	4	15%	151	13%
30-34	5	15%	3	10%	5	16%	3	16%	3	11%	247	21%
35-39	7	21%	2	7%	4	13%	-	-	1	4%	217	18%
40-44	10	30%	5	17%	6	19%	1	5%	5	19%	227	19%
45-49	2	6%	7	24%	4	13%	4	21%	3	11%	135	11%
50-54	4	12%	8	28%	3	10%	4	21%	6	22%	86	7%
55-64	2	6%	-	-	3	10%	1	5%	2	7%	63	5%
65+	-	-	1	3%	-	-	1	5%	-	-	13	1%
Race/Ethnicity												
White, not Hispanic	17	52%	13	45%	13	42%	12	63%	16	59%	794	66%
Black/African American, not Hispanic	12	36%	14	48%	13	42%	4	21%	9	33%	365	30%
Hispanic/Latino	4	12%	1	3%	2	6%	1	5%	-	-	30	2%
Asian/Pacific Islander	-	-	-	-	1	3%	1	5%	-	-	3	<1%
American Indian/Alaska Native	-	-	1	3%	-	-	-	-	-	-	2	<1%
Unknown	-	-	-	-	2	6%	1	5%	2	7%	10	1%
Race/Sex												
White, not Hispanic Males	16	48%	11	38%	13	42%	9	47%	13	48%	714	59%
White, not Hispanic Females	1	3%	2	7%	-	-	3	16%	3	11%	80	7%
Black/African American, not Hispanic Males	6	18%	11	38%	10	32%	3	16%	6	22%	293	24%
Black/African American, not Hispanic Females	6	18%	3	10%	3	10%	1	5%	3	11%	72	6%
Hispanic/Latino Males	4	12%	-	-	1	3%	-	-	-	-	23	2%
Hispanic/Latina Females	-	-	1	3%	1	3%	1	5%	-	-	7	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	1	5%	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	1	3%	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	1	3%	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	2	6%	1	5%	2	7%	10	1%
Total	33		29		31		19		27		1,204	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 3

Table 50. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 3, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	15	58%	13	57%	17	65%	10	71%	16	76%	732	71%
Injection drug use (IDU)	3	12%	1	4%	3	12%	1	7%	1	5%	76	7%
Male-to-male sexual contact and IDU	1	4%	1	4%	-	-	-	-	-	-	65	6%
Heterosexual contact	3	12%	3	13%	4	15%	1	7%	3	14%	55	5%
Other/unknown	4	15%	5	22%	2	8%	2	14%	1	5%	109	11%
Subtotal	26	100%	23	100%	26	100%	14	100%	21	100%	1,037	100%
Female adult or adolescent												
Injection drug use	2	29%	3	50%	-	-	-	-	-	-	23	14%
Heterosexual contact	4	57%	2	33%	3	60%	2	40%	1	17%	94	58%
Other/unknown	1	14%	1	17%	2	40%	3	60%	5	83%	44	27%
Subtotal	7	100%	6	100%	5	100%	5	100%	6	100%	161	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	33		29		31		19		27		1,204	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 51. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 4

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	131.3	593	72%	321	73%	272	71%
Females	47.6	226	28%	117	27%	109	29%
Age at diagnosis (yr)							
<13	*	1	<1%	1	<1%	-	-
13-14	*	1	<1%	-	-	1	<1%
15-19	8.1	5	1%	4	1%	1	<1%
20-24	78.7	43	5%	36	8%	7	2%
25-29	111.5	56	7%	43	10%	13	3%
30-34	91.2	47	6%	31	7%	16	4%
35-39	130.8	68	8%	37	8%	31	8%
40-44	218.3	128	16%	68	16%	60	16%
45-49	216.9	141	17%	67	15%	74	19%
50-54	201.0	150	18%	65	15%	85	22%
55-64	106.6	143	17%	67	15%	76	20%
65+	22.8	36	4%	19	4%	17	4%
Race/Ethnicity							
White, not Hispanic	54.7	442	54%	237	54%	205	54%
Black/African American, not Hispanic	413.4	299	37%	148	34%	151	40%
Hispanic/Latino	195.3	43	5%	25	6%	18	5%
Asian/Pacific Islander	*	2	<1%	2	<1%	-	-
American Indian/Alaska Native	*	3	<1%	2	<1%	1	<1%
Unknown	*	30	4%	24	5%	6	2%
Race/Sex							
White, not Hispanic Males	89.4	351	43%	186	42%	165	43%
White, not Hispanic Females	21.9	91	11%	51	12%	40	10%
Black/African American, not Hispanic Males	535.6	190	23%	101	23%	89	23%
Black/African American, not Hispanic Females	295.8	109	13%	47	11%	62	16%
Hispanic/Latino Males	215.9	26	3%	14	3%	12	3%
Hispanic/Latina Females	170.4	17	2%	11	3%	6	2%
Asian/Pacific Islander Males	*	1	<1%	1	<1%	-	-
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	30	4%	24	5%	6	2%
Total	88.5	819		438		381	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 52. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 4

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	361	61%	185	58%	176	65%
Injection drug use (IDU)	21	4%	8	3%	13	5%
Male-to-male sexual contact and IDU	13	2%	5	2%	8	3%
Heterosexual contact	57	10%	24	8%	33	12%
Other/unknown	137	23%	97	30%	40	15%
Subtotal	589	100%	319	100%	270	100%
Female adult or adolescent						
Injection drug use	21	10%	6	5%	15	14%
Heterosexual contact	125	58%	57	52%	68	64%
Other/unknown	70	32%	47	43%	23	22%
Subtotal	216	100%	110	100%	106	100%
Child (<13 yrs at diagnosis)						
Perinatal	11	79%	7	78%	4	80%
Other/unknown	3	21%	2	22%	1	20%
Subtotal	14	100%	9	100%	5	100%
Total	819		438		381	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 53. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 4, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	434	71%	465	71%	498	71%	547	72%	593	72%
Females	179	29%	192	29%	205	29%	211	28%	226	28%
Age at diagnosis (yr)										
<13	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	4	1%	4	1%	4	1%	5	1%	5	1%
20-24	5	1%	10	2%	15	2%	26	3%	43	5%
25-29	19	3%	27	4%	41	6%	44	6%	56	7%
30-34	28	5%	31	5%	36	5%	43	6%	47	6%
35-39	52	8%	56	9%	62	9%	63	8%	68	8%
40-44	99	16%	106	16%	112	16%	123	16%	128	16%
45-49	121	20%	126	19%	128	18%	137	18%	141	17%
50-54	124	20%	134	20%	139	20%	145	19%	150	18%
55-64	126	21%	128	19%	130	18%	136	18%	143	17%
65+	33	5%	33	5%	34	5%	34	4%	36	4%
Race/Ethnicity										
White, not Hispanic	331	54%	353	54%	380	54%	410	54%	442	54%
Black/African American, not Hispanic	225	37%	244	37%	258	37%	277	37%	299	37%
Hispanic/Latino	35	6%	37	6%	39	6%	41	5%	43	5%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	2	<1%	3	<1%
Unknown	20	3%	21	3%	24	3%	27	4%	30	4%
Race/Sex										
White, not Hispanic Males	261	43%	279	42%	301	43%	327	43%	351	43%
White, not Hispanic Females	70	11%	74	11%	79	11%	83	11%	91	11%
Black/African American, not Hispanic Males	136	22%	147	22%	155	22%	172	23%	190	23%
Black/African American, not Hispanic Females	89	15%	97	15%	103	15%	105	14%	109	13%
Hispanic/Latino Males	21	3%	22	3%	23	3%	25	3%	26	3%
Hispanic/Latina Females	14	2%	15	2%	16	2%	16	2%	17	2%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	2	<1%	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	20	3%	21	3%	24	3%	27	4%	30	4%
Total	613		657		703		758		819	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 54. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 4, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	256	60%	279	61%	303	61%	334	62%	361	61%
Injection drug use (IDU)	17	4%	19	4%	19	4%	21	4%	21	4%
Male-to-male sexual contact and IDU	10	2%	10	2%	11	2%	13	2%	13	2%
Heterosexual contact	40	9%	42	9%	48	10%	55	10%	57	10%
Other/unknown	107	25%	111	24%	113	23%	120	22%	137	23%
Subtotal	430	100%	461	100%	494	100%	543	100%	589	100%
Female adult or adolescent										
Injection drug use	17	10%	17	9%	17	9%	18	9%	21	10%
Heterosexual contact	95	56%	105	58%	116	59%	119	59%	125	58%
Other/unknown	57	34%	60	33%	62	32%	64	32%	70	32%
Subtotal	169	100%	182	100%	195	100%	201	100%	216	100%
Child (<13 yrs at diagnosis)										
Perinatal	11	79%	11	79%	11	79%	11	79%	11	79%
Other/unknown	3	21%	3	21%	3	21%	3	21%	3	21%
Subtotal	14	100%	14	100%	14	100%	14	100%	14	100%
Total	613		657		703		758		819	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 55. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 4, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	6.0	27	75%	7.3	33	70%	8.4	38	73%	11.1	50	89%	10.6	48	76%
Females	1.9	9	25%	3.0	14	30%	3.0	14	27%	1.3	6	11%	3.2	15	24%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	3%	*	3	6%	*	2	4%	11.4	7	13%	8.1	5	8%
20-24	*	2	6%	16.5	9	19%	25.6	14	27%	12.8	7	13%	23.8	13	21%
25-29	*	4	11%	10.0	5	11%	*	4	8%	*	4	7%	23.9	12	19%
30-34	9.7	5	14%	*	3	6%	17.5	9	17%	*	4	7%	*	4	6%
35-39	9.6	5	14%	13.5	7	15%	11.5	6	12%	15.4	8	14%	*	4	6%
40-44	13.6	8	22%	8.5	5	11%	*	3	6%	20.5	12	21%	10.2	6	10%
45-49	13.8	9	25%	15.4	10	21%	7.7	5	10%	*	4	7%	*	4	6%
50-54	*	-	-	*	4	9%	*	4	8%	9.4	7	13%	6.7	5	8%
55-64	*	2	6%	*	1	2%	3.7	5	10%	*	3	5%	6.0	8	13%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	2	3%
Race/Ethnicity															
White, not Hispanic	2.2	18	50%	2.8	23	49%	4.0	32	62%	3.8	31	55%	4.0	32	51%
Black/African American, not Hispanic	20.7	15	42%	27.7	20	43%	20.7	15	29%	26.3	19	34%	31.8	23	37%
Hispanic/Latino	*	2	6%	*	3	6%	*	2	4%	*	2	4%	*	2	3%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	2%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	2%	*	1	2%
Unknown	*	1	3%	*	1	2%	*	3	6%	*	3	5%	*	4	6%
Race/Sex															
White, not Hispanic Males	3.8	15	42%	4.8	19	40%	6.9	27	52%	6.9	27	48%	6.1	24	38%
White, not Hispanic Females	*	3	8%	*	4	9%	1.2	5	10%	*	4	7%	1.9	8	13%
Black/African American, not Hispanic Males	28.2	10	28%	31.0	11	23%	22.6	8	15%	47.9	17	30%	53.6	19	30%
Black/African American, not Hispanic Females	13.6	5	14%	24.4	9	19%	19.0	7	13%	*	2	4%	*	4	6%
Hispanic/Latino Males	*	1	3%	*	2	4%	*	1	2%	*	2	4%	*	1	2%
Hispanic/Latina Females	*	1	3%	*	1	2%	*	1	2%	*	-	-	*	1	2%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	2%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	2%	*	1	2%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	3%	*	1	2%	*	3	6%	*	3	5%	*	4	6%
Total	3.9	36		5.1	47		5.6	52		6.0	56		6.8	63	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 56. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 4, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	18	67%	25	76%	26	68%	31	62%	27	56%
Injection drug use (IDU)	-	-	2	6%	1	3%	2	4%	1	2%
Male-to-male sexual contact and IDU	-	-	-	-	1	3%	2	4%	-	-
Heterosexual contact	4	15%	2	6%	6	16%	7	14%	2	4%
Other/unknown	5	19%	4	12%	4	11%	8	16%	18	38%
Subtotal	27	100%	33	100%	38	100%	50	100%	48	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	1	17%	3	20%
Heterosexual contact	7	78%	10	71%	11	79%	3	50%	6	40%
Other/unknown	2	22%	4	29%	3	21%	2	33%	6	40%
Subtotal	9	100%	14	100%	14	100%	6	100%	15	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	36		47		52		56		63	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 57. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristic, Region 4, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	20	69%	20	74%	18	69%	18	67%	28	78%	688	78%
Females	9	31%	7	26%	8	31%	9	33%	8	22%	191	22%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	7	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	1	4%	-	-	12	1%
20-24	-	-	1	4%	-	-	2	7%	5	14%	37	4%
25-29	1	3%	4	15%	1	4%	2	7%	4	11%	94	11%
30-34	4	14%	2	7%	6	23%	1	4%	2	6%	168	19%
35-39	4	14%	4	15%	1	4%	3	11%	5	14%	192	22%
40-44	6	21%	5	19%	5	19%	7	26%	6	17%	150	17%
45-49	6	21%	7	26%	5	19%	6	22%	3	8%	104	12%
50-54	2	7%	3	11%	5	19%	3	11%	3	8%	51	6%
55-64	6	21%	1	4%	3	12%	2	7%	7	19%	53	6%
65+	-	-	-	-	-	-	-	-	1	3%	11	1%
Race/Ethnicity												
White, not Hispanic	13	45%	17	63%	19	73%	10	37%	15	42%	546	62%
Black/African American, not Hispanic	14	48%	8	30%	6	23%	14	52%	18	50%	294	33%
Hispanic/Latino	1	3%	1	4%	-	-	1	4%	1	3%	31	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	1	4%	1	4%	2	7%	2	6%	8	1%
Race/Sex												
White, not Hispanic Males	11	38%	13	48%	14	54%	9	33%	12	33%	470	53%
White, not Hispanic Females	2	7%	4	15%	5	19%	1	4%	3	8%	76	9%
Black/African American, not Hispanic Males	8	28%	5	19%	3	12%	6	22%	15	42%	190	22%
Black/African American, not Hispanic Females	6	21%	3	11%	3	12%	8	30%	3	8%	104	12%
Hispanic/Latino Males	-	-	1	4%	-	-	1	4%	-	-	21	2%
Hispanic/Latina Females	1	3%	-	-	-	-	-	-	1	3%	10	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	1	4%	1	4%	2	7%	2	6%	8	1%
Total	29		27		26		27		36		879	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 4

Table 58. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 4, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	12	60%	16	80%	8	44%	11	61%	13	46%	435	64%
Injection drug use (IDU)	-	-	1	5%	1	6%	2	11%	1	4%	42	6%
Male-to-male sexual contact and IDU	1	5%	-	-	1	6%	-	-	-	-	35	5%
Heterosexual contact	3	15%	2	10%	3	17%	1	6%	6	21%	55	8%
Other/unknown	4	20%	1	5%	5	28%	4	22%	8	29%	116	17%
Subtotal	20	100%	20	100%	18	100%	18	100%	28	100%	683	100%
Female adult or adolescent												
Injection drug use	-	-	1	14%	2	25%	1	11%	4	50%	43	23%
Heterosexual contact	7	78%	6	86%	3	38%	5	56%	3	38%	95	51%
Other/unknown	2	22%	-	-	3	38%	3	33%	1	13%	48	26%
Subtotal	9	100%	7	100%	8	100%	9	100%	8	100%	186	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	8	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	20%
Subtotal	-	-	-	-	-	-	-	-	-	-	10	100%
Total	29		27		26		27		36		879	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 59. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 5

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	540.0	3,101	82%	1,755	81%	1,346	83%
Females	113.5	686	18%	408	19%	278	17%
Age at diagnosis (yr)							
<13	5.8	12	<1%	5	<1%	7	<1%
13-14	*	-	-	-	-	-	-
15-19	25.5	20	1%	15	1%	5	<1%
20-24	175.7	180	5%	142	7%	38	2%
25-29	291.0	303	8%	217	10%	86	5%
30-34	392.9	364	10%	243	11%	121	7%
35-39	520.1	411	11%	235	11%	176	11%
40-44	727.5	578	15%	327	15%	251	15%
45-49	923.6	725	19%	379	18%	346	21%
50-54	684.6	552	15%	274	13%	278	17%
55-64	419.7	539	14%	266	12%	273	17%
65+	86.8	103	3%	60	3%	43	3%
Race/Ethnicity							
White, not Hispanic	260.2	2,063	54%	1,196	55%	867	53%
Black/African American, not Hispanic	590.4	1,464	39%	822	38%	642	40%
Hispanic/Latino	237.8	138	4%	70	3%	68	4%
Asian/Pacific Islander	42.0	20	1%	11	1%	9	1%
American Indian/Alaska Native	*	4	<1%	3	<1%	1	<1%
Unknown	*	98	3%	61	3%	37	2%
Race/Sex							
White, not Hispanic Males	485.2	1,881	50%	1,078	50%	803	49%
White, not Hispanic Females	44.9	182	5%	118	5%	64	4%
Black/African American, not Hispanic Males	863.3	1,007	27%	557	26%	450	28%
Black/African American, not Hispanic Females	348.0	457	12%	265	12%	192	12%
Hispanic/Latino Males	363.8	112	3%	58	3%	54	3%
Hispanic/Latina Females	95.4	26	1%	12	1%	14	1%
Asian/Pacific Islander Males	63.9	15	<1%	8	<1%	7	<1%
Asian/Pacific Islander Females	20.7	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	98	3%	61	3%	37	2%
Total	321.3	3,787		2,163		1,624	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 60. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 5

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,116	69%	1,186	68%	930	69%
Injection drug use (IDU)	61	2%	19	1%	42	3%
Male-to-male sexual contact and IDU	82	3%	36	2%	46	3%
Heterosexual contact	198	6%	75	4%	123	9%
Other/unknown	629	20%	430	25%	199	15%
Subtotal	3,086	100%	1,746	100%	1,340	100%
Female adult or adolescent						
Injection drug use	56	8%	28	7%	28	10%
Heterosexual contact	356	53%	195	49%	161	60%
Other/unknown	258	39%	178	44%	80	30%
Subtotal	670	100%	401	100%	269	100%
Child (<13 yrs at diagnosis)						
Perinatal	25	81%	13	81%	12	80%
Other/unknown	6	19%	3	19%	3	20%
Subtotal	31	100%	16	100%	15	100%
Total	3,787		2,163		1,624	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 61. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 5, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,255	82%	2,468	82%	2,673	82%	2,881	82%	3,101	82%
Females	511	18%	553	18%	604	18%	639	18%	686	18%
Age at diagnosis (yr)										
<13	8	<1%	10	<1%	11	<1%	11	<1%	12	<1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	8	<1%	8	<1%	11	<1%	13	<1%	20	1%
20-24	34	1%	51	2%	94	3%	134	4%	180	5%
25-29	114	4%	161	5%	212	6%	255	7%	303	8%
30-34	210	8%	245	8%	285	9%	322	9%	364	10%
35-39	269	10%	303	10%	337	10%	371	11%	411	11%
40-44	447	16%	484	16%	520	16%	550	16%	578	15%
45-49	633	23%	665	22%	680	21%	704	20%	725	19%
50-54	482	17%	505	17%	519	16%	534	15%	552	15%
55-64	473	17%	497	16%	512	16%	526	15%	539	14%
65+	88	3%	92	3%	96	3%	100	3%	103	3%
Race/Ethnicity										
White, not Hispanic	1,564	57%	1,697	56%	1,820	56%	1,939	55%	2,063	54%
Black/African American, not Hispanic	1,025	37%	1,128	37%	1,243	38%	1,348	38%	1,464	39%
Hispanic/Latino	92	3%	101	3%	110	3%	122	3%	138	4%
Asian/Pacific Islander	12	<1%	14	<1%	16	<1%	18	1%	20	1%
American Indian/Alaska Native	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	70	3%	77	3%	84	3%	89	3%	98	3%
Race/Sex										
White, not Hispanic Males	1,430	52%	1,552	51%	1,662	51%	1,770	50%	1,881	50%
White, not Hispanic Females	134	5%	145	5%	158	5%	169	5%	182	5%
Black/African American, not Hispanic Males	682	25%	756	25%	837	26%	921	26%	1,007	27%
Black/African American, not Hispanic Females	343	12%	372	12%	406	12%	427	12%	457	12%
Hispanic/Latino Males	72	3%	81	3%	89	3%	100	3%	112	3%
Hispanic/Latina Females	20	1%	20	1%	21	1%	22	1%	26	1%
Asian/Pacific Islander Males	9	<1%	11	<1%	12	<1%	13	<1%	15	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	5	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	70	3%	77	3%	84	3%	89	3%	98	3%
Total	2,766		3,021		3,277		3,520		3,787	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 62. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 5, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,517	68%	1,683	69%	1,836	69%	1,974	69%	2,116	69%
Injection drug use (IDU)	56	3%	57	2%	58	2%	58	2%	61	2%
Male-to-male sexual contact and IDU	56	3%	63	3%	69	3%	77	3%	82	3%
Heterosexual contact	146	7%	171	7%	186	7%	195	7%	198	6%
Other/unknown	465	21%	479	20%	509	19%	562	20%	629	20%
Subtotal	2,240	100%	2,453	100%	2,658	100%	2,866	100%	3,086	100%
Female adult or adolescent										
Injection drug use	49	10%	52	10%	53	9%	56	9%	56	8%
Heterosexual contact	270	54%	293	54%	326	55%	340	54%	356	53%
Other/unknown	181	36%	195	36%	210	36%	228	37%	258	39%
Subtotal	500	100%	540	100%	589	100%	624	100%	670	100%
Child (<13 yrs at diagnosis)										
Perinatal	20	77%	22	79%	24	80%	24	80%	25	81%
Other/unknown	6	23%	6	21%	6	20%	6	20%	6	19%
Subtotal	26	100%	28	100%	30	100%	30	100%	31	100%
Total	2,766		3,021		3,277		3,520		3,787	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 63. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 5, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	38.1	219	82%	40.2	231	84%	37.8	217	81%	38.1	219	86%	38.8	223	82%
Females	7.8	47	18%	7.1	43	16%	8.6	52	19%	6.0	36	14%	8.1	49	18%
Age at diagnosis (yr)															
<13	*	3	1%	*	2	1%	*	2	1%	*	-	-	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	16.6	13	5%	16.6	13	5%	33.2	26	10%	15.3	12	5%	19.1	15	6%
20-24	39.0	40	15%	44.9	46	17%	48.8	50	19%	50.8	52	20%	48.8	50	18%
25-29	52.8	55	21%	33.6	35	13%	50.9	53	20%	38.4	40	16%	42.3	44	16%
30-34	27.0	25	9%	44.3	41	15%	34.5	32	12%	38.9	36	14%	44.3	41	15%
35-39	51.9	41	15%	48.1	38	14%	48.1	38	14%	41.8	33	13%	50.6	40	15%
40-44	59.2	47	18%	47.8	38	14%	32.7	26	10%	39.0	31	12%	39.0	31	11%
45-49	25.5	20	8%	35.7	28	10%	20.4	16	6%	31.8	25	10%	22.9	18	7%
50-54	17.4	14	5%	28.5	23	8%	16.1	13	5%	12.4	10	4%	22.3	18	7%
55-64	6.2	8	3%	4.7	6	2%	7.8	10	4%	10.9	14	5%	8.6	11	4%
65+	*	-	-	*	4	1%	*	3	1%	*	2	1%	*	3	1%
Race/Ethnicity															
White, not Hispanic	16.9	134	50%	17.9	142	52%	16.5	131	49%	15.8	125	49%	16.0	127	47%
Black/African American, not Hispanic	43.6	108	41%	45.2	112	41%	48.4	120	45%	44.8	111	44%	47.6	118	43%
Hispanic/Latino	22.4	13	5%	17.2	10	4%	15.5	9	3%	20.7	12	5%	27.6	16	6%
Asian/Pacific Islander	*	3	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	3%	*	7	3%	*	5	2%	*	9	3%
Race/Sex															
White, not Hispanic Males	31.0	120	45%	33.8	131	48%	30.4	118	44%	29.4	114	45%	29.1	113	42%
White, not Hispanic Females	3.5	14	5%	2.7	11	4%	3.2	13	5%	2.7	11	4%	3.5	14	5%
Black/African American, not Hispanic Males	66.9	78	29%	71.2	83	30%	72.9	85	32%	76.3	89	35%	74.6	87	32%
Black/African American, not Hispanic Females	22.8	30	11%	22.1	29	11%	26.7	35	13%	16.8	22	9%	23.6	31	11%
Hispanic/Latino Males	39.0	12	5%	29.2	9	3%	26.0	8	3%	35.7	11	4%	39.0	12	4%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	4	1%
Asian/Pacific Islander Males	*	3	1%	*	2	1%	*	1	<1%	*	1	<1%	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	1	<1%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	3%	*	7	3%	*	5	2%	*	9	3%
Total	22.6	266		23.2	274		22.8	269		21.6	255		23.1	272	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 64. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 5, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	173	79%	179	77%	161	74%	144	66%	143	64%
Injection drug use (IDU)	6	3%	2	1%	1	<1%	-	-	3	1%
Male-to-male sexual contact and IDU	6	3%	9	4%	6	3%	8	4%	5	2%
Heterosexual contact	9	4%	26	11%	15	7%	9	4%	3	1%
Other/unknown	24	11%	15	6%	34	16%	58	26%	69	31%
Subtotal	218	100%	231	100%	217	100%	219	100%	223	100%
Female adult or adolescent										
Injection drug use	4	9%	3	7%	1	2%	3	8%	-	-
Heterosexual contact	24	53%	22	54%	33	66%	14	39%	16	33%
Other/unknown	17	38%	16	39%	16	32%	19	53%	32	67%
Subtotal	45	100%	41	100%	50	100%	36	100%	48	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	2	100%	2	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	2	100%	-	-	1	100%
Total	266		274		269		255		272	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 65. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 5, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	92	79%	99	83%	123	79%	124	86%	119	80%	2,974	87%
Females	25	21%	20	17%	32	21%	20	14%	30	20%	435	13%
Age at AIDS diagnosis (yr)												
<13	2	2%	2	2%	1	1%	-	-	1	1%	17	<1%
13-14	1	1%	-	-	-	-	-	-	-	-	4	<1%
15-19	5	4%	1	1%	3	2%	3	2%	3	2%	29	1%
20-24	8	7%	10	8%	12	8%	15	10%	15	10%	181	5%
25-29	12	10%	12	10%	27	17%	15	10%	13	9%	455	13%
30-34	12	10%	21	18%	18	12%	22	15%	27	18%	736	22%
35-39	18	15%	20	17%	19	12%	19	13%	30	20%	676	20%
40-44	23	20%	19	16%	22	14%	27	19%	21	14%	564	17%
45-49	18	15%	17	14%	28	18%	21	15%	15	10%	379	11%
50-54	11	9%	11	9%	13	8%	8	6%	9	6%	195	6%
55-64	7	6%	5	4%	9	6%	11	8%	15	10%	149	4%
65+	-	-	1	1%	3	2%	3	2%	-	-	23	1%
Race/Ethnicity												
White, not Hispanic	55	47%	53	45%	66	43%	72	50%	62	42%	2,141	63%
Black/African American, not Hispanic	49	42%	55	46%	78	50%	67	47%	69	46%	1,117	33%
Hispanic/Latino	8	7%	4	3%	6	4%	4	3%	9	6%	95	3%
Asian/Pacific Islander	1	1%	1	1%	1	1%	-	-	2	1%	14	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	4	3%	6	5%	4	3%	1	1%	6	4%	40	1%
Race/Sex												
White, not Hispanic Males	48	41%	51	43%	58	37%	70	49%	56	38%	2,014	59%
White, not Hispanic Females	7	6%	2	2%	8	5%	2	1%	6	4%	127	4%
Black/African American, not Hispanic Males	32	27%	39	33%	56	36%	50	35%	49	33%	834	24%
Black/African American, not Hispanic Females	17	15%	16	13%	22	14%	17	12%	20	13%	283	8%
Hispanic/Latino Males	7	6%	3	3%	6	4%	4	3%	6	4%	79	2%
Hispanic/Latina Females	1	1%	1	1%	-	-	-	-	3	2%	16	<1%
Asian/Pacific Islander Males	1	1%	1	1%	-	-	-	-	2	1%	12	<1%
Asian/Pacific Islander Females	-	-	-	-	1	1%	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	4	3%	6	5%	4	3%	1	1%	6	4%	40	1%
Total	117		119		155		144		149		3,409	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 5

Table 66. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 5, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	64	71%	66	67%	87	71%	71	57%	72	61%	2,179	74%
Injection drug use (IDU)	1	1%	3	3%	3	2%	-	-	2	2%	148	5%
Male-to-male sexual contact and IDU	5	6%	7	7%	3	2%	6	5%	4	3%	114	4%
Heterosexual contact	12	13%	15	15%	16	13%	9	7%	5	4%	155	5%
Other/unknown	8	9%	7	7%	14	11%	38	31%	36	30%	368	12%
Subtotal	90	100%	98	100%	123	100%	124	100%	119	100%	2,964	100%
Female adult or adolescent												
Injection drug use	4	17%	1	6%	2	6%	2	10%	-	-	61	14%
Heterosexual contact	14	58%	10	56%	18	58%	10	50%	13	45%	229	54%
Other/unknown	6	25%	7	39%	11	35%	8	40%	16	55%	132	31%
Subtotal	24	100%	18	100%	31	100%	20	100%	29	100%	422	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	100%	2	67%	1	100%	-	-	1	100%	18	78%
Other/unknown	-	-	1	33%	-	-	-	-	-	-	5	22%
Subtotal	3	100%	3	100%	1	100%	-	-	1	100%	23	100%
Total	117		119		155		144		149		3,409	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 67. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 6

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	477.4	1,835	79%	898	77%	937	81%
Females	115.6	481	21%	267	23%	214	19%
Age at diagnosis (yr)							
<13	*	2	<1%	1	<1%	1	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	45.2	25	1%	23	2%	2	<1%
20-24	219.5	130	6%	97	8%	33	3%
25-29	293.3	172	7%	117	10%	55	5%
30-34	328.6	170	7%	97	8%	73	6%
35-39	536.4	243	10%	139	12%	104	9%
40-44	581.4	291	13%	134	12%	157	14%
45-49	793.2	446	19%	193	17%	253	22%
50-54	604.6	367	16%	184	16%	183	16%
55-64	376.4	379	16%	142	12%	237	21%
65+	84.3	90	4%	38	3%	52	5%
Race/Ethnicity							
White, not Hispanic	168.6	912	39%	465	40%	447	39%
Black/African American, not Hispanic	651.0	1,330	57%	658	56%	672	58%
Hispanic/Latino	172.0	37	2%	20	2%	17	1%
Asian/Pacific Islander	64.5	11	<1%	7	1%	4	<1%
American Indian/Alaska Native	*	2	<1%	-	-	2	<1%
Unknown	*	24	1%	15	1%	9	1%
Race/Sex							
White, not Hispanic Males	299.6	789	34%	383	33%	406	35%
White, not Hispanic Females	44.3	123	5%	82	7%	41	4%
Black/African American, not Hispanic Males	1,057.2	988	43%	482	41%	506	44%
Black/African American, not Hispanic Females	308.5	342	15%	176	15%	166	14%
Hispanic/Latino Males	247.1	28	1%	15	1%	13	1%
Hispanic/Latina Females	88.4	9	<1%	5	<1%	4	<1%
Asian/Pacific Islander Males	107.7	9	<1%	6	1%	3	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	-	-	2	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	24	1%	15	1%	9	1%
Total	289.4	2,316		1,165		1,151	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 68. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 6

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,231	67%	567	63%	664	71%
Injection drug use (IDU)	58	3%	26	3%	32	3%
Male-to-male sexual contact and IDU	75	4%	27	3%	48	5%
Heterosexual contact	109	6%	46	5%	63	7%
Other/unknown	354	19%	227	25%	127	14%
Subtotal	1,827	100%	893	100%	934	100%
Female adult or adolescent						
Injection drug use	44	9%	18	7%	26	12%
Heterosexual contact	252	53%	129	49%	123	58%
Other/unknown	178	38%	116	44%	62	29%
Subtotal	474	100%	263	100%	211	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	67%	7	78%	3	50%
Other/unknown	5	33%	2	22%	3	50%
Subtotal	15	100%	9	100%	6	100%
Total	2,316		1,165		1,151	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 69. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 6, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	1,422	81%	1,506	80%	1,618	80%	1,733	80%	1,835	79%
Females	343	19%	371	20%	410	20%	444	20%	481	21%
Age at diagnosis (yr)										
<13	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	4	<1%	4	<1%	5	<1%	14	1%	25	1%
20-24	21	1%	35	2%	67	3%	99	5%	130	6%
25-29	67	4%	96	5%	125	6%	151	7%	172	7%
30-34	109	6%	119	6%	136	7%	156	7%	170	7%
35-39	188	11%	197	10%	212	10%	227	10%	243	10%
40-44	244	14%	255	14%	265	13%	279	13%	291	13%
45-49	385	22%	405	22%	420	21%	436	20%	446	19%
50-54	318	18%	328	17%	341	17%	353	16%	367	16%
55-64	340	19%	349	19%	367	18%	372	17%	379	16%
65+	86	5%	86	5%	87	4%	87	4%	90	4%
Race/Ethnicity										
White, not Hispanic	757	43%	790	42%	826	41%	869	40%	912	39%
Black/African American, not Hispanic	954	54%	1,028	55%	1,136	56%	1,239	57%	1,330	57%
Hispanic/Latino	26	1%	30	2%	34	2%	34	2%	37	2%
Asian/Pacific Islander	8	<1%	8	<1%	9	<1%	9	<1%	11	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	18	1%	19	1%	21	1%	24	1%	24	1%
Race/Sex										
White, not Hispanic Males	665	38%	691	37%	720	36%	754	35%	789	34%
White, not Hispanic Females	92	5%	99	5%	106	5%	115	5%	123	5%
Black/African American, not Hispanic Males	715	41%	768	41%	845	42%	924	42%	988	43%
Black/African American, not Hispanic Females	239	14%	260	14%	291	14%	315	14%	342	15%
Hispanic/Latino Males	19	1%	23	1%	26	1%	26	1%	28	1%
Hispanic/Latina Females	7	<1%	7	<1%	8	<1%	8	<1%	9	<1%
Asian/Pacific Islander Males	7	<1%	7	<1%	8	<1%	8	<1%	9	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	18	1%	19	1%	21	1%	24	1%	24	1%
Total	1,765		1,877		2,028		2,177		2,316	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 70. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 6, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	954	67%	1,013	68%	1,091	68%	1,172	68%	1,231	67%
Injection drug use (IDU)	53	4%	56	4%	58	4%	58	3%	58	3%
Male-to-male sexual contact and IDU	65	5%	68	5%	70	4%	73	4%	75	4%
Heterosexual contact	83	6%	90	6%	96	6%	106	6%	109	6%
Other/unknown	259	18%	271	18%	295	18%	316	18%	354	19%
Subtotal	1,414	100%	1,498	100%	1,610	100%	1,725	100%	1,827	100%
Female adult or adolescent										
Injection drug use	30	9%	34	9%	40	10%	43	10%	44	9%
Heterosexual contact	205	61%	219	60%	231	57%	242	55%	252	53%
Other/unknown	101	30%	111	30%	132	33%	152	35%	178	38%
Subtotal	336	100%	364	100%	403	100%	437	100%	474	100%
Child (<13 yrs at diagnosis)										
Perinatal	10	67%	10	67%	10	67%	10	67%	10	67%
Other/unknown	5	33%	5	33%	5	33%	5	33%	5	33%
Subtotal	15	100%	15	100%	15	100%	15	100%	15	100%
Total	1,765		1,877		2,028		2,177		2,316	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 71. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 6, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	25.2	97	82%	23.9	92	75%	30.4	117	73%	30.4	117	76%	27.1	104	73%
Females	5.0	21	18%	7.2	30	25%	10.3	43	27%	8.7	36	24%	9.1	38	27%
Age at diagnosis (yr)															
<13	*	1	1%	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	14.5	8	7%	12.7	7	6%	29.0	16	10%	32.6	18	12%	21.7	12	8%
20-24	37.1	22	19%	57.4	34	28%	62.5	37	23%	57.4	34	22%	55.7	33	23%
25-29	25.6	15	13%	18.8	11	9%	37.5	22	14%	46.0	27	18%	37.5	22	15%
30-34	30.9	16	14%	17.4	9	7%	34.8	18	11%	32.9	17	11%	30.9	16	11%
35-39	24.3	11	9%	30.9	14	11%	22.1	10	6%	30.9	14	9%	24.3	11	8%
40-44	34.0	17	14%	50.0	25	20%	24.0	12	8%	30.0	15	10%	26.0	13	9%
45-49	24.9	14	12%	14.2	8	7%	28.5	16	10%	33.8	19	12%	28.5	16	11%
50-54	11.5	7	6%	16.5	10	8%	24.7	15	9%	13.2	8	5%	18.1	11	8%
55-64	6.0	6	5%	*	4	3%	12.9	13	8%	*	1	1%	6.0	6	4%
65+	*	1	1%	*	-	-	*	1	1%	*	-	-	*	2	1%
Race/Ethnicity															
White, not Hispanic	6.8	37	31%	6.1	33	27%	7.0	38	24%	8.3	45	29%	8.3	45	32%
Black/African American, not Hispanic	36.7	75	64%	40.6	83	68%	56.3	115	72%	50.9	104	68%	45.0	92	65%
Hispanic/Latino	*	4	3%	23.2	5	4%	*	4	3%	*	-	-	*	3	2%
Asian/Pacific Islander	*	1	1%	*	-	-	*	1	1%	*	-	-	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	1	1%	*	2	1%	*	4	3%	*	-	-
Race/Sex															
White, not Hispanic Males	12.2	32	27%	9.9	26	21%	11.4	30	19%	13.3	35	23%	14.1	37	26%
White, not Hispanic Females	1.8	5	4%	2.5	7	6%	2.9	8	5%	3.6	10	7%	2.9	8	6%
Black/African American, not Hispanic Males	66.3	62	53%	64.2	60	49%	86.7	81	51%	85.6	80	52%	68.5	64	45%
Black/African American, not Hispanic Females	11.7	13	11%	20.7	23	19%	30.7	34	21%	21.6	24	16%	25.3	28	20%
Hispanic/Latino Males	*	2	2%	44.1	5	4%	*	3	2%	*	-	-	*	2	1%
Hispanic/Latina Females	*	2	2%	*	-	-	*	1	1%	*	-	-	*	1	1%
Asian/Pacific Islander Males	*	1	1%	*	-	-	*	1	1%	*	-	-	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	1	1%	*	2	1%	*	4	3%	*	-	-
Total	14.7	118		15.2	122		20.0	160		19.1	153		17.7	142	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 72. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 6, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	71	73%	62	67%	81	69%	83	71%	60	58%
Injection drug use (IDU)	1	1%	3	3%	2	2%	-	-	-	-
Male-to-male sexual contact and IDU	2	2%	3	3%	2	2%	3	3%	3	3%
Heterosexual contact	15	15%	7	8%	6	5%	10	9%	3	3%
Other/unknown	8	8%	17	18%	26	22%	21	18%	38	37%
Subtotal	97	100%	92	100%	117	100%	117	100%	104	100%
Female adult or adolescent										
Injection drug use	1	5%	4	13%	7	16%	3	8%	1	3%
Heterosexual contact	15	75%	15	50%	13	30%	11	31%	10	26%
Other/unknown	4	20%	11	37%	23	53%	22	61%	27	71%
Subtotal	20	100%	30	100%	43	100%	36	100%	38	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-
Total	118		122		160		153		142	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 73. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 6, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	40	83%	64	77%	74	72%	70	81%	56	76%	1,990	86%
Females	8	17%	19	23%	29	28%	16	19%	18	24%	311	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	16	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	4	5%	6	6%	-	-	2	3%	21	1%
20-24	6	13%	8	10%	11	11%	7	8%	15	20%	114	5%
25-29	4	8%	5	6%	16	16%	21	24%	7	9%	347	15%
30-34	5	10%	6	7%	10	10%	12	14%	13	18%	473	21%
35-39	7	15%	15	18%	13	13%	9	10%	8	11%	487	21%
40-44	9	19%	22	27%	20	19%	10	12%	3	4%	375	16%
45-49	9	19%	10	12%	10	10%	12	14%	8	11%	212	9%
50-54	2	4%	9	11%	11	11%	8	9%	10	14%	139	6%
55-64	5	10%	4	5%	5	5%	6	7%	7	9%	92	4%
65+	1	2%	-	-	1	1%	1	1%	1	1%	25	1%
Race/Ethnicity												
White, not Hispanic	13	27%	26	31%	29	28%	25	29%	22	30%	1,095	48%
Black/African American, not Hispanic	33	69%	55	66%	67	65%	60	70%	51	69%	1,164	51%
Hispanic/Latino	2	4%	1	1%	3	3%	-	-	1	1%	24	1%
Asian/Pacific Islander	-	-	-	-	1	1%	-	-	-	-	6	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	-	-	1	1%	3	3%	1	1%	-	-	10	<1%
Race/Sex												
White, not Hispanic Males	11	23%	24	29%	25	24%	23	27%	18	24%	1,028	45%
White, not Hispanic Females	2	4%	2	2%	4	4%	2	2%	4	5%	67	3%
Black/African American, not Hispanic Males	28	58%	38	46%	46	45%	46	53%	37	50%	929	40%
Black/African American, not Hispanic Females	5	10%	17	20%	21	20%	14	16%	14	19%	235	10%
Hispanic/Latino Males	1	2%	1	1%	1	1%	-	-	1	1%	18	1%
Hispanic/Latina Females	1	2%	-	-	2	2%	-	-	-	-	6	<1%
Asian/Pacific Islander Males	-	-	-	-	1	1%	-	-	-	-	5	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	1%	3	3%	1	1%	-	-	10	<1%
Total	48		83		103		86		74		2,301	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 6

Table 74. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 6, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	29	73%	46	72%	48	65%	52	74%	34	61%	1,440	73%
Injection drug use (IDU)	1	3%	1	2%	2	3%	-	-	-	-	88	4%
Male-to-male sexual contact and IDU	1	3%	1	2%	3	4%	3	4%	2	4%	125	6%
Heterosexual contact	7	18%	7	11%	9	12%	3	4%	1	2%	85	4%
Other/unknown	2	5%	9	14%	12	16%	12	17%	19	34%	239	12%
Subtotal	40	100%	64	100%	74	100%	70	100%	56	100%	1,977	100%
Female adult or adolescent												
Injection drug use	-	-	1	5%	5	17%	1	6%	2	12%	50	16%
Heterosexual contact	6	75%	16	84%	14	48%	4	25%	5	29%	167	55%
Other/unknown	2	25%	2	11%	10	34%	11	69%	10	59%	88	29%
Subtotal	8	100%	19	100%	29	100%	16	100%	17	100%	305	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	1	100%	14	74%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	26%
Subtotal	-	-	-	-	-	-	-	-	1	100%	19	100%
Total	48		83		103		86		74		2,301	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 75. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 7

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	199.5	1,286	78%	606	76%	680	81%
Females	53.3	360	22%	196	24%	164	19%
Age at diagnosis (yr)							
<13	*	4	<1%	4	<1%	-	-
13-14	*	2	<1%	2	<1%	-	-
15-19	15.4	14	1%	9	1%	5	1%
20-24	81.6	70	4%	51	6%	19	2%
25-29	155.9	124	8%	77	10%	47	6%
30-34	164.1	133	8%	72	9%	61	7%
35-39	215.2	171	10%	93	12%	78	9%
40-44	241.4	218	13%	99	12%	119	14%
45-49	318.7	306	19%	131	16%	175	21%
50-54	260.1	260	16%	124	15%	136	16%
55-64	161.9	272	17%	107	13%	165	20%
65+	39.1	72	4%	33	4%	39	5%
Race/Ethnicity							
White, not Hispanic	78.1	844	51%	394	49%	450	53%
Black/African American, not Hispanic	491.2	731	44%	371	46%	360	43%
Hispanic/Latino	90.0	32	2%	15	2%	17	2%
Asian/Pacific Islander	30.6	9	1%	5	1%	4	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	29	2%	17	2%	12	1%
Race/Sex							
White, not Hispanic Males	135.8	718	44%	325	41%	393	47%
White, not Hispanic Females	22.8	126	8%	69	9%	57	7%
Black/African American, not Hispanic Males	717.2	511	31%	252	31%	259	31%
Black/African American, not Hispanic Females	283.6	220	13%	119	15%	101	12%
Hispanic/Latino Males	150.6	28	2%	13	2%	15	2%
Hispanic/Latina Females	*	4	<1%	2	<1%	2	<1%
Asian/Pacific Islander Males	57.9	8	<1%	5	1%	3	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	29	2%	17	2%	12	1%
Total	124.7	1,646		802		844	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 76. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 7

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	891	70%	390	65%	501	74%
Injection drug use (IDU)	35	3%	18	3%	17	3%
Male-to-male sexual contact and IDU	40	3%	12	2%	28	4%
Heterosexual contact	79	6%	33	5%	46	7%
Other/unknown	232	18%	148	25%	84	12%
Subtotal	1,277	100%	601	100%	676	100%
Female adult or adolescent						
Injection drug use	39	11%	23	12%	16	10%
Heterosexual contact	208	59%	101	53%	107	67%
Other/unknown	104	30%	67	35%	37	23%
Subtotal	351	100%	191	100%	160	100%
Child (<13 yrs at diagnosis)						
Perinatal	16	89%	9	90%	7	88%
Other/unknown	2	11%	1	10%	1	13%
Subtotal	18	100%	10	100%	8	100%
Total	1,646		802		844	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 77. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 7, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	997	78%	1,058	78%	1,112	78%	1,193	78%	1,286	78%
Females	274	22%	296	22%	315	22%	333	22%	360	22%
Age at diagnosis (yr)										
<13	2	<1%	2	<1%	2	<1%	4	<1%	4	<1%
13-14	1	<1%	1	<1%	2	<1%	2	<1%	2	<1%
15-19	7	1%	7	1%	7	<1%	11	1%	14	1%
20-24	12	1%	20	1%	28	2%	43	3%	70	4%
25-29	61	5%	74	5%	87	6%	111	7%	124	8%
30-34	78	6%	89	7%	103	7%	116	8%	133	8%
35-39	131	10%	141	10%	150	11%	155	10%	171	10%
40-44	179	14%	189	14%	193	14%	205	13%	218	13%
45-49	270	21%	281	21%	288	20%	296	19%	306	19%
50-54	215	17%	226	17%	235	16%	245	16%	260	16%
55-64	250	20%	259	19%	265	19%	267	17%	272	17%
65+	65	5%	65	5%	67	5%	71	5%	72	4%
Race/Ethnicity										
White, not Hispanic	706	56%	737	54%	761	53%	798	52%	844	51%
Black/African American, not Hispanic	517	41%	559	41%	605	42%	662	43%	731	44%
Hispanic/Latino	20	2%	28	2%	29	2%	30	2%	32	2%
Asian/Pacific Islander	7	1%	8	1%	8	1%	9	1%	9	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	20	2%	21	2%	23	2%	26	2%	29	2%
Race/Sex										
White, not Hispanic Males	601	47%	626	46%	646	45%	679	44%	718	44%
White, not Hispanic Females	105	8%	111	8%	115	8%	119	8%	126	8%
Black/African American, not Hispanic Males	358	28%	386	29%	417	29%	462	30%	511	31%
Black/African American, not Hispanic Females	159	13%	173	13%	188	13%	200	13%	220	13%
Hispanic/Latino Males	17	1%	24	2%	25	2%	26	2%	28	2%
Hispanic/Latina Females	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Asian/Pacific Islander Males	6	<1%	7	1%	7	<1%	8	1%	8	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	20	2%	21	2%	23	2%	26	2%	29	2%
Total	1,271		1,354		1,427		1,526		1,646	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 78. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 7, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	700	71%	741	70%	779	70%	839	71%	891	70%
Injection drug use (IDU)	32	3%	32	3%	34	3%	34	3%	35	3%
Male-to-male sexual contact and IDU	38	4%	38	4%	39	4%	40	3%	40	3%
Heterosexual contact	60	6%	65	6%	67	6%	70	6%	79	6%
Other/unknown	161	16%	176	17%	186	17%	201	17%	232	18%
Subtotal	991	100%	1,052	100%	1,105	100%	1,184	100%	1,277	100%
Female adult or adolescent										
Injection drug use	34	13%	34	12%	35	11%	36	11%	39	11%
Heterosexual contact	171	65%	185	64%	193	63%	200	62%	208	59%
Other/unknown	60	23%	68	24%	78	25%	88	27%	104	30%
Subtotal	265	100%	287	100%	306	100%	324	100%	351	100%
Child (<13 yrs at diagnosis)										
Perinatal	14	93%	14	93%	15	94%	16	89%	16	89%
Other/unknown	1	7%	1	7%	1	6%	2	11%	2	11%
Subtotal	15	100%	15	100%	16	100%	18	100%	18	100%
Total	1,271		1,354		1,427		1,526		1,646	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 79. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 7, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	11.2	72	76%	10.5	68	75%	8.8	57	73%	13.0	84	82%	14.9	96	78%
Females	3.4	23	24%	3.4	23	25%	3.1	21	27%	2.8	19	18%	4.0	27	22%
Age at diagnosis (yr)															
<13	*	1	1%	*	-	-	*	1	1%	*	2	2%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	4	4%	6.6	6	7%	5.5	5	6%	11.0	10	10%	6.6	6	5%
20-24	15.2	13	14%	15.2	13	14%	14.0	12	15%	26.8	23	22%	35.0	30	24%
25-29	13.8	11	12%	16.3	13	14%	16.3	13	17%	23.9	19	18%	15.1	12	10%
30-34	14.8	12	13%	12.3	10	11%	14.8	12	15%	6.2	5	5%	23.4	19	15%
35-39	20.1	16	17%	12.6	10	11%	6.3	5	6%	13.8	11	11%	15.1	12	10%
40-44	13.3	12	13%	14.4	13	14%	7.7	7	9%	11.1	10	10%	15.5	14	11%
45-49	9.4	9	9%	11.5	11	12%	10.4	10	13%	9.4	9	9%	11.5	11	9%
50-54	10.0	10	11%	8.0	8	9%	7.0	7	9%	7.0	7	7%	13.0	13	11%
55-64	*	4	4%	3.6	6	7%	3.6	6	8%	*	2	2%	*	4	3%
65+	*	3	3%	*	1	1%	*	-	-	2.7	5	5%	*	2	2%
Race/Ethnicity															
White, not Hispanic	3.6	39	41%	3.5	38	42%	2.4	26	33%	3.6	39	38%	4.3	47	38%
Black/African American, not Hispanic	32.3	48	51%	28.9	43	47%	32.9	49	63%	39.6	59	57%	47.7	71	58%
Hispanic/Latino	*	3	3%	22.5	8	9%	*	1	1%	*	1	1%	*	2	2%
Asian/Pacific Islander	*	2	2%	*	1	1%	*	-	-	*	1	1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	3%	*	1	1%	*	2	3%	*	3	3%	*	3	2%
Race/Sex															
White, not Hispanic Males	6.6	35	37%	5.9	31	34%	4.2	22	28%	6.6	35	34%	7.6	40	33%
White, not Hispanic Females	*	4	4%	1.3	7	8%	*	4	5%	*	4	4%	1.3	7	6%
Black/African American, not Hispanic Males	43.5	31	33%	40.7	29	32%	44.9	32	41%	64.6	46	45%	71.6	51	41%
Black/African American, not Hispanic Females	21.9	17	18%	18.0	14	15%	21.9	17	22%	16.8	13	13%	25.8	20	16%
Hispanic/Latino Males	*	3	3%	37.6	7	8%	*	1	1%	*	1	1%	*	2	2%
Hispanic/Latina Females	*	-	-	*	1	1%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	2%	*	1	1%	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	3%	*	1	1%	*	2	3%	*	3	3%	*	3	2%
Total	7.2	95		6.9	91		5.9	78		7.8	103		9.3	123	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 80. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 7, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	52	72%	45	66%	39	70%	61	74%	52	54%
Injection drug use (IDU)	5	7%	-	-	2	4%	1	1%	1	1%
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	1	1%	-	-
Heterosexual contact	7	10%	5	7%	2	4%	3	4%	9	9%
Other/unknown	8	11%	18	26%	12	21%	16	20%	34	35%
Subtotal	72	100%	68	100%	56	100%	82	100%	96	100%
Female adult or adolescent										
Injection drug use	2	9%	-	-	1	5%	1	5%	3	11%
Heterosexual contact	15	68%	14	61%	8	38%	7	37%	8	30%
Other/unknown	5	23%	9	39%	12	57%	11	58%	16	59%
Subtotal	22	100%	23	100%	21	100%	19	100%	27	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	1	100%	1	50%	-	-
Other/unknown	-	-	-	-	-	-	1	50%	-	-
Subtotal	1	100%	-	-	1	100%	2	100%	-	-
Total	95		91		78		103		123	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 81. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 7, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	47	80%	48	75%	38	73%	36	75%	46	81%	1,418	85%
Females	12	20%	16	25%	14	27%	12	25%	11	19%	251	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	14	1%
13-14	-	-	-	-	-	-	-	-	-	-	4	<1%
15-19	-	-	1	2%	1	2%	2	4%	2	4%	24	1%
20-24	5	8%	7	11%	5	10%	4	8%	10	18%	92	6%
25-29	5	8%	10	16%	8	15%	8	17%	7	12%	212	13%
30-34	6	10%	7	11%	6	12%	7	15%	6	11%	309	19%
35-39	12	20%	6	9%	7	13%	3	6%	6	11%	335	20%
40-44	15	25%	11	17%	7	13%	5	10%	7	12%	287	17%
45-49	6	10%	7	11%	7	13%	8	17%	8	14%	189	11%
50-54	5	8%	7	11%	7	13%	5	10%	5	9%	105	6%
55-64	1	2%	8	13%	4	8%	3	6%	5	9%	73	4%
65+	4	7%	-	-	-	-	3	6%	1	2%	25	1%
Race/Ethnicity												
White, not Hispanic	25	42%	27	42%	21	40%	23	48%	27	47%	1,022	61%
Black/African American, not Hispanic	32	54%	32	50%	29	56%	24	50%	24	42%	608	36%
Hispanic/Latino	-	-	3	5%	1	2%	-	-	2	4%	22	1%
Asian/Pacific Islander	-	-	2	3%	-	-	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	2	3%	-	-	1	2%	1	2%	4	7%	12	1%
Race/Sex												
White, not Hispanic Males	24	41%	24	38%	17	33%	20	42%	23	40%	929	56%
White, not Hispanic Females	1	2%	3	5%	4	8%	3	6%	4	7%	93	6%
Black/African American, not Hispanic Males	22	37%	19	30%	19	37%	15	31%	19	33%	455	27%
Black/African American, not Hispanic Females	10	17%	13	20%	10	19%	9	19%	5	9%	153	9%
Hispanic/Latino Males	-	-	3	5%	1	2%	-	-	2	4%	21	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	-	-	2	3%	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	3%	-	-	1	2%	1	2%	4	7%	12	1%
Total	59		64		52		48		57		1,669	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 7

Table 82. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 7, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	32	68%	36	75%	27	71%	28	78%	31	67%	1,008	72%
Injection drug use (IDU)	1	2%	1	2%	1	3%	1	3%	-	-	59	4%
Male-to-male sexual contact and IDU	2	4%	-	-	-	-	2	6%	1	2%	91	6%
Heterosexual contact	4	9%	4	8%	4	11%	1	3%	5	11%	80	6%
Other/unknown	8	17%	7	15%	6	16%	4	11%	9	20%	166	12%
Subtotal	47	100%	48	100%	38	100%	36	100%	46	100%	1,404	100%
Female adult or adolescent												
Injection drug use	1	8%	-	-	-	-	-	-	-	-	42	17%
Heterosexual contact	10	83%	12	75%	9	64%	6	50%	5	45%	133	55%
Other/unknown	1	8%	4	25%	5	36%	6	50%	6	55%	68	28%
Subtotal	12	100%	16	100%	14	100%	12	100%	11	100%	243	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	15	68%
Other/unknown	-	-	-	-	-	-	-	-	-	-	7	32%
Subtotal	-	-	-	-	-	-	-	-	-	-	22	100%
Total	59		64		52		48		57		1,669	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 83. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 8A

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	87.9	390	80%	195	80%	195	81%
Females	21.4	96	20%	49	20%	47	19%
Age at diagnosis (yr)							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	7.9	5	1%	1	<1%	4	2%
20-24	32.6	17	3%	13	5%	4	2%
25-29	40.7	20	4%	15	6%	5	2%
30-34	87.6	47	10%	27	11%	20	8%
35-39	76.3	43	9%	21	9%	22	9%
40-44	106.2	66	14%	29	12%	37	15%
45-49	157.6	104	21%	52	21%	52	21%
50-54	129.0	88	18%	41	17%	47	19%
55-64	67.0	77	16%	38	16%	39	16%
65+	16.5	20	4%	8	3%	12	5%
Race/Ethnicity							
White, not Hispanic	45.7	371	76%	188	77%	183	76%
Black/African American, not Hispanic	281.8	86	18%	42	17%	44	18%
Hispanic/Latino	90.5	20	4%	11	4%	9	4%
Asian/Pacific Islander	*	3	1%	2	1%	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	7	1%	2	1%	5	2%
Race/Sex							
White, not Hispanic Males	77.2	310	64%	158	64%	152	63%
White, not Hispanic Females	14.9	61	13%	30	12%	31	13%
Black/African American, not Hispanic Males	341.0	57	12%	26	11%	31	13%
Black/African American, not Hispanic Females	210.1	29	6%	16	7%	13	5%
Hispanic/Latino Males	129.2	15	3%	8	3%	7	3%
Hispanic/Latina Females	47.6	5	1%	3	1%	2	1%
Asian/Pacific Islander Males	*	3	1%	2	1%	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	7	1%	2	1%	5	2%
Total	54.5	487		245		242	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 84. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 8A

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	254	66%	124	64%	130	67%
Injection drug use (IDU)	14	4%	5	3%	9	5%
Male-to-male sexual contact and IDU	15	4%	9	5%	6	3%
Heterosexual contact	26	7%	9	5%	17	9%
Other/unknown	78	20%	46	24%	32	16%
Subtotal	387	100%	193	100%	194	100%
Female adult or adolescent						
Injection drug use	10	11%	3	6%	7	16%
Heterosexual contact	58	63%	29	59%	29	67%
Other/unknown	24	26%	17	35%	7	16%
Subtotal	92	100%	49	100%	43	100%
Child (<13 yrs at diagnosis)						
Perinatal	6	86%	1	50%	5	100%
Other/unknown	1	14%	1	50%	-	-
Subtotal	7	100%	2	100%	5	100%
Total	487		245		242	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 85. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 8A, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	294	78%	313	78%	337	79%	366	80%	390	80%
Females	82	22%	86	22%	89	21%	92	20%	96	20%
Age at diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	4	1%	4	1%	4	1%	4	1%	5	1%
20-24	3	1%	3	1%	6	1%	12	3%	17	3%
25-29	8	2%	9	2%	14	3%	19	4%	20	4%
30-34	26	7%	33	8%	38	9%	43	9%	47	10%
35-39	32	8%	35	9%	37	9%	39	8%	43	9%
40-44	55	15%	58	15%	59	14%	62	14%	66	14%
45-49	87	23%	89	22%	95	22%	97	21%	104	21%
50-54	73	19%	78	20%	82	19%	87	19%	88	18%
55-64	70	19%	72	18%	72	17%	76	17%	77	16%
65+	19	5%	19	5%	20	5%	20	4%	20	4%
Race/Ethnicity										
White, not Hispanic	284	75%	304	76%	324	76%	351	76%	371	76%
Black/African American, not Hispanic	69	18%	71	18%	77	18%	81	18%	86	18%
Hispanic/Latino	16	4%	17	4%	18	4%	18	4%	20	4%
Asian/Pacific Islander	2	1%	2	1%	2	<1%	2	<1%	3	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	6	2%	6	2%	6	1%	7	2%	7	1%
Race/Sex										
White, not Hispanic Males	234	62%	250	63%	268	63%	292	64%	310	64%
White, not Hispanic Females	50	13%	54	14%	56	13%	59	13%	61	13%
Black/African American, not Hispanic Males	43	11%	45	11%	50	12%	54	12%	57	12%
Black/African American, not Hispanic Females	26	7%	26	7%	27	6%	27	6%	29	6%
Hispanic/Latino Males	11	3%	12	3%	13	3%	13	3%	15	3%
Hispanic/Latina Females	5	1%	5	1%	5	1%	5	1%	5	1%
Asian/Pacific Islander Males	2	1%	2	1%	2	<1%	2	<1%	3	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	6	2%	6	2%	6	1%	7	2%	7	1%
Total	377		400		427		459		487	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 86. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 8A, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	193	66%	206	66%	225	67%	241	66%	254	66%
Injection drug use (IDU)	13	4%	13	4%	13	4%	13	4%	14	4%
Male-to-male sexual contact and IDU	14	5%	14	5%	14	4%	15	4%	15	4%
Heterosexual contact	18	6%	22	7%	24	7%	25	7%	26	7%
Other/unknown	53	18%	55	18%	58	17%	69	19%	78	20%
Subtotal	291	100%	310	100%	334	100%	363	100%	387	100%
Female adult or adolescent										
Injection drug use	8	10%	9	11%	10	12%	10	11%	10	11%
Heterosexual contact	49	63%	52	63%	53	62%	56	64%	58	63%
Other/unknown	21	27%	21	26%	22	26%	22	25%	24	26%
Subtotal	78	100%	82	100%	85	100%	88	100%	92	100%
Child (<13 yrs at diagnosis)										
Perinatal	6	86%	6	86%	6	86%	6	86%	6	86%
Other/unknown	1	14%	1	14%	1	14%	1	14%	1	14%
Subtotal	7	100%	7	100%	7	100%	7	100%	7	100%
Total	377		400		427		459		487	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 87. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 8A, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	5.6	25	96%	4.5	20	77%	5.9	26	90%	6.3	28	90%	5.6	25	86%
Females	*	1	4%	1.3	6	23%	*	3	10%	*	3	10%	*	4	14%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	1	3%	*	1	3%	*	1	3%
20-24	*	3	12%	*	1	4%	*	3	10%	11.5	6	19%	11.5	6	21%
25-29	12.2	6	23%	10.2	5	19%	14.2	7	24%	12.2	6	19%	*	1	3%
30-34	*	3	12%	9.3	5	19%	*	3	10%	*	3	10%	*	4	14%
35-39	*	4	15%	*	3	12%	*	1	3%	*	3	10%	*	4	14%
40-44	*	4	15%	*	2	8%	11.3	7	24%	*	2	6%	11.3	7	24%
45-49	*	1	4%	10.6	7	27%	*	4	14%	7.6	5	16%	*	4	14%
50-54	*	4	15%	*	2	8%	*	1	3%	*	3	10%	*	1	3%
55-64	*	1	4%	*	1	4%	*	2	7%	*	2	6%	*	1	3%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity															
White, not Hispanic	2.7	22	85%	2.5	20	77%	2.6	21	72%	3.3	27	87%	2.6	21	72%
Black/African American, not Hispanic	*	2	8%	*	4	15%	22.9	7	24%	*	3	10%	16.4	5	17%
Hispanic/Latino	*	1	4%	*	2	8%	*	1	3%	*	-	-	*	2	7%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	-	-	*	-	-	*	1	3%	*	-	-
Race/Sex															
White, not Hispanic Males	5.2	21	81%	4.0	16	62%	4.7	19	66%	6.0	24	77%	4.7	19	66%
White, not Hispanic Females	*	1	4%	*	4	15%	*	2	7%	*	3	10%	*	2	7%
Black/African American, not Hispanic Males	*	2	8%	*	3	12%	35.9	6	21%	*	3	10%	*	3	10%
Black/African American, not Hispanic Females	*	-	-	*	1	4%	*	1	3%	*	-	-	*	2	7%
Hispanic/Latino Males	*	1	4%	*	1	4%	*	1	3%	*	-	-	*	2	7%
Hispanic/Latina Females	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	-	-	*	-	-	*	1	3%	*	-	-
Total	2.9	26		2.9	26		3.2	29		3.5	31		3.2	29	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 88. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 8A, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	20	80%	12	60%	21	81%	15	54%	13	52%
Injection drug use (IDU)	-	-	2	10%	-	-	-	-	1	4%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	4%	-	-
Heterosexual contact	1	4%	4	20%	2	8%	1	4%	1	4%
Other/unknown	4	16%	2	10%	3	12%	11	39%	10	40%
Subtotal	25	100%	20	100%	26	100%	28	100%	25	100%
Female adult or adolescent										
Injection drug use	-	-	2	33%	1	33%	-	-	-	-
Heterosexual contact	1	100%	3	50%	1	33%	3	100%	2	50%
Other/unknown	-	-	1	17%	1	33%	-	-	2	50%
Subtotal	1	100%	6	100%	3	100%	3	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	26		26		29		31		29	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 89. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 8A, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	10	77%	14	74%	7	47%	13	87%	13	87%	423	85%
Females	3	23%	5	26%	8	53%	2	13%	2	13%	75	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	1	7%	5	1%
20-24	-	-	1	5%	1	7%	-	-	1	7%	15	3%
25-29	1	8%	3	16%	2	13%	4	27%	3	20%	73	15%
30-34	2	15%	3	16%	2	13%	3	20%	1	7%	95	19%
35-39	1	8%	2	11%	-	-	1	7%	3	20%	92	18%
40-44	1	8%	3	16%	4	27%	1	7%	2	13%	70	14%
45-49	2	15%	5	26%	4	27%	4	27%	1	7%	71	14%
50-54	5	38%	1	5%	1	7%	2	13%	1	7%	38	8%
55-64	1	8%	1	5%	1	7%	-	-	2	13%	26	5%
65+	-	-	-	-	-	-	-	-	-	-	7	1%
Race/Ethnicity												
White, not Hispanic	10	77%	13	68%	10	67%	13	87%	11	73%	418	84%
Black/African American, not Hispanic	2	15%	6	32%	4	27%	-	-	3	20%	58	12%
Hispanic/Latino	-	-	-	-	-	-	-	-	1	7%	14	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	1	8%	-	-	1	7%	2	13%	-	-	6	1%
Race/Sex												
White, not Hispanic Males	9	69%	10	53%	5	33%	11	73%	10	67%	370	74%
White, not Hispanic Females	1	8%	3	16%	5	33%	2	13%	1	7%	48	10%
Black/African American, not Hispanic Males	1	8%	4	21%	2	13%	-	-	2	13%	36	7%
Black/African American, not Hispanic Females	1	8%	2	11%	2	13%	-	-	1	7%	22	4%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	1	7%	11	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	3	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	8%	-	-	1	7%	2	13%	-	-	6	1%
Total	13		19		15		15		15		498	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8A

Table 90. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 8A, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	7	70%	9	64%	6	86%	8	62%	8	62%	281	67%
Injection drug use (IDU)	-	-	1	7%	1	14%	-	-	1	8%	20	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	18	4%
Heterosexual contact	1	10%	2	14%	-	-	1	8%	-	-	26	6%
Other/unknown	2	20%	2	14%	-	-	4	31%	4	31%	76	18%
Subtotal	10	100%	14	100%	7	100%	13	100%	13	100%	421	100%
Female adult or adolescent												
Injection drug use	1	33%	1	20%	1	13%	-	-	-	-	14	20%
Heterosexual contact	2	67%	3	60%	5	63%	2	100%	1	50%	47	67%
Other/unknown	-	-	1	20%	2	25%	-	-	1	50%	9	13%
Subtotal	3	100%	5	100%	8	100%	2	100%	2	100%	70	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	57%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	43%
Subtotal	-	-	-	-	-	-	-	-	-	-	7	100%
Total	13		19		15		15		15		498	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 91. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 8B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	80.9	448	75%	209	74%	239	76%
Females	25.9	147	25%	73	26%	74	24%
Age at diagnosis (yr)							
<13	*	3	1%	2	1%	1	<1%
13-14	*	3	1%	2	1%	1	<1%
15-19	*	2	<1%	1	<1%	1	<1%
20-24	36.6	24	4%	18	6%	6	2%
25-29	66.1	40	7%	24	9%	16	5%
30-34	79.5	51	9%	26	9%	25	8%
35-39	90.4	60	10%	32	11%	28	9%
40-44	113.2	85	14%	43	15%	42	13%
45-49	129.9	106	18%	41	15%	65	21%
50-54	113.7	99	17%	38	13%	61	19%
55-64	57.8	88	15%	37	13%	51	16%
65+	20.2	34	6%	18	6%	16	5%
Race/Ethnicity							
White, not Hispanic	38.1	382	64%	192	68%	190	61%
Black/African American, not Hispanic	250.6	124	21%	48	17%	76	24%
Hispanic/Latino	182.5	75	13%	38	13%	37	12%
Asian/Pacific Islander	*	3	1%	1	<1%	2	1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	10	2%	3	1%	7	2%
Race/Sex							
White, not Hispanic Males	61.1	301	51%	150	53%	151	48%
White, not Hispanic Females	15.9	81	14%	42	15%	39	12%
Black/African American, not Hispanic Males	326.8	89	15%	31	11%	58	19%
Black/African American, not Hispanic Females	157.4	35	6%	17	6%	18	6%
Hispanic/Latino Males	244.3	51	9%	26	9%	25	8%
Hispanic/Latina Females	118.7	24	4%	12	4%	12	4%
Asian/Pacific Islander Males	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	10	2%	3	1%	7	2%
Total	53.0	595		282		313	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 92. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 8B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	279	63%	123	59%	156	66%
Injection drug use (IDU)	22	5%	13	6%	9	4%
Male-to-male sexual contact and IDU	24	5%	5	2%	19	8%
Heterosexual contact	25	6%	8	4%	17	7%
Other/unknown	94	21%	58	28%	36	15%
Subtotal	444	100%	207	100%	237	100%
Female adult or adolescent						
Injection drug use	10	7%	6	8%	4	6%
Heterosexual contact	103	72%	50	70%	53	74%
Other/unknown	30	21%	15	21%	15	21%
Subtotal	143	100%	71	100%	72	100%
Child (<13 yrs at diagnosis)						
Perinatal	7	88%	3	75%	4	100%
Other/unknown	1	13%	1	25%	-	-
Subtotal	8	100%	4	100%	4	100%
Total	595		282		313	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 93. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 8B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	358	75%	376	75%	401	75%	422	75%	448	75%
Females	120	25%	127	25%	134	25%	138	25%	147	25%
Age at diagnosis (yr)										
<13	2	<1%	2	<1%	3	1%	3	1%	3	1%
13-14	3	1%	3	1%	3	1%	3	1%	3	1%
15-19	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
20-24	2	<1%	5	1%	11	2%	15	3%	24	4%
25-29	23	5%	28	6%	30	6%	36	6%	40	7%
30-34	41	9%	42	8%	45	8%	48	9%	51	9%
35-39	40	8%	43	9%	47	9%	54	10%	60	10%
40-44	76	16%	78	16%	82	15%	83	15%	85	14%
45-49	89	19%	93	18%	100	19%	102	18%	106	18%
50-54	92	19%	95	19%	97	18%	98	18%	99	17%
55-64	78	16%	81	16%	83	16%	83	15%	88	15%
65+	31	6%	32	6%	33	6%	33	6%	34	6%
Race/Ethnicity										
White, not Hispanic	312	65%	325	65%	341	64%	357	64%	382	64%
Black/African American, not Hispanic	96	20%	103	20%	112	21%	119	21%	124	21%
Hispanic/Latino	60	13%	62	12%	69	13%	71	13%	75	13%
Asian/Pacific Islander	2	<1%	2	<1%	2	<1%	2	<1%	3	1%
American Indian/Alaska Native	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	8	2%	10	2%	10	2%	10	2%	10	2%
Race/Sex										
White, not Hispanic Males	246	51%	258	51%	270	50%	284	51%	301	51%
White, not Hispanic Females	66	14%	67	13%	71	13%	73	13%	81	14%
Black/African American, not Hispanic Males	67	14%	71	14%	78	15%	84	15%	89	15%
Black/African American, not Hispanic Females	29	6%	32	6%	34	6%	35	6%	35	6%
Hispanic/Latino Males	39	8%	40	8%	46	9%	47	8%	51	9%
Hispanic/Latina Females	21	4%	22	4%	23	4%	24	4%	24	4%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	8	2%	10	2%	10	2%	10	2%	10	2%
Total	478		503		535		560		595	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 94. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 8B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	223	63%	235	63%	252	63%	264	63%	279	63%
Injection drug use (IDU)	21	6%	21	6%	22	6%	22	5%	22	5%
Male-to-male sexual contact and IDU	19	5%	20	5%	20	5%	22	5%	24	5%
Heterosexual contact	21	6%	22	6%	23	6%	23	6%	25	6%
Other/unknown	70	20%	74	20%	80	20%	87	21%	94	21%
Subtotal	354	100%	372	100%	397	100%	418	100%	444	100%
Female adult or adolescent										
Injection drug use	7	6%	7	6%	7	5%	7	5%	10	7%
Heterosexual contact	84	72%	90	73%	96	74%	99	74%	103	72%
Other/unknown	26	22%	27	22%	27	21%	28	21%	30	21%
Subtotal	117	100%	124	100%	130	100%	134	100%	143	100%
Child (<13 yrs at diagnosis)										
Perinatal	7	100%	7	100%	7	88%	7	88%	7	88%
Other/unknown	-	-	-	-	1	13%	1	13%	1	13%
Subtotal	7	100%	7	100%	8	100%	8	100%	8	100%
Total	478		503		535		560		595	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 95. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 8B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	5.2	29	81%	3.2	18	75%	4.5	25	78%	4.2	23	85%	4.7	26	74%
Females	1.2	7	19%	1.1	6	25%	1.2	7	22%	*	4	15%	1.6	9	26%
Age at diagnosis (yr)															
<13	*	1	3%	*	-	-	*	1	3%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	4%	*	3	9%	*	1	4%	*	1	3%
20-24	7.6	5	14%	7.6	5	21%	7.6	5	16%	9.2	6	22%	13.7	9	26%
25-29	*	3	8%	*	-	-	*	3	9%	8.3	5	19%	9.9	6	17%
30-34	*	3	8%	*	2	8%	*	3	9%	9.4	6	22%	*	2	6%
35-39	12.0	8	22%	*	3	13%	*	3	9%	7.5	5	19%	*	4	11%
40-44	13.3	10	28%	6.7	5	21%	6.7	5	16%	*	-	-	*	2	6%
45-49	6.1	5	14%	*	3	13%	7.4	6	19%	*	2	7%	6.1	5	14%
50-54	*	-	-	*	3	13%	*	-	-	*	1	4%	*	2	6%
55-64	*	1	3%	*	2	8%	*	2	6%	*	1	4%	*	3	9%
65+	*	-	-	*	-	-	*	1	3%	*	-	-	*	1	3%
Race/Ethnicity															
White, not Hispanic	2.4	24	67%	1.4	14	58%	1.7	17	53%	1.8	18	67%	2.5	25	71%
Black/African American, not Hispanic	14.1	7	19%	12.1	6	25%	16.2	8	25%	14.1	7	26%	10.1	5	14%
Hispanic/Latino	12.2	5	14%	*	2	8%	17.0	7	22%	*	2	7%	*	4	11%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-
Race/Sex															
White, not Hispanic Males	3.7	18	50%	2.4	12	50%	2.6	13	41%	3.2	16	59%	3.4	17	49%
White, not Hispanic Females	1.2	6	17%	*	2	8%	*	4	13%	*	2	7%	1.6	8	23%
Black/African American, not Hispanic Males	25.7	7	19%	*	4	17%	22.0	6	19%	22.0	6	22%	18.4	5	14%
Black/African American, not Hispanic Females	*	-	-	*	2	8%	*	2	6%	*	1	4%	*	-	-
Hispanic/Latino Males	*	4	11%	*	1	4%	28.7	6	19%	*	1	4%	*	4	11%
Hispanic/Latina Females	*	1	3%	*	1	4%	*	1	3%	*	1	4%	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-
Total	3.2	36		2.1	24		2.9	32		2.4	27		3.1	35	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 96. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 8B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	20	71%	11	61%	16	64%	14	61%	15	58%
Injection drug use (IDU)	-	-	-	-	1	4%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	6%	-	-	2	9%	2	8%
Heterosexual contact	5	18%	1	6%	1	4%	-	-	2	8%
Other/unknown	3	11%	5	28%	7	28%	7	30%	7	27%
Subtotal	28	100%	18	100%	25	100%	23	100%	26	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	3	33%
Heterosexual contact	4	57%	4	67%	6	100%	3	75%	4	44%
Other/unknown	3	43%	2	33%	-	-	1	25%	2	22%
Subtotal	7	100%	6	100%	6	100%	4	100%	9	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-
Subtotal	1	100%	-	-	1	100%	-	-	-	-
Total	36		24		32		27		35	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 97. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 8B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	13	68%	11	69%	16	64%	10	77%	10	63%	558	81%
Females	6	32%	5	31%	9	36%	3	23%	6	38%	132	19%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	11	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	2	8%	-	-	-	-	5	1%
20-24	3	16%	-	-	3	12%	2	15%	1	6%	38	6%
25-29	3	16%	2	13%	1	4%	1	8%	6	38%	99	14%
30-34	-	-	2	13%	-	-	4	31%	2	13%	112	16%
35-39	4	21%	3	19%	4	16%	4	31%	1	6%	151	22%
40-44	3	16%	4	25%	5	20%	1	8%	3	19%	116	17%
45-49	5	26%	3	19%	4	16%	-	-	2	13%	82	12%
50-54	1	5%	-	-	4	16%	-	-	1	6%	36	5%
55-64	-	-	2	13%	1	4%	1	8%	-	-	31	4%
65+	-	-	-	-	1	4%	-	-	-	-	7	1%
Race/Ethnicity												
White, not Hispanic	15	79%	9	56%	13	52%	8	62%	9	56%	493	71%
Black/African American, not Hispanic	2	11%	6	38%	8	32%	4	31%	5	31%	117	17%
Hispanic/Latino	2	11%	-	-	4	16%	1	8%	1	6%	71	10%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	1	6%	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	1	6%	5	1%
Race/Sex												
White, not Hispanic Males	10	53%	8	50%	8	32%	6	46%	5	31%	420	61%
White, not Hispanic Females	5	26%	1	6%	5	20%	2	15%	4	25%	73	11%
Black/African American, not Hispanic Males	2	11%	3	19%	5	20%	3	23%	3	19%	89	13%
Black/African American, not Hispanic Females	-	-	3	19%	3	12%	1	8%	2	13%	28	4%
Hispanic/Latino Males	1	5%	-	-	3	12%	1	8%	1	6%	44	6%
Hispanic/Latina Females	1	5%	-	-	1	4%	-	-	-	-	27	4%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	6%	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	1	6%	5	1%
Total	19		16		25		13		16		690	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8B

Table 98. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 8B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	69%	7	64%	10	63%	9	90%	4	40%	354	64%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	41	7%
Male-to-male sexual contact and IDU	-	-	1	9%	-	-	-	-	1	10%	43	8%
Heterosexual contact	2	15%	-	-	1	6%	1	10%	-	-	31	6%
Other/unknown	2	15%	3	27%	5	31%	-	-	5	50%	83	15%
Subtotal	13	100%	11	100%	16	100%	10	100%	10	100%	552	100%
Female adult or adolescent												
Injection drug use	1	17%	-	-	2	22%	-	-	-	-	19	15%
Heterosexual contact	4	67%	3	60%	7	78%	1	33%	3	50%	82	65%
Other/unknown	1	17%	2	40%	-	-	2	67%	3	50%	26	20%
Subtotal	6	100%	5	100%	9	100%	3	100%	6	100%	127	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	8	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	27%
Subtotal	-	-	-	-	-	-	-	-	-	-	11	100%
Total	19		16		25		13		16		690	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 99. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 8C

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	93.0	494	80%	221	78%	273	83%
Females	22.3	120	20%	63	22%	57	17%
Age at diagnosis (yr)							
<13	*	3	<1%	2	1%	1	<1%
13-14	*	1	<1%	1	<1%	-	-
15-19	*	2	<1%	1	<1%	1	<1%
20-24	49.7	35	6%	30	11%	5	2%
25-29	58.4	37	6%	23	8%	14	4%
30-34	80.1	52	8%	35	12%	17	5%
35-39	108.5	70	11%	40	14%	30	9%
40-44	113.6	82	13%	33	12%	49	15%
45-49	150.2	116	19%	33	12%	83	25%
50-54	124.4	102	17%	35	12%	67	20%
55-64	59.5	84	14%	33	12%	51	15%
65+	19.7	30	5%	18	6%	12	4%
Race/Ethnicity							
White, not Hispanic	47.5	464	76%	209	74%	255	77%
Black/African American, not Hispanic	261.2	114	19%	59	21%	55	17%
Hispanic/Latino	93.8	16	3%	4	1%	12	4%
Asian/Pacific Islander	*	3	<1%	2	1%	1	<1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	16	3%	9	3%	7	2%
Race/Sex							
White, not Hispanic Males	78.2	377	61%	162	57%	215	65%
White, not Hispanic Females	17.6	87	14%	47	17%	40	12%
Black/African American, not Hispanic Males	339.7	86	14%	47	17%	39	12%
Black/African American, not Hispanic Females	152.8	28	5%	12	4%	16	5%
Hispanic/Latino Males	165.7	15	2%	4	1%	11	3%
Hispanic/Latina Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	*	3	<1%	2	1%	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	16	3%	9	3%	7	2%
Total	57.5	614		284		330	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 100. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 8C

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	339	69%	145	66%	194	72%
Injection drug use (IDU)	19	4%	7	3%	12	4%
Male-to-male sexual contact and IDU	19	4%	9	4%	10	4%
Heterosexual contact	40	8%	14	6%	26	10%
Other/unknown	72	15%	44	20%	28	10%
Subtotal	489	100%	219	100%	270	100%
Female adult or adolescent						
Injection drug use	5	4%	4	6%	1	2%
Heterosexual contact	77	65%	34	55%	43	77%
Other/unknown	36	31%	24	39%	12	21%
Subtotal	118	100%	62	100%	56	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	57%	3	100%	1	25%
Other/unknown	3	43%	-	-	3	75%
Subtotal	7	100%	3	100%	4	100%
Total	614		284		330	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 101. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 8C, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	357	79%	393	80%	430	81%	460	81%	494	80%
Females	96	21%	100	20%	104	19%	109	19%	120	20%
Age at diagnosis (yr)										
<13	1	<1%	2	<1%	2	<1%	3	1%	3	<1%
13-14	-	-	-	-	-	-	1	<1%	1	<1%
15-19	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
20-24	1	<1%	4	1%	14	3%	25	4%	35	6%
25-29	20	4%	23	5%	31	6%	31	5%	37	6%
30-34	29	6%	34	7%	39	7%	47	8%	52	8%
35-39	39	9%	50	10%	57	11%	60	11%	70	11%
40-44	61	13%	68	14%	70	13%	77	14%	82	13%
45-49	105	23%	110	22%	111	21%	113	20%	116	19%
50-54	89	20%	90	18%	95	18%	97	17%	102	17%
55-64	78	17%	81	16%	84	16%	84	15%	84	14%
65+	28	6%	29	6%	29	5%	29	5%	30	5%
Race/Ethnicity										
White, not Hispanic	348	77%	380	77%	409	77%	432	76%	464	76%
Black/African American, not Hispanic	79	17%	86	17%	96	18%	105	18%	114	19%
Hispanic/Latino	12	3%	12	2%	13	2%	14	2%	16	3%
Asian/Pacific Islander	2	<1%	2	<1%	2	<1%	2	<1%	3	<1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	11	2%	12	2%	13	2%	15	3%	16	3%
Race/Sex										
White, not Hispanic Males	277	61%	307	62%	332	62%	353	62%	377	61%
White, not Hispanic Females	71	16%	73	15%	77	14%	79	14%	87	14%
Black/African American, not Hispanic Males	58	13%	63	13%	73	14%	79	14%	86	14%
Black/African American, not Hispanic Females	21	5%	23	5%	23	4%	26	5%	28	5%
Hispanic/Latino Males	11	2%	11	2%	12	2%	13	2%	15	2%
Hispanic/Latina Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Males	2	<1%	2	<1%	2	<1%	2	<1%	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	11	2%	12	2%	13	2%	15	3%	16	3%
Total	453		493		534		569		614	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 102. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 8C, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	243	69%	272	70%	298	70%	321	71%	339	69%
Injection drug use (IDU)	16	5%	17	4%	17	4%	17	4%	19	4%
Male-to-male sexual contact and IDU	15	4%	16	4%	17	4%	18	4%	19	4%
Heterosexual contact	32	9%	33	8%	39	9%	39	9%	40	8%
Other/unknown	48	14%	51	13%	55	13%	60	13%	72	15%
Subtotal	354	100%	389	100%	426	100%	455	100%	489	100%
Female adult or adolescent										
Injection drug use	1	1%	3	3%	3	3%	3	3%	5	4%
Heterosexual contact	67	71%	69	70%	73	71%	74	69%	77	65%
Other/unknown	27	28%	27	27%	27	26%	30	28%	36	31%
Subtotal	95	100%	99	100%	103	100%	107	100%	118	100%
Child (<13 yrs at diagnosis)										
Perinatal	2	50%	2	40%	2	40%	4	57%	4	57%
Other/unknown	2	50%	3	60%	3	60%	3	43%	3	43%
Subtotal	4	100%	5	100%	5	100%	7	100%	7	100%
Total	453		493		534		569		614	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 103. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 8C, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	3.4	18	60%	7.0	37	90%	8.5	45	92%	5.6	30	86%	6.6	35	76%
Females	2.2	12	40%	*	4	10%	*	4	8%	0.9	5	14%	2.0	11	24%
Age at diagnosis (yr)															
<13	*	-	-	*	1	2%	*	-	-	*	2	6%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	3	7%	*	4	8%	6.8	5	14%	*	1	2%
20-24	*	4	13%	*	3	7%	17.0	12	24%	8.5	6	17%	17.0	12	26%
25-29	7.9	5	17%	*	4	10%	12.6	8	16%	*	4	11%	*	4	9%
30-34	7.7	5	17%	17.0	11	27%	9.2	6	12%	9.2	6	17%	12.3	8	17%
35-39	*	4	13%	12.4	8	20%	9.3	6	12%	*	4	11%	12.4	8	17%
40-44	*	3	10%	6.9	5	12%	*	1	2%	6.9	5	14%	*	4	9%
45-49	6.5	5	17%	*	2	5%	*	4	8%	*	1	3%	7.8	6	13%
50-54	*	2	7%	*	2	5%	*	3	6%	*	2	6%	*	2	4%
55-64	*	1	3%	*	1	2%	3.5	5	10%	*	-	-	*	-	-
65+	*	1	3%	*	1	2%	*	-	-	*	-	-	*	1	2%
Race/Ethnicity															
White, not Hispanic	2.5	24	80%	3.4	33	80%	3.6	35	71%	2.4	23	66%	3.4	33	72%
Black/African American, not Hispanic	13.7	6	20%	16.0	7	17%	25.2	11	22%	20.6	9	26%	20.6	9	20%
Hispanic/Latino	*	-	-	*	-	-	*	2	4%	*	1	3%	*	2	4%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	2%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	2%	*	1	2%	*	2	6%	*	1	2%
Race/Sex															
White, not Hispanic Males	2.9	14	47%	6.4	31	76%	6.4	31	63%	4.4	21	60%	5.2	25	54%
White, not Hispanic Females	2.0	10	33%	*	2	5%	*	4	8%	*	2	6%	1.6	8	17%
Black/African American, not Hispanic Males	*	4	13%	19.8	5	12%	43.5	11	22%	23.7	6	17%	27.7	7	15%
Black/African American, not Hispanic Females	*	2	7%	*	2	5%	*	-	-	*	3	9%	*	2	4%
Hispanic/Latino Males	*	-	-	*	-	-	*	2	4%	*	1	3%	*	2	4%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	1	2%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	2%	*	1	2%	*	2	6%	*	1	2%
Total	2.8	30		3.8	41		4.6	49		3.3	35		4.3	46	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 104. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 8C, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	12	67%	30	83%	31	69%	23	79%	18	51%
Injection drug use (IDU)	-	-	1	3%	-	-	-	-	2	6%
Male-to-male sexual contact and IDU	-	-	1	3%	1	2%	1	3%	1	3%
Heterosexual contact	5	28%	1	3%	6	13%	-	-	1	3%
Other/unknown	1	6%	3	8%	7	16%	5	17%	13	37%
Subtotal	18	100%	36	100%	45	100%	29	100%	35	100%
Female adult or adolescent										
Injection drug use	-	-	2	50%	-	-	-	-	2	18%
Heterosexual contact	10	83%	2	50%	4	100%	1	25%	3	27%
Other/unknown	2	17%	-	-	-	-	3	75%	6	55%
Subtotal	12	100%	4	100%	4	100%	4	100%	11	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	2	100%	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	2	100%	-	-
Total	30		41		49		35		46	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 105. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 8C, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	23	79%	22	88%	22	88%	16	80%	13	72%	671	87%
Females	6	21%	3	12%	3	12%	4	20%	5	28%	100	13%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	5%	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	2	10%	-	-	8	1%
20-24	2	7%	1	4%	2	8%	2	10%	3	17%	35	5%
25-29	5	17%	1	4%	5	20%	2	10%	1	6%	101	13%
30-34	5	17%	2	8%	-	-	4	20%	4	22%	176	23%
35-39	3	10%	9	36%	3	12%	1	5%	2	11%	141	18%
40-44	7	24%	1	4%	8	32%	4	20%	3	17%	143	19%
45-49	2	7%	6	24%	4	16%	2	10%	2	11%	74	10%
50-54	2	7%	4	16%	2	8%	2	10%	2	11%	45	6%
55-64	2	7%	1	4%	1	4%	-	-	1	6%	28	4%
65+	1	3%	-	-	-	-	-	-	-	-	13	2%
Race/Ethnicity												
White, not Hispanic	24	83%	21	84%	16	64%	12	60%	17	94%	629	82%
Black/African American, not Hispanic	3	10%	3	12%	6	24%	6	30%	1	6%	117	15%
Hispanic/Latino	1	3%	-	-	1	4%	1	5%	-	-	14	2%
Asian/Pacific Islander	1	3%	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	4%	2	8%	1	5%	-	-	8	1%
Race/Sex												
White, not Hispanic Males	19	66%	18	72%	14	56%	10	50%	13	72%	554	72%
White, not Hispanic Females	5	17%	3	12%	2	8%	2	10%	4	22%	75	10%
Black/African American, not Hispanic Males	2	7%	3	12%	5	20%	4	20%	-	-	94	12%
Black/African American, not Hispanic Females	1	3%	-	-	1	4%	2	10%	1	6%	23	3%
Hispanic/Latino Males	1	3%	-	-	1	4%	1	5%	-	-	13	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	1	3%	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	4%	2	8%	1	5%	-	-	8	1%
Total	29		25		25		20		18		771	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8C

Table 106. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 8C, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	15	65%	15	68%	15	68%	11	73%	8	62%	461	69%
Injection drug use (IDU)	1	4%	-	-	-	-	-	-	-	-	41	6%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	28	4%
Heterosexual contact	5	22%	5	23%	2	9%	-	-	1	8%	37	6%
Other/unknown	2	9%	2	9%	5	23%	4	27%	4	31%	98	15%
Subtotal	23	100%	22	100%	22	100%	15	100%	13	100%	665	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	1	20%	11	11%
Heterosexual contact	5	83%	3	100%	3	100%	1	25%	3	60%	61	63%
Other/unknown	1	17%	-	-	-	-	3	75%	1	20%	25	26%
Subtotal	6	100%	3	100%	3	100%	4	100%	5	100%	97	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	44%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	5	56%
Subtotal	-	-	-	-	-	-	1	100%	-	-	9	100%
Total	29		25		25		20		18		771	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 107. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Region 8D

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	90.1	364	76%	163	73%	201	79%
Females	27.7	114	24%	61	27%	53	21%
Age at diagnosis (yr)							
<13	*	3	1%	2	1%	1	<1%
13-14	*	-	-	-	-	-	-
15-19	*	1	<1%	-	-	1	<1%
20-24	33.9	20	4%	14	6%	6	2%
25-29	43.8	20	4%	17	8%	3	1%
30-34	76.5	36	8%	22	10%	14	6%
35-39	73.7	35	7%	18	8%	17	7%
40-44	135.6	73	15%	34	15%	39	15%
45-49	201.2	117	24%	47	21%	70	28%
50-54	105.2	66	14%	24	11%	42	17%
55-64	81.3	89	19%	41	18%	48	19%
65+	15.0	18	4%	5	2%	13	5%
Race/Ethnicity							
White, not Hispanic	52.0	394	82%	181	81%	213	84%
Black/African American, not Hispanic	219.8	59	12%	29	13%	30	12%
Hispanic/Latino	146.0	14	3%	6	3%	8	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	11	2%	8	4%	3	1%
Race/Sex							
White, not Hispanic Males	82.0	306	64%	134	60%	172	68%
White, not Hispanic Females	22.9	88	18%	47	21%	41	16%
Black/African American, not Hispanic Males	265.2	40	8%	19	8%	21	8%
Black/African American, not Hispanic Females	161.6	19	4%	10	4%	9	4%
Hispanic/Latino Males	205.6	10	2%	5	2%	5	2%
Hispanic/Latina Females	*	4	1%	1	<1%	3	1%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	11	2%	8	4%	3	1%
Total	58.6	478		224		254	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 108. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Region 8D

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	225	63%	100	61%	125	63%
Injection drug use (IDU)	13	4%	2	1%	11	6%
Male-to-male sexual contact and IDU	12	3%	5	3%	7	4%
Heterosexual contact	36	10%	13	8%	23	12%
Other/unknown	74	21%	43	26%	31	16%
Subtotal	360	100%	163	100%	197	100%
Female adult or adolescent						
Injection drug use	12	11%	5	9%	7	13%
Heterosexual contact	63	58%	27	48%	36	68%
Other/unknown	34	31%	24	43%	10	19%
Subtotal	109	100%	56	100%	53	100%
Child (<13 yrs at diagnosis)						
Perinatal	7	78%	5	100%	2	50%
Other/unknown	2	22%	-	-	2	50%
Subtotal	9	100%	5	100%	4	100%
Total	478		224		254	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 109. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Region 8D, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	282	75%	292	75%	311	75%	339	76%	364	76%
Females	96	25%	98	25%	103	25%	110	24%	114	24%
Age at diagnosis (yr)										
<13	2	1%	2	1%	3	1%	3	1%	3	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
20-24	4	1%	6	2%	7	2%	12	3%	20	4%
25-29	4	1%	5	1%	9	2%	16	4%	20	4%
30-34	22	6%	23	6%	25	6%	31	7%	36	8%
35-39	27	7%	30	8%	33	8%	35	8%	35	7%
40-44	58	15%	58	15%	64	15%	70	16%	73	15%
45-49	102	27%	104	27%	108	26%	114	25%	117	24%
50-54	61	16%	62	16%	62	15%	64	14%	66	14%
55-64	81	21%	82	21%	84	20%	85	19%	89	19%
65+	16	4%	17	4%	18	4%	18	4%	18	4%
Race/Ethnicity										
White, not Hispanic	318	84%	326	84%	345	83%	368	82%	394	82%
Black/African American, not Hispanic	42	11%	46	12%	49	12%	57	13%	59	12%
Hispanic/Latino	9	2%	9	2%	11	3%	13	3%	14	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	9	2%	9	2%	9	2%	11	2%	11	2%
Race/Sex										
White, not Hispanic Males	242	64%	248	64%	264	64%	284	63%	306	64%
White, not Hispanic Females	76	20%	78	20%	81	20%	84	19%	88	18%
Black/African American, not Hispanic Males	28	7%	32	8%	34	8%	38	8%	40	8%
Black/African American, not Hispanic Females	14	4%	14	4%	15	4%	19	4%	19	4%
Hispanic/Latino Males	6	2%	6	2%	7	2%	9	2%	10	2%
Hispanic/Latina Females	3	1%	3	1%	4	1%	4	1%	4	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	9	2%	9	2%	9	2%	11	2%	11	2%
Total	378		390		414		449		478	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 110. Reported persons living with a diagnosis of HIV infection by year and transmission category, Region 8D, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	171	61%	178	62%	190	62%	210	63%	225	63%
Injection drug use (IDU)	13	5%	13	4%	13	4%	13	4%	13	4%
Male-to-male sexual contact and IDU	10	4%	10	3%	11	4%	12	4%	12	3%
Heterosexual contact	31	11%	32	11%	33	11%	34	10%	36	10%
Other/unknown	54	19%	56	19%	60	20%	66	20%	74	21%
Subtotal	279	100%	289	100%	307	100%	335	100%	360	100%
Female adult or adolescent										
Injection drug use	10	11%	10	11%	10	10%	10	10%	12	11%
Heterosexual contact	53	58%	55	59%	59	60%	61	58%	63	58%
Other/unknown	28	31%	28	30%	29	30%	34	32%	34	31%
Subtotal	91	100%	93	100%	98	100%	105	100%	109	100%
Child (<13 yrs at diagnosis)										
Perinatal	6	75%	6	75%	7	78%	7	78%	7	78%
Other/unknown	2	25%	2	25%	2	22%	2	22%	2	22%
Subtotal	8	100%	8	100%	9	100%	9	100%	9	100%
Total	378		390		414		449		478	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 111. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Region 8D, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	5.0	20	83%	3.0	12	80%	4.7	19	76%	6.7	27	79%	6.2	25	86%
Females	*	4	17%	*	3	20%	1.5	6	24%	1.7	7	21%	*	4	14%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	1	3%	*	1	3%
20-24	*	1	4%	*	3	20%	*	3	12%	15.3	9	26%	15.3	9	31%
25-29	*	3	13%	*	1	7%	*	3	12%	*	3	9%	*	3	10%
30-34	*	-	-	*	3	20%	*	2	8%	10.6	5	15%	*	4	14%
35-39	10.5	5	21%	*	1	7%	10.5	5	20%	*	4	12%	*	2	7%
40-44	13.0	7	29%	*	4	27%	11.1	6	24%	16.7	9	26%	*	1	3%
45-49	*	4	17%	*	-	-	*	1	4%	*	1	3%	*	3	10%
50-54	*	2	8%	*	2	13%	*	2	8%	*	1	3%	*	3	10%
55-64	*	1	4%	*	-	-	*	2	8%	*	1	3%	*	3	10%
65+	*	1	4%	*	1	7%	*	-	-	*	-	-	*	-	-
Race/Ethnicity															
White, not Hispanic	2.6	20	83%	1.5	11	73%	2.5	19	76%	2.9	22	65%	3.4	26	90%
Black/African American, not Hispanic	*	3	13%	*	4	27%	*	4	16%	29.8	8	24%	*	2	7%
Hispanic/Latino	*	1	4%	*	-	-	*	2	8%	*	2	6%	*	1	3%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	2	6%	*	-	-
Race/Sex															
White, not Hispanic Males	4.8	18	75%	2.1	8	53%	4.3	16	64%	5.1	19	56%	5.9	22	76%
White, not Hispanic Females	*	2	8%	*	3	20%	*	3	12%	*	3	9%	*	4	14%
Black/African American, not Hispanic Males	*	1	4%	*	4	27%	*	2	8%	*	4	12%	*	2	7%
Black/African American, not Hispanic Females	*	2	8%	*	-	-	*	2	8%	*	4	12%	*	-	-
Hispanic/Latino Males	*	1	4%	*	-	-	*	1	4%	*	2	6%	*	1	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	2	6%	*	-	-
Total	2.9	24		1.8	15		3.1	25		4.2	34		3.6	29	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 112. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Region 8D, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	10	50%	9	75%	12	67%	20	74%	15	60%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	10%	-	-	1	6%	-	-	-	-
Heterosexual contact	4	20%	1	8%	1	6%	1	4%	2	8%
Other/unknown	4	20%	2	17%	4	22%	6	22%	8	32%
Subtotal	20	100%	12	100%	18	100%	27	100%	25	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	2	50%
Heterosexual contact	4	100%	2	67%	4	67%	2	29%	2	50%
Other/unknown	-	-	1	33%	2	33%	5	71%	-	-
Subtotal	4	100%	3	100%	6	100%	7	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-
Total	24		15		25		34		29	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 113. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Region 8D, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	22	92%	11	69%	4	44%	14	74%	13	87%	441	86%
Females	2	8%	5	31%	5	56%	5	26%	2	13%	74	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	1	11%	-	-	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	-	-	-	-	-	-	-	-	-	-	3	1%
20-24	-	-	2	13%	1	11%	1	5%	1	7%	19	4%
25-29	3	13%	1	6%	-	-	1	5%	2	13%	57	11%
30-34	-	-	2	13%	-	-	3	16%	3	20%	100	19%
35-39	2	8%	2	13%	1	11%	3	16%	1	7%	100	19%
40-44	10	42%	5	31%	2	22%	7	37%	2	13%	109	21%
45-49	4	17%	2	13%	1	11%	1	5%	2	13%	47	9%
50-54	3	13%	2	13%	1	11%	1	5%	1	7%	34	7%
55-64	1	4%	-	-	2	22%	2	11%	3	20%	33	6%
65+	1	4%	-	-	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	20	83%	14	88%	5	56%	15	79%	15	100%	455	88%
Black/African American, not Hispanic	2	8%	1	6%	3	33%	4	21%	-	-	44	9%
Hispanic/Latino	1	4%	1	6%	1	11%	-	-	-	-	10	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	-	-	-	-	-	-	-	-	5	1%
Race/Sex												
White, not Hispanic Males	18	75%	9	56%	3	33%	13	68%	13	87%	395	77%
White, not Hispanic Females	2	8%	5	31%	2	22%	2	11%	2	13%	60	12%
Black/African American, not Hispanic Males	2	8%	1	6%	1	11%	1	5%	-	-	32	6%
Black/African American, not Hispanic Females	-	-	-	-	2	22%	3	16%	-	-	12	2%
Hispanic/Latino Males	1	4%	1	6%	-	-	-	-	-	-	8	2%
Hispanic/Latina Females	-	-	-	-	1	11%	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	-	-	-	-	-	-	-	-	5	1%
Total	24		16		9		19		15		515	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

HIV Prevention Planning Region 8D

Table 114. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Region 8D, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	12	55%	7	64%	3	100%	8	57%	7	54%	276	64%
Injection drug use (IDU)	-	-	1	9%	-	-	-	-	-	-	23	5%
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-	-	-	19	4%
Heterosexual contact	5	23%	1	9%	-	-	-	-	-	-	30	7%
Other/unknown	4	18%	2	18%	-	-	6	43%	6	46%	86	20%
Subtotal	22	100%	11	100%	3	100%	14	100%	13	100%	434	100%
Female adult or adolescent												
Injection drug use	1	50%	-	-	-	-	-	-	-	-	10	14%
Heterosexual contact	1	50%	5	100%	3	60%	2	40%	2	100%	44	61%
Other/unknown	-	-	-	-	2	40%	3	60%	-	-	18	25%
Subtotal	2	100%	5	100%	5	100%	5	100%	2	100%	72	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	1	100%	-	-	-	-	6	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	33%
Subtotal	-	-	-	-	1	100%	-	-	-	-	9	100%
Total	24		16		9		19		15		515	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

This page is intentionally blank