

**Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Brown**

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Brown**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status				
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS		
				No.	%	No.	%	
<b>Sex</b>								
Males	100.5	22	76%	7	64%	15	83%	
Females	31.5	7	24%	4	36%	3	17%	
<b>Age at end of year</b>								
<13	*	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	
20-24	*	1	3%	1	9%	-	-	
25-29	*	2	7%	2	18%	-	-	
30-34	*	1	3%	-	-	1	6%	
35-39	228.1	6	21%	2	18%	4	22%	
40-44	*	2	7%	2	18%	-	-	
45-49	207.7	6	21%	1	9%	5	28%	
50-54	144.5	5	17%	2	18%	3	17%	
55-64	*	4	14%	-	-	4	22%	
65+	*	2	7%	1	9%	1	6%	
<b>Race/Ethnicity</b>								
White, not Hispanic	58.7	25	86%	9	82%	16	89%	
Black/African American, not Hispanic	*	-	-	-	-	-	-	
Hispanic/Latino	*	2	7%	2	18%	-	-	
Asian/Pacific Islander	*	1	3%	-	-	1	6%	
American Indian/Alaska Native	*	-	-	-	-	-	-	
Unknown	*	1	3%	-	-	1	6%	
<b>Race/Sex</b>								
White, not Hispanic Males	85.3	18	62%	5	45%	13	72%	
White, not Hispanic Females	32.5	7	24%	4	36%	3	17%	
Black/African American, not Hispanic Males	*	-	-	-	-	-	-	
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	
Hispanic/Latino Males	*	2	7%	2	18%	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	1	3%	-	-	1	6%	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	
Unknown	*	1	3%	-	-	1	6%	
<b>Total</b>	<b>65.7</b>	<b>29</b>		<b>11</b>		<b>18</b>		

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Brown**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	14	64%	4	57%	10	67%
Injection drug use (IDU)	2	9%	2	29%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	4	18%	-	-	4	27%
Other/unknown	2	9%	1	14%	1	7%
Subtotal	22	100%	7	100%	15	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	43%	1	25%	2	67%
Heterosexual contact	3	43%	2	50%	1	33%
Other/unknown	1	14%	1	25%	-	-
Subtotal	7	100%	4	100%	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>29</b>		<b>11</b>		<b>18</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Brown**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	14	48%	4	36%	10	56%
Injection drug use (IDU) only	2	7%	2	18%	-	-
Heterosexual contact only	7	24%	2	18%	5	28%
MSM & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	3	10%	1	9%	2	11%
MSM & Heterosexual contact	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	3	10%	2	18%	1	6%
<b>Total</b>	<b>29</b>		<b>11</b>		<b>18</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Brown**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	12	67%	-	-	<10	*	<10	*	-	-	-	-
Injection drug use (IDU)	1	6%	-	-	<10	*	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	17%	-	-	-	-	-	-	-	-	<10	*
Other/unknown	2	11%	-	-	-	-	-	-	-	-	-	-
Subtotal	18	100%	-	-	<10	*	<10	*	-	-	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	3	43%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	43%	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	14%	-	-	-	-	-	-	-	-	-	-
Subtotal	7	100%	-	-	-	-	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>25</b>		<b>0</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Brown**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	48%	-	-	<10	*	<10	*	-	-	-	-
Injection drug use (IDU) only	2	8%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	6	24%	-	-	-	-	-	-	-	-	<10	*
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	8%	-	-	<10	*	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	12%	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>25</b>		<b>0</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Brown**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	15	71%	19	73%	20	74%	20	74%	22	76%
Females	6	29%	7	27%	7	26%	7	26%	7	24%
<b>Age at end of year</b>										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	-	-	1	4%	-	-	-	-	1	3%
25-29	1	5%	1	4%	2	7%	2	7%	2	7%
30-34	3	14%	1	4%	-	-	-	-	1	3%
35-39	2	10%	5	19%	7	26%	7	26%	6	21%
40-44	3	14%	4	15%	2	7%	1	4%	2	7%
45-49	6	29%	7	27%	9	33%	10	37%	6	21%
50-54	2	10%	3	12%	3	11%	2	7%	5	17%
55-64	3	14%	3	12%	3	11%	3	11%	4	14%
65+	1	5%	1	4%	1	4%	2	7%	2	7%
<b>Race/Ethnicity</b>										
White, not Hispanic	18	86%	22	85%	23	85%	23	85%	25	86%
Black/African American, not Hispanic	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino	1	5%	2	8%	2	7%	2	7%	2	7%
Asian/Pacific Islander	1	5%	1	4%	1	4%	1	4%	1	3%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	1	4%	1	4%	1	4%	1	3%
<b>Race/Sex</b>										
White, not Hispanic Males	12	57%	15	58%	16	59%	16	59%	18	62%
White, not Hispanic Females	6	29%	7	27%	7	26%	7	26%	7	24%
Black/African American, not Hispanic Males	-	-	-	-	-	-	-	-	-	-
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino Males	1	5%	2	8%	2	7%	2	7%	2	7%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	1	5%	1	4%	1	4%	1	4%	1	3%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	1	4%	1	4%	1	4%	1	3%
<b>Total</b>	<b>21</b>		<b>26</b>		<b>27</b>		<b>27</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Brown

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	9	60%	12	63%	13	65%	13	65%	14	64%
Injection drug use (IDU)	-	-	1	5%	1	5%	1	5%	2	9%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	4	27%	4	21%	4	20%	4	20%	4	18%
Other/unknown	2	13%	2	11%	2	10%	2	10%	2	9%
Subtotal	15	100%	19	100%	20	100%	20	100%	22	100%
<b>Female adult or adolescent</b>										
Injection drug use	3	50%	3	43%	3	43%	3	43%	3	43%
Heterosexual contact	2	33%	3	43%	3	43%	3	43%	3	43%
Other/unknown	1	17%	1	14%	1	14%	1	14%	1	14%
Subtotal	6	100%	7	100%	7	100%	7	100%	7	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>21</b>		<b>26</b>		<b>27</b>		<b>27</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Brown

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	9	43%	12	46%	13	48%	13	48%	14	48%
Injection drug use (IDU) only	1	5%	1	4%	1	4%	1	4%	2	7%
Heterosexual contact only	6	29%	7	27%	7	26%	7	26%	7	24%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	10%	3	12%	3	11%	3	11%	3	10%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	14%	3	12%	3	11%	3	11%	3	10%
<b>Total</b>	<b>21</b>		<b>26</b>		<b>27</b>		<b>27</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.