

Diagnoses of HIV and/or AIDS  
Reported in HIV Prevention Planning  
Butler

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Butler

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
<b>Sex</b>									
Males	8.2	15	79%	11	85%	4	67%	-	-
Females	*	4	21%	2	15%	2	33%	-	-
<b>Age at diagnosis (yr)</b>									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	2	11%	2	15%	-	-	-	-
20-24	*	3	16%	3	23%	-	-	-	-
25-29	*	2	11%	1	8%	1	17%	-	-
30-34	*	2	11%	2	15%	-	-	-	-
35-39	*	2	11%	-	-	2	33%	-	-
40-44	*	2	11%	2	15%	-	-	-	-
45-49	*	1	5%	1	8%	-	-	-	-
50-54	*	3	16%	-	-	3	50%	-	-
55-64	*	2	11%	2	15%	-	-	-	-
65+	*	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>									
White, not Hispanic	3.6	11	58%	6	46%	5	83%	-	-
Black/African American, not Hispanic	24.1	7	37%	6	46%	1	17%	-	-
Hispanic/Latino	*	1	5%	1	8%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
<b>Race/Sex</b>									
White, not Hispanic Males	6.6	10	53%	6	46%	4	67%	-	-
White, not Hispanic Females	*	1	5%	-	-	1	17%	-	-
Black/African American, not Hispanic Males	*	4	21%	4	31%	-	-	-	-
Black/African American, not Hispanic Females	*	3	16%	2	15%	1	17%	-	-
Hispanic/Latino Males	*	1	5%	1	8%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
<b>Total</b>	<b>5.1</b>	<b>19</b>		<b>13</b>		<b>6</b>			

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Butler

Transmission Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	8	53%	7	64%	1	25%	-	-
Injection drug use (IDU)	1	7%	-	-	1	25%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	7%	-	-	1	25%	-	-
Other/unknown	5	33%	4	36%	1	25%	-	-
Subtotal	15	100%	11	100%	4	100%	-	-
<b>Female adult or adolescent</b>								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	75%	2	100%	1	50%	-	-
Other/unknown	1	25%	-	-	1	50%	-	-
Subtotal	4	100%	2	100%	2	100%	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
<b>Total</b>	<b>19</b>		<b>13</b>		<b>6</b>		<b>-</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Butler

Exposure Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	8	42%	7	54%	1	17%	-	-
Injection drug use (IDU) only	1	5%	-	-	1	17%	-	-
Heterosexual contact only	4	21%	2	15%	2	33%	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	6	32%	4	31%	2	33%	-	-
<b>Total</b>	<b>19</b>		<b>13</b>		<b>6</b>		<b>-</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Butler

Transmission Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	6	60%	2	50%	-	-	-	-	-	-	-	-
Injection drug use (IDU)	1	10%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	10%	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	20%	2	50%	1	100%	-	-	-	-	-	-
Subtotal	10	100%	4	100%	1	100%	-	-	-	-	-	-
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	3	100%	-	-	-	-	-	-	-	-
Other/unknown	1	100%	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	3	100%	-	-	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>11</b>		<b>7</b>		<b>1</b>		<b>0</b>		<b>0</b>		<b>0</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Butler

Exposure Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	6	55%	2	29%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	1	9%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	9%	3	43%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	27%	2	29%	1	100%	-	-	-	-	-	-
<b>Total</b>	<b>11</b>		<b>7</b>		<b>1</b>		<b>0</b>		<b>0</b>		<b>0</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Butler

### Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate <sup>a</sup>	No.	%									
<b>Sex</b>												
Males	7.6	14	78%	8.2	15	71%	4.9	9	90%	10.4	19	76%
Females	*	4	22%	3.1	6	29%	*	1	10%	3.1	6	24%
<b>Age at diagnosis (yr)</b>												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	1	4%
20-24	*	3	17%	*	4	19%	*	2	20%	*	3	12%
25-29	*	3	17%	*	3	14%	*	2	20%	33.3	7	28%
30-34	*	-	-	*	4	19%	*	1	10%	*	4	16%
35-39	22.5	5	28%	*	2	10%	*	1	10%	31.6	7	28%
40-44	*	2	11%	*	3	14%	*	1	10%	*	-	-
45-49	*	2	11%	*	2	10%	*	2	20%	*	2	8%
50-54	*	1	6%	*	2	10%	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	1	10%	*	1	4%
65+	*	2	11%	*	1	5%	*	-	-	*	-	-
<b>Race/Ethnicity</b>												
White, not Hispanic	2.3	7	39%	2.3	7	33%	1.6	5	50%	3.2	10	40%
Black/African American, not Hispanic	27.5	8	44%	48.2	14	67%	*	3	30%	31.0	9	36%
Hispanic/Latino	*	1	6%	*	-	-	*	1	10%	*	3	12%
Asian/Pacific Islander	*	1	6%	*	-	-	*	-	-	*	1	4%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	6%	*	-	-	*	1	10%	*	2	8%
<b>Race/Sex</b>												
White, not Hispanic Males	4.0	6	33%	4.6	7	33%	*	4	40%	5.9	9	36%
White, not Hispanic Females	*	1	6%	*	-	-	*	1	10%	*	1	4%
Black/African American, not Hispanic Males	42.5	6	33%	56.6	8	38%	*	3	30%	49.6	7	28%
Black/African American, not Hispanic Females	*	2	11%	40.2	6	29%	*	-	-	*	2	8%
Hispanic/Latino Males	*	1	6%	*	-	-	*	1	10%	*	1	4%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	2	8%
Asian/Pacific Islander Males	*	1	6%	*	-	-	*	-	-	*	1	4%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	6%	*	-	-	*	1	10%	*	2	8%
<b>Total</b>	<b>4.8</b>	<b>18</b>		<b>5.6</b>	<b>21</b>		<b>2.7</b>	<b>10</b>		<b>6.7</b>	<b>25</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Butler

### Diagnosis of HIV Infection by Year

Transmission Category <sup>a</sup>	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	7	50%	8	53%	5	56%	14	74%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	7	50%	7	47%	4	44%	5	26%
Subtotal	14	100%	15	100%	9	100%	19	100%
<b>Female adult or adolescent</b>								
Injection drug use	1	25%	-	-	-	-	1	17%
Heterosexual contact	2	50%	5	83%	-	-	4	67%
Other/unknown	1	25%	1	17%	1	100%	1	17%
Subtotal	4	100%	6	100%	1	100%	6	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
<b>Total</b>	<b>18</b>		<b>21</b>		<b>10</b>		<b>25</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Butler

### Diagnosis of HIV Infection by Year

Exposure Category <sup>a</sup>	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	39%	8	38%	5	50%	14	56%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	2	11%	5	24%	-	-	4	16%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	6%	-	-	-	-	1	4%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	8	44%	8	38%	5	50%	6	24%
<b>Total</b>	<b>18</b>		<b>21</b>		<b>10</b>		<b>25</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Butler**

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Sex</b>										
Males	6	75%	5	63%	5	83%	12	75%	11	85%
Females	2	25%	3	38%	1	17%	4	25%	2	15%
<b>Age at HIV diagnosis (yr)</b>										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	1	6%	2	15%
20-24	2	25%	-	-	2	33%	1	6%	3	23%
25-29	1	13%	2	25%	1	17%	3	19%	1	8%
30-34	-	-	2	25%	-	-	4	25%	2	15%
35-39	3	38%	1	13%	1	17%	5	31%	-	-
40-44	1	13%	1	13%	-	-	-	-	2	15%
45-49	1	13%	-	-	2	33%	2	13%	1	8%
50-54	-	-	2	25%	-	-	-	-	-	-
55-64	-	-	-	-	-	-	-	-	2	15%
65+	-	-	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>										
White, not Hispanic	3	38%	3	38%	3	50%	7	44%	6	46%
Black/African American, not Hispanic	3	38%	5	63%	2	33%	5	31%	6	46%
Hispanic/Latino	1	13%	-	-	-	-	1	6%	1	8%
Asian/Pacific Islander	1	13%	-	-	-	-	1	6%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	17%	2	13%	-	-
<b>Race/Sex</b>										
White, not Hispanic Males	2	25%	3	38%	2	33%	6	38%	6	46%
White, not Hispanic Females	1	13%	-	-	1	17%	1	6%	-	-
Black/African American, not Hispanic Males	2	25%	2	25%	2	33%	4	25%	4	31%
Black/African American, not Hispanic Females	1	13%	3	38%	-	-	1	6%	2	15%
Hispanic/Latino Males	1	13%	-	-	-	-	-	-	1	8%
Hispanic/Latina Females	-	-	-	-	-	-	1	6%	-	-
Asian/Pacific Islander Males	1	13%	-	-	-	-	1	6%	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	17%	2	13%	-	-
<b>Total</b>	<b>8</b>		<b>8</b>		<b>6</b>		<b>16</b>		<b>13</b>	

*Note:* Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Butler**

Transmission Category <sup>a</sup>	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	3	50%	1	20%	3	60%	9	75%	7	64%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	50%	4	80%	2	40%	3	25%	4	36%
Subtotal	6	100%	5	100%	5	100%	12	100%	11	100%
<b>Female adult or adolescent</b>										
Injection drug use	1	50%	-	-	-	-	1	25%	-	-
Heterosexual contact	-	-	2	67%	-	-	2	50%	2	100%
Other/unknown	1	50%	1	33%	1	100%	1	25%	-	-
Subtotal	2	100%	3	100%	1	100%	4	100%	2	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>8</b>		<b>8</b>		<b>6</b>		<b>16</b>		<b>13</b>	

*Note:* Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Butler**

Exposure Category <sup>a</sup>	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	3	38%	1	13%	3	50%	9	56%	7	54%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	2	25%	-	-	2	13%	2	15%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	13%	-	-	-	-	1	6%	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	50%	5	63%	3	50%	4	25%	4	31%
<b>Total</b>	<b>8</b>		<b>8</b>		<b>6</b>		<b>16</b>		<b>13</b>	

*Note:* Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Butler**

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Sex</b>												
Males	8	80%	17	85%	4	100%	7	70%	7	70%	249	86%
Females	2	20%	3	15%	-	-	3	30%	3	30%	41	14%
<b>Age at AIDS diagnosis (yr)</b>												
<13	-	-	-	-	-	-	-	-	-	-	2	1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	1	5%	-	-	-	-	-	-	3	1%
20-24	1	10%	5	25%	-	-	1	10%	2	20%	31	11%
25-29	3	30%	2	10%	1	25%	4	40%	1	10%	50	17%
30-34	2	20%	2	10%	2	50%	-	-	1	10%	57	20%
35-39	-	-	3	15%	-	-	2	20%	2	20%	48	17%
40-44	-	-	2	10%	-	-	1	10%	1	10%	34	12%
45-49	1	10%	2	10%	-	-	1	10%	-	-	21	7%
50-54	1	10%	1	5%	-	-	-	-	3	30%	22	8%
55-64	-	-	1	5%	1	25%	1	10%	-	-	16	6%
65+	2	20%	1	5%	-	-	-	-	-	-	5	2%
<b>Race/Ethnicity</b>												
White, not Hispanic	4	40%	8	40%	2	50%	2	20%	9	90%	192	66%
Black/African American, not Hispanic	6	60%	9	45%	1	25%	5	50%	1	10%	79	27%
Hispanic/Latino	-	-	1	5%	1	25%	3	30%	-	-	14	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	10%	-	-	-	-	-	-	5	2%
<b>Race/Sex</b>												
White, not Hispanic Males	4	40%	8	40%	2	50%	2	20%	7	70%	180	62%
White, not Hispanic Females	-	-	-	-	-	-	-	-	2	20%	12	4%
Black/African American, not Hispanic Males	4	40%	7	35%	1	25%	3	30%	-	-	53	18%
Black/African American, not Hispanic Females	2	20%	2	10%	-	-	2	20%	1	10%	26	9%
Hispanic/Latino Males	-	-	1	5%	1	25%	2	20%	-	-	13	4%
Hispanic/Latina Females	-	-	-	-	-	-	1	10%	-	-	1	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	10%	-	-	-	-	-	-	5	2%
<b>Total</b>	<b>10</b>		<b>20</b>		<b>4</b>		<b>10</b>		<b>10</b>		<b>290</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Butler**

Transmission Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	5	63%	12	71%	1	25%	5	71%	3	43%	169	68%
Injection drug use (IDU)	-	-	-	-	1	25%	-	-	1	14%	12	5%
Male-to-male sexual contact and IDU	1	13%	-	-	-	-	-	-	-	-	12	5%
Heterosexual contact	-	-	1	6%	-	-	-	-	1	14%	12	5%
Other/unknown	2	25%	4	24%	2	50%	2	29%	2	29%	43	17%
Subtotal	8	100%	17	100%	4	100%	7	100%	7	100%	248	100%
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	4	10%
Heterosexual contact	2	100%	3	100%	-	-	3	100%	2	67%	31	78%
Other/unknown	-	-	-	-	-	-	-	-	1	33%	5	13%
Subtotal	2	100%	3	100%	-	-	3	100%	3	100%	40	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	1	50%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	50%
Subtotal	-	-	-	-	-	-	-	-	-	-	2	100%
<b>Total</b>	<b>10</b>		<b>20</b>		<b>4</b>		<b>10</b>		<b>10</b>		<b>290</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Butler**

Exposure Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	5	50%	12	60%	1	25%	5	50%	3	30%	162	56%
Injection drug use (IDU) only	-	-	-	-	1	25%	-	-	1	10%	9	3%
Heterosexual contact only	2	20%	4	20%	-	-	3	30%	3	30%	43	15%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	8	3%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	2%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	2%
MSM & IDU & Heterosexual contact	1	10%	-	-	-	-	-	-	-	-	4	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	1	<1%
Other/unknown	2	20%	4	20%	2	50%	2	20%	3	30%	49	17%
<b>Total</b>	<b>10</b>		<b>20</b>		<b>4</b>		<b>10</b>		<b>10</b>		<b>290</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Butler

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Butler**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	147.8	271	79%	106	73%	165	84%
Females	36.7	70	21%	39	27%	31	16%
<b>Age at end of year</b>							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	2	1%	1	1%	1	1%
20-24	31.1	10	3%	6	4%	4	2%
25-29	90.4	19	6%	13	9%	6	3%
30-34	147.5	34	10%	15	10%	19	10%
35-39	180.3	40	12%	21	14%	19	10%
40-44	210.7	50	15%	24	17%	26	13%
45-49	208.3	51	15%	20	14%	31	16%
50-54	219.9	59	17%	18	12%	41	21%
55-64	105.9	50	15%	18	12%	32	16%
65+	52.7	26	8%	9	6%	17	9%
<b>Race/Ethnicity</b>							
White, not Hispanic	62.4	193	57%	82	57%	111	57%
Black/African American, not Hispanic	368.3	107	31%	45	31%	62	32%
Hispanic/Latino	160.6	27	8%	11	8%	16	8%
Asian/Pacific Islander	*	4	1%	2	1%	2	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	10	3%	5	3%	5	3%
<b>Race/Sex</b>							
White, not Hispanic Males	110.2	167	49%	68	47%	99	51%
White, not Hispanic Females	16.5	26	8%	14	10%	12	6%
Black/African American, not Hispanic Males	531.0	75	22%	26	18%	49	25%
Black/African American, not Hispanic Females	214.3	32	9%	19	13%	13	7%
Hispanic/Latino Males	222.0	19	6%	6	4%	13	7%
Hispanic/Latina Females	96.9	8	2%	5	3%	3	2%
Asian/Pacific Islander Males	*	3	1%	2	1%	1	1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	10	3%	5	3%	5	3%
<b>Total</b>	<b>91.1</b>	<b>341</b>		<b>145</b>		<b>196</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Butler

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	173	65%	58	56%	115	71%
Injection drug use (IDU)	7	3%	3	3%	4	2%
Male-to-male sexual contact and IDU	6	2%	2	2%	4	2%
Heterosexual contact	18	7%	9	9%	9	6%
Other/unknown	63	24%	32	31%	31	19%
Subtotal	267	100%	104	100%	163	100%
<b>Female adult or adolescent</b>						
Injection drug use	9	13%	2	5%	7	23%
Heterosexual contact	45	65%	24	63%	21	68%
Other/unknown	15	22%	12	32%	3	10%
Subtotal	69	100%	38	100%	31	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	3	60%	2	67%	1	50%
Other/unknown	2	40%	1	33%	1	50%
Subtotal	5	100%	3	100%	2	100%
<b>Total</b>	<b>341</b>		<b>145</b>		<b>196</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Butler

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	169	50%	58	40%	111	57%
Injection drug use (IDU) only	6	2%	1	1%	5	3%
Heterosexual contact only	63	18%	33	23%	30	15%
MSM & IDU	4	1%	2	1%	2	1%
IDU & Heterosexual contact	10	3%	4	3%	6	3%
MSM & Heterosexual contact	4	1%	-	-	4	2%
MSM & IDU & Heterosexual contact	2	1%	-	-	2	1%
Perinatal exposure	3	1%	2	1%	1	1%
Other/unknown	80	23%	45	31%	35	18%
<b>Total</b>	<b>341</b>		<b>145</b>		<b>196</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Butler**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	122	74%	35	48%	9	47%	2	67%	-	-	5	71%
Injection drug use (IDU)	-	-	4	5%	3	16%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	6	4%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	6	4%	8	11%	3	16%	1	33%	-	-	-	-
Other/unknown	31	19%	26	36%	4	21%	-	-	-	-	2	29%
Subtotal	165	100%	73	100%	19	100%	3	100%	-	-	7	100%
<b>Female adult or adolescent</b>												
Injection drug use	4	16%	3	9%	1	13%	1	100%	-	-	-	-
Heterosexual contact	13	52%	24	75%	5	63%	-	-	-	-	3	100%
Other/unknown	8	32%	5	16%	2	25%	-	-	-	-	-	-
Subtotal	25	100%	32	100%	8	100%	1	100%	-	-	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	3	100%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	2	100%	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	-	-	-	-	-	-	-	-
<b>Total</b>	<b>193</b>		<b>107</b>		<b>27</b>		<b>4</b>		<b>0</b>		<b>10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Butler**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	121	63%	34	32%	7	26%	2	50%	-	-	5	50%
Injection drug use (IDU) only	2	1%	2	2%	1	4%	1	25%	-	-	-	-
Heterosexual contact only	19	10%	32	30%	8	30%	1	25%	-	-	3	30%
MSM & IDU	4	2%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	1%	5	5%	3	11%	-	-	-	-	-	-
MSM & Heterosexual contact	1	1%	1	1%	2	7%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	2	1%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	3	2%	-	-	-	-	-	-	-	-	-	-
Other/unknown	39	20%	33	31%	6	22%	-	-	-	-	2	20%
<b>Total</b>	<b>193</b>		<b>107</b>		<b>27</b>		<b>4</b>		<b>0</b>		<b>10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Butler**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	221	80%	234	80%	245	80%	259	79%	271	79%
Females	56	20%	60	20%	61	20%	67	21%	70	21%
<b>Age at end of year</b>										
<13	-	-	-	-	-	-	-	-	-	-
13-14	1	<1%	-	-	-	-	-	-	-	-
15-19	3	1%	3	1%	3	1%	3	1%	2	1%
20-24	10	4%	11	4%	9	3%	10	3%	10	3%
25-29	17	6%	16	5%	19	6%	20	6%	19	6%
30-34	30	11%	32	11%	27	9%	29	9%	34	10%
35-39	41	15%	37	13%	42	14%	46	14%	40	12%
40-44	41	15%	44	15%	43	14%	46	14%	50	15%
45-49	53	19%	58	20%	61	20%	54	17%	51	15%
50-54	37	13%	41	14%	46	15%	53	16%	59	17%
55-64	31	11%	36	12%	39	13%	41	13%	50	15%
65+	13	5%	16	5%	17	6%	24	7%	26	8%
<b>Race/Ethnicity</b>										
White, not Hispanic	166	60%	172	59%	179	58%	186	57%	193	57%
Black/African American, not Hispanic	79	29%	89	30%	91	30%	100	31%	107	31%
Hispanic/Latino	20	7%	21	7%	24	8%	26	8%	27	8%
Asian/Pacific Islander	3	1%	3	1%	3	1%	4	1%	4	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	9	3%	9	3%	10	3%	10	3%
<b>Race/Sex</b>										
White, not Hispanic Males	143	52%	149	51%	155	51%	161	49%	167	49%
White, not Hispanic Females	23	8%	23	8%	24	8%	25	8%	26	8%
Black/African American, not Hispanic Males	55	20%	61	21%	63	21%	70	21%	75	22%
Black/African American, not Hispanic Females	24	9%	28	10%	28	9%	30	9%	32	9%
Hispanic/Latino Males	14	5%	15	5%	18	6%	18	6%	19	6%
Hispanic/Latina Females	6	2%	6	2%	6	2%	8	2%	8	2%
Asian/Pacific Islander Males	2	1%	2	1%	2	1%	3	1%	3	1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	9	3%	9	3%	10	3%	10	3%
<b>Total</b>	<b>277</b>		<b>294</b>		<b>306</b>		<b>326</b>		<b>341</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Butler

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	141	65%	149	65%	156	65%	166	65%	173	65%
Injection drug use (IDU)	6	3%	6	3%	6	2%	7	3%	7	3%
Male-to-male sexual contact and IDU	6	3%	6	3%	6	2%	6	2%	6	2%
Heterosexual contact	17	8%	17	7%	17	7%	17	7%	18	7%
Other/unknown	47	22%	52	23%	56	23%	59	23%	63	24%
Subtotal	217	100%	230	100%	241	100%	255	100%	267	100%
<b>Female adult or adolescent</b>										
Injection drug use	8	15%	8	14%	8	13%	9	14%	9	13%
Heterosexual contact	36	65%	39	66%	39	65%	43	65%	45	65%
Other/unknown	11	20%	12	20%	13	22%	14	21%	15	22%
Subtotal	55	100%	59	100%	60	100%	66	100%	69	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	3	60%	3	60%	3	60%	3	60%	3	60%
Other/unknown	2	40%	2	40%	2	40%	2	40%	2	40%
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
<b>Total</b>	<b>277</b>		<b>294</b>		<b>306</b>		<b>326</b>		<b>341</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Butler

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	137	49%	145	49%	152	50%	162	50%	169	50%
Injection drug use (IDU) only	6	2%	6	2%	6	2%	6	2%	6	2%
Heterosexual contact only	53	19%	56	19%	56	18%	60	18%	63	18%
MSM & IDU	4	1%	4	1%	4	1%	4	1%	4	1%
IDU & Heterosexual contact	8	3%	8	3%	8	3%	10	3%	10	3%
MSM & Heterosexual contact	4	1%	4	1%	4	1%	4	1%	4	1%
MSM & IDU & Heterosexual contact	2	1%	2	1%	2	1%	2	1%	2	1%
Perinatal exposure	3	1%	3	1%	3	1%	3	1%	3	1%
Other/unknown	60	22%	66	22%	71	23%	75	23%	80	23%
<b>Total</b>	<b>277</b>		<b>294</b>		<b>306</b>		<b>326</b>		<b>341</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.