

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Lucas

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Lucas

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	18.0	38	88%	33	87%	5	100%	-	-
Females	2.2	5	12%	5	13%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	17.6	5	12%	5	13%	-	-	-	-
20-24	34.6	11	26%	11	29%	-	-	-	-
25-29	15.8	5	12%	5	13%	-	-	-	-
30-34	18.4	5	12%	4	11%	1	20%	-	-
35-39	*	-	-	-	-	-	-	-	-
40-44	22.1	6	14%	5	13%	1	20%	-	-
45-49	*	4	9%	3	8%	1	20%	-	-
50-54	*	2	5%	1	3%	1	20%	-	-
55-64	*	4	9%	3	8%	1	20%	-	-
65+	*	1	2%	1	3%	-	-	-	-
Race/Ethnicity									
White, not Hispanic	4.6	14	33%	11	29%	3	60%	-	-
Black/African American, not Hispanic	26.6	22	51%	21	55%	1	20%	-	-
Hispanic/Latino	17.1	5	12%	4	11%	1	20%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	2	5%	2	5%	-	-	-	-
Race/Sex									
White, not Hispanic Males	8.1	12	28%	9	24%	3	60%	-	-
White, not Hispanic Females	*	2	5%	2	5%	-	-	-	-
Black/African American, not Hispanic Males	51.2	20	47%	19	50%	1	20%	-	-
Black/African American, not Hispanic Females	*	2	5%	2	5%	-	-	-	-
Hispanic/Latino Males	*	4	9%	3	8%	1	20%	-	-
Hispanic/Latina Females	*	1	2%	1	3%	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	2	5%	2	5%	-	-	-	-
Total	9.9	43		38		5			

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	31	82%	27	82%	4	80%	-	-
Injection drug use (IDU)	1	3%	1	3%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	3	8%	3	9%	-	-	-	-
Other/unknown	3	8%	2	6%	1	20%	-	-
Subtotal	38	100%	33	100%	5	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	4	80%	4	80%	-	-	-	-
Other/unknown	1	20%	1	20%	-	-	-	-
Subtotal	5	100%	5	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	43		38		5		0	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	30	70%	26	68%	4	80%	-	-
Injection drug use (IDU) only	1	2%	1	3%	-	-	-	-
Heterosexual contact only	7	16%	7	18%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	2%	1	3%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	4	9%	3	8%	1	20%	-	-
Total	43		38		5		-	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	10	83%	15	75%	4	100%	-	-	-	-	2	100%
Injection drug use (IDU)	-	-	1	5%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	3	15%	-	-	-	-	-	-	-	-
Other/unknown	2	17%	1	5%	-	-	-	-	-	-	-	-
Subtotal	12	100%	20	100%	4	100%	-	-	-	-	2	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	2	100%	2	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	14		22		5		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	9	64%	15	68%	4	80%	-	-	-	-	2	100%
Injection drug use (IDU) only	-	-	1	5%	-	-	-	-	-	-	-	-
Heterosexual contact only	2	14%	5	23%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	7%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	14%	1	5%	1	20%	-	-	-	-	-	-
Total	14		22		5		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Lucas

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	9.5	20	74%	15.2	32	76%	17.1	36	82%	12.8	27	90%
Females	3.1	7	26%	4.5	10	24%	3.6	8	18%	*	3	10%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	*	1	4%	*	4	10%	*	-	-	*	3	10%
20-24	18.9	6	22%	22.0	7	17%	44.0	14	32%	25.1	8	27%
25-29	*	1	4%	15.8	5	12%	19.0	6	14%	*	4	13%
30-34	18.4	5	19%	18.4	5	12%	22.0	6	14%	*	2	7%
35-39	36.5	9	33%	28.4	7	17%	20.3	5	11%	*	2	7%
40-44	*	-	-	18.4	5	12%	22.1	6	14%	*	1	3%
45-49	*	2	7%	*	4	10%	*	2	5%	18.8	5	17%
50-54	*	2	7%	*	2	5%	*	2	5%	*	1	3%
55-64	*	1	4%	*	3	7%	*	2	5%	*	3	10%
65+	*	-	-	*	-	-	*	1	2%	*	1	3%
Race/Ethnicity												
White, not Hispanic	4.3	13	48%	4.9	15	36%	8.2	25	57%	3.0	9	30%
Black/African American, not Hispanic	14.5	12	44%	31.4	26	62%	19.3	16	36%	21.7	18	60%
Hispanic/Latino	*	1	4%	*	1	2%	*	-	-	*	3	10%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	-	-	*	3	7%	*	-	-
Race/Sex												
White, not Hispanic Males	6.8	10	37%	9.5	14	33%	12.8	19	43%	4.7	7	23%
White, not Hispanic Females	*	3	11%	*	1	2%	3.8	6	14%	*	2	7%
Black/African American, not Hispanic Males	20.5	8	30%	43.5	17	40%	35.9	14	32%	43.5	17	57%
Black/African American, not Hispanic Females	*	4	15%	20.6	9	21%	*	2	5%	*	1	3%
Hispanic/Latino Males	*	1	4%	*	1	2%	*	-	-	*	3	10%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	-	-	*	3	7%	*	-	-
Total	6.2	27		9.6	42		10.1	44		6.9	30	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Lucas

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	17	85%	21	66%	26	72%	22	81%
Injection drug use (IDU)	-	-	1	3%	1	3%	1	4%
Male-to-male sexual contact and IDU	-	-	1	3%	3	8%	1	4%
Heterosexual contact	3	15%	4	13%	-	-	1	4%
Other/unknown	-	-	5	16%	6	17%	2	7%
Subtotal	20	100%	32	100%	36	100%	27	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	1	13%	-	-
Heterosexual contact	7	100%	10	100%	6	75%	3	100%
Other/unknown	-	-	-	-	1	13%	-	-
Subtotal	7	100%	10	100%	8	100%	3	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	27		42		44		30	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Lucas

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	59%	21	50%	25	57%	20	67%
Injection drug use (IDU) only	-	-	1	2%	-	-	1	3%
Heterosexual contact only	10	37%	14	33%	6	14%	4	13%
MSM & IDU	-	-	1	2%	3	7%	1	3%
IDU & Heterosexual contact	-	-	-	-	2	5%	-	-
MSM & Heterosexual contact	1	4%	-	-	1	2%	2	7%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	-	-	5	12%	7	16%	2	7%
Total	27		42		44		30	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Lucas

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	11	73%	19	76%	29	85%	20	87%	33	87%
Females	4	27%	6	24%	5	15%	3	13%	5	13%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	7%	3	12%	-	-	3	13%	5	13%
20-24	5	33%	6	24%	13	38%	5	22%	11	29%
25-29	1	7%	5	20%	5	15%	4	17%	5	13%
30-34	-	-	3	12%	5	15%	1	4%	4	11%
35-39	7	47%	2	8%	4	12%	-	-	-	-
40-44	-	-	2	8%	4	12%	1	4%	5	13%
45-49	-	-	2	8%	1	3%	5	22%	3	8%
50-54	-	-	1	4%	1	3%	1	4%	1	3%
55-64	1	7%	1	4%	-	-	3	13%	3	8%
65+	-	-	-	-	1	3%	-	-	1	3%
Race/Ethnicity										
White, not Hispanic	5	33%	9	36%	19	56%	7	30%	11	29%
Black/African American, not Hispanic	9	60%	15	60%	14	41%	15	65%	21	55%
Hispanic/Latino	-	-	1	4%	-	-	1	4%	4	11%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	7%	-	-	1	3%	-	-	2	5%
Race/Sex										
White, not Hispanic Males	3	20%	8	32%	16	47%	5	22%	9	24%
White, not Hispanic Females	2	13%	1	4%	3	9%	2	9%	2	5%
Black/African American, not Hispanic Males	7	47%	10	40%	12	35%	14	61%	19	50%
Black/African American, not Hispanic Females	2	13%	5	20%	2	6%	1	4%	2	5%
Hispanic/Latino Males	-	-	1	4%	-	-	1	4%	3	8%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	1	3%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	7%	-	-	1	3%	-	-	2	5%
Total	15		25		34		23		38	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	8	73%	14	74%	20	69%	16	80%	27	82%
Injection drug use (IDU)	-	-	-	-	1	3%	1	5%	1	3%
Male-to-male sexual contact and IDU	-	-	1	5%	3	10%	-	-	-	-
Heterosexual contact	3	27%	1	5%	-	-	1	5%	3	9%
Other/unknown	-	-	3	16%	5	17%	2	10%	2	6%
Subtotal	11	100%	19	100%	29	100%	20	100%	33	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	1	20%	-	-	-	-
Heterosexual contact	4	100%	6	100%	4	80%	3	100%	4	80%
Other/unknown	-	-	-	-	-	-	-	-	1	20%
Subtotal	4	100%	6	100%	5	100%	3	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	15		25		34		23		38	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	47%	14	56%	19	56%	14	61%	26	68%
Injection drug use (IDU) only	-	-	-	-	-	-	1	4%	1	3%
Heterosexual contact only	7	47%	7	28%	4	12%	4	17%	7	18%
MSM & IDU	-	-	1	4%	3	9%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	2	6%	-	-	-	-
MSM & Heterosexual contact	1	7%	-	-	1	3%	2	9%	1	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	3	12%	5	15%	2	9%	3	8%
Total	15		25		34		23		38	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Lucas

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	18	75%	24	80%	8	73%	11	92%	6	100%	702	78%
Females	6	25%	6	20%	3	27%	1	8%	-	-	193	22%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	15	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	3	13%	1	3%	-	-	-	-	-	-	27	3%
20-24	1	4%	3	10%	1	9%	5	42%	-	-	87	10%
25-29	5	21%	1	3%	1	9%	1	8%	-	-	152	17%
30-34	6	25%	5	17%	1	9%	2	17%	1	17%	181	20%
35-39	2	8%	7	23%	2	18%	2	17%	-	-	174	19%
40-44	2	8%	6	20%	2	18%	1	8%	2	33%	107	12%
45-49	3	13%	3	10%	1	9%	-	-	1	17%	61	7%
50-54	2	8%	1	3%	1	9%	-	-	1	17%	50	6%
55-64	-	-	3	10%	2	18%	-	-	1	17%	35	4%
65+	-	-	-	-	-	-	1	8%	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	12	50%	12	40%	6	55%	5	42%	4	67%	478	53%
Black/African American, not Hispanic	9	38%	16	53%	2	18%	3	25%	1	17%	360	40%
Hispanic/Latino	3	13%	2	7%	1	9%	4	33%	1	17%	44	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	2	18%	-	-	-	-	10	1%
Race/Sex												
White, not Hispanic Males	9	38%	11	37%	3	27%	5	42%	4	67%	413	46%
White, not Hispanic Females	3	13%	1	3%	3	27%	-	-	-	-	65	7%
Black/African American, not Hispanic Males	6	25%	11	37%	2	18%	3	25%	1	17%	246	27%
Black/African American, not Hispanic Females	3	13%	5	17%	-	-	-	-	-	-	114	13%
Hispanic/Latino Males	3	13%	2	7%	1	9%	3	25%	1	17%	33	4%
Hispanic/Latina Females	-	-	-	-	-	-	1	8%	-	-	11	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	2	18%	-	-	-	-	10	1%
Total	24		30		11		12		6		895	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	14	78%	15	63%	7	88%	9	82%	4	67%	506	73%
Injection drug use (IDU)	-	-	1	4%	-	-	-	-	-	-	43	6%
Male-to-male sexual contact and IDU	1	6%	1	4%	-	-	1	9%	1	17%	54	8%
Heterosexual contact	3	17%	3	13%	-	-	-	-	-	-	46	7%
Other/unknown	-	-	4	17%	1	13%	1	9%	1	17%	46	7%
Subtotal	18	100%	24	100%	8	100%	11	100%	6	100%	695	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	30	16%
Heterosexual contact	6	100%	6	100%	2	67%	1	100%	-	-	144	78%
Other/unknown	-	-	-	-	1	33%	-	-	-	-	11	6%
Subtotal	6	100%	6	100%	3	100%	1	100%	-	-	185	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	14	93%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	7%
Subtotal	-	-	-	-	-	-	-	-	-	-	15	100%
Total	24		30		11		12		6		895	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	14	58%	15	50%	7	64%	9	75%	4	67%	480	54%
Injection drug use (IDU) only	-	-	1	3%	-	-	-	-	-	-	35	4%
Heterosexual contact only	9	38%	9	30%	2	18%	1	8%	-	-	190	21%
MSM & IDU	1	4%	1	3%	-	-	1	8%	1	17%	40	4%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	38	4%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	26	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	14	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	14	2%
Other/unknown	-	-	4	13%	2	18%	1	8%	1	17%	58	6%
Total	24		30		11		12		6		895	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Lucas

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Lucas

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	333.6	704	73%	363	72%	341	74%
Females	117.7	264	27%	142	28%	122	26%
Age at end of year							
<13	*	3	<1%	2	<1%	1	<1%
13-14	*	2	<1%	-	-	2	<1%
15-19	35.1	10	1%	8	2%	2	<1%
20-24	122.6	39	4%	32	6%	7	2%
25-29	189.9	60	6%	51	10%	9	2%
30-34	235.1	64	7%	44	9%	20	4%
35-39	381.4	94	10%	55	11%	39	8%
40-44	585.1	159	16%	74	15%	85	18%
45-49	578.5	154	16%	75	15%	79	17%
50-54	555.3	172	18%	69	14%	103	22%
55-64	293.5	173	18%	77	15%	96	21%
65+	59.9	38	4%	18	4%	20	4%
Race/Ethnicity							
White, not Hispanic	142.4	433	45%	223	44%	210	45%
Black/African American, not Hispanic	547.0	453	47%	237	47%	216	47%
Hispanic/Latino	181.4	53	5%	26	5%	27	6%
Asian/Pacific Islander	66.0	5	1%	4	1%	1	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	23	2%	15	3%	8	2%
Race/Sex							
White, not Hispanic Males	231.0	342	35%	168	33%	174	38%
White, not Hispanic Females	58.3	91	9%	55	11%	36	8%
Black/African American, not Hispanic Males	763.2	298	31%	158	31%	140	30%
Black/African American, not Hispanic Females	354.1	155	16%	79	16%	76	16%
Hispanic/Latino Males	277.0	40	4%	21	4%	19	4%
Hispanic/Latina Females	88.0	13	1%	5	1%	8	2%
Asian/Pacific Islander Males	*	4	<1%	3	1%	1	<1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	23	2%	15	3%	8	2%
Total	222.4	968		505		463	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	493	70%	253	70%	240	71%
Injection drug use (IDU)	38	5%	15	4%	23	7%
Male-to-male sexual contact and IDU	43	6%	22	6%	21	6%
Heterosexual contact	67	10%	32	9%	35	10%
Other/unknown	60	9%	40	11%	20	6%
Subtotal	701	100%	362	100%	339	100%
Female adult or adolescent						
Injection drug use	24	9%	8	6%	16	14%
Heterosexual contact	208	82%	114	82%	94	82%
Other/unknown	22	9%	17	12%	5	4%
Subtotal	254	100%	139	100%	115	100%
Child (<13 yrs at diagnosis)						
Perinatal	13	100%	4	100%	9	100%
Other/unknown	-	-	-	-	-	-
Subtotal	13	100%	4	100%	9	100%
Total	968		505		463	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	453	47%	233	46%	220	48%
Injection drug use (IDU) only	26	3%	11	2%	15	3%
Heterosexual contact only	275	28%	146	29%	129	28%
MSM & IDU	32	3%	19	4%	13	3%
IDU & Heterosexual contact	36	4%	12	2%	24	5%
MSM & Heterosexual contact	40	4%	20	4%	20	4%
MSM & IDU & Heterosexual contact	11	1%	3	1%	8	2%
Perinatal exposure	13	1%	4	1%	9	2%
Other/unknown	82	8%	57	11%	25	5%
Total	968		505		463	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Lucas

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	266	78%	180	61%	33	83%	2	50%	-	-	12	60%
Injection drug use (IDU)	16	5%	18	6%	3	8%	-	-	-	-	1	5%
Male-to-male sexual contact and IDU	22	6%	16	5%	3	8%	-	-	-	-	2	10%
Heterosexual contact	15	4%	49	16%	1	3%	-	-	-	-	2	10%
Other/unknown	21	6%	34	11%	-	-	2	50%	-	-	3	15%
Subtotal	340	100%	297	100%	40	100%	4	100%	-	-	20	100%
Female adult or adolescent												
Injection drug use	6	7%	18	12%	-	-	-	-	-	-	-	-
Heterosexual contact	79	89%	118	79%	9	75%	1	100%	1	100%	-	-
Other/unknown	4	4%	13	9%	3	25%	-	-	-	-	2	100%
Subtotal	89	100%	149	100%	12	100%	1	100%	1	100%	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	4	100%	7	100%	1	100%	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	7	100%	1	100%	-	-	-	-	1	100%
Total	433		453		53		5		1		23	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Lucas

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	247	57%	161	36%	31	58%	2	40%	-	-	12	52%
Injection drug use (IDU) only	10	2%	16	4%	-	-	-	-	-	-	-	-
Heterosexual contact only	94	22%	167	37%	10	19%	1	20%	1	100%	2	9%
MSM & IDU	14	3%	15	3%	2	4%	-	-	-	-	1	4%
IDU & Heterosexual contact	12	3%	20	4%	3	6%	-	-	-	-	1	4%
MSM & Heterosexual contact	19	4%	19	4%	2	4%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	8	2%	1	0%	1	2%	-	-	-	-	1	4%
Perinatal exposure	4	1%	7	2%	1	2%	-	-	-	-	1	4%
Other/unknown	25	6%	47	10%	3	6%	2	40%	-	-	5	22%
Total	433		453		53		5		1		23	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Lucas

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	577	71%	607	71%	642	71%	666	72%	704	73%
Females	238	29%	248	29%	256	29%	260	28%	264	27%
Age at end of year										
<13	6	1%	6	1%	5	1%	4	<1%	3	<1%
13-14	2	<1%	2	<1%	1	<1%	2	<1%	2	<1%
15-19	6	1%	7	1%	8	1%	9	1%	10	1%
20-24	28	3%	31	4%	40	4%	39	4%	39	4%
25-29	44	5%	47	5%	52	6%	52	6%	60	6%
30-34	72	9%	69	8%	62	7%	67	7%	64	7%
35-39	124	15%	111	13%	111	12%	100	11%	94	10%
40-44	145	18%	154	18%	159	18%	159	17%	159	16%
45-49	158	19%	161	19%	159	18%	156	17%	154	16%
50-54	111	14%	134	16%	148	16%	157	17%	172	18%
55-64	101	12%	111	13%	125	14%	150	16%	173	18%
65+	18	2%	22	3%	28	3%	31	3%	38	4%
Race/Ethnicity										
White, not Hispanic	374	46%	388	45%	409	46%	419	45%	433	45%
Black/African American, not Hispanic	373	46%	398	47%	416	46%	432	47%	453	47%
Hispanic/Latino	44	5%	45	5%	46	5%	48	5%	53	5%
Asian/Pacific Islander	5	1%	5	1%	5	1%	5	1%	5	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	18	2%	18	2%	21	2%	21	2%	23	2%
Race/Sex										
White, not Hispanic Males	295	36%	307	36%	323	36%	330	36%	342	35%
White, not Hispanic Females	79	10%	81	9%	86	10%	89	10%	91	9%
Black/African American, not Hispanic Males	231	28%	248	29%	263	29%	278	30%	298	31%
Black/African American, not Hispanic Females	142	17%	150	18%	153	17%	154	17%	155	16%
Hispanic/Latino Males	32	4%	33	4%	34	4%	36	4%	40	4%
Hispanic/Latina Females	12	1%	12	1%	12	1%	12	1%	13	1%
Asian/Pacific Islander Males	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	18	2%	18	2%	21	2%	21	2%	23	2%
Total	815		855		898		926		968	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Lucas

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	397	69%	418	69%	443	69%	463	70%	493	70%
Injection drug use (IDU)	35	6%	35	6%	36	6%	37	6%	38	5%
Male-to-male sexual contact and IDU	40	7%	40	7%	43	7%	43	6%	43	6%
Heterosexual contact	59	10%	63	10%	63	10%	64	10%	67	10%
Other/unknown	43	7%	48	8%	54	8%	56	8%	60	9%
Subtotal	574	100%	604	100%	639	100%	663	100%	701	100%
Female adult or adolescent										
Injection drug use	23	10%	23	10%	24	10%	24	10%	24	9%
Heterosexual contact	184	81%	194	82%	201	82%	205	82%	208	82%
Other/unknown	21	9%	21	9%	21	9%	21	8%	22	9%
Subtotal	228	100%	238	100%	246	100%	250	100%	254	100%
Child (<13 yrs at diagnosis)										
Perinatal	13	100%	13	100%	13	100%	13	100%	13	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	13	100%	13	100%	13	100%	13	100%	13	100%
Total	815		855		898		926		968	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Lucas

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	362	44%	383	45%	406	45%	424	46%	453	47%
Injection drug use (IDU) only	24	3%	24	3%	24	3%	25	3%	26	3%
Heterosexual contact only	243	30%	257	30%	264	29%	269	29%	275	28%
MSM & IDU	29	4%	29	3%	32	4%	32	3%	32	3%
IDU & Heterosexual contact	34	4%	34	4%	36	4%	36	4%	36	4%
MSM & Heterosexual contact	35	4%	35	4%	37	4%	39	4%	40	4%
MSM & IDU & Heterosexual contact	11	1%	11	1%	11	1%	11	1%	11	1%
Perinatal exposure	13	2%	13	2%	13	1%	13	1%	13	1%
Other/unknown	64	8%	69	8%	75	8%	77	8%	82	8%
Total	815		855		898		926		968	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.