

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Sandusky

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Sandusky**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	128.3	38	86%	14	93%	24	83%
Females	19.6	6	14%	1	7%	5	17%
<b>Age at end of year</b>							
<13	*	1	2%	-	-	1	3%
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	2%	1	7%	-	-
25-29	153.1	5	11%	3	20%	2	7%
30-34	*	-	-	-	-	-	-
35-39	202.4	7	16%	2	13%	5	17%
40-44	159.8	6	14%	4	27%	2	7%
45-49	153.7	6	14%	-	-	6	21%
50-54	157.5	7	16%	2	13%	5	17%
55-64	91.0	8	18%	2	13%	6	21%
65+	*	3	7%	1	7%	2	7%
<b>Race/Ethnicity</b>							
White, not Hispanic	56.6	29	66%	11	73%	18	62%
Black/African American, not Hispanic	344.0	6	14%	3	20%	3	10%
Hispanic/Latino	122.0	7	16%	-	-	7	24%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	5%	1	7%	1	3%
<b>Race/Sex</b>							
White, not Hispanic Males	107.5	27	61%	10	67%	17	59%
White, not Hispanic Females	*	2	5%	1	7%	1	3%
Black/African American, not Hispanic Males	535.3	5	11%	3	20%	2	7%
Black/African American, not Hispanic Females	*	1	2%	-	-	1	3%
Hispanic/Latino Males	*	4	9%	-	-	4	14%
Hispanic/Latina Females	*	3	7%	-	-	3	10%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	5%	1	7%	1	3%
<b>Total</b>	<b>73.1</b>	<b>44</b>		<b>15</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Sandusky**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	28	76%	10	71%	18	78%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	11%	3	21%	1	4%
Heterosexual contact	3	8%	1	7%	2	9%
Other/unknown	2	5%	-	-	2	9%
Subtotal	37	100%	14	100%	23	100%
<b>Female adult or adolescent</b>						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	6	100%	1	100%	5	100%
Other/unknown	-	-	-	-	-	-
Subtotal	6	100%	1	100%	5	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	1	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
<b>Total</b>	<b>44</b>		<b>15</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Sandusky**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	26	59%	10	67%	16	55%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	9	20%	2	13%	7	24%
MSM & IDU	4	9%	3	20%	1	3%
IDU & Heterosexual contact	-	-	-	-	-	-
MSM & Heterosexual contact	2	5%	-	-	2	7%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	2%	-	-	1	3%
Other/unknown	2	5%	-	-	2	7%
<b>Total</b>	<b>44</b>		<b>15</b>		<b>29</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Sandusky**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	21	78%	<10	*	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	15%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	7%	<10	*	-	-	-	-	-	-	-	-
Other/unknown	-	-	<10	*	<10	*	-	-	-	-	-	-
Subtotal	27	100%	<10	*	<10	*	-	-	-	-	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	<10	*	<10	*	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	<10	*	<10	*	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	<10	*
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	<10	*
<b>Total</b>	<b>29</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Sandusky**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	20	69%	<10	*	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	4	14%	<10	*	<10	*	-	-	-	-	-	-
MSM & IDU	4	14%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	3%	-	-	<10	*	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	<10	*
Other/unknown	-	-	<10	*	<10	*	-	-	-	-	-	-
<b>Total</b>	<b>29</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Sandusky**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	28	85%	32	86%	35	85%	36	86%	38	86%
Females	5	15%	5	14%	6	15%	6	14%	6	14%
<b>Age at end of year</b>										
<13	1	3%	1	3%	1	2%	1	2%	1	2%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	-	-	2	5%	3	7%	2	5%	1	2%
25-29	-	-	-	-	-	-	2	5%	5	11%
30-34	4	12%	1	3%	1	2%	-	-	-	-
35-39	6	18%	10	27%	7	17%	8	19%	7	16%
40-44	4	12%	2	5%	7	17%	7	17%	6	14%
45-49	7	21%	8	22%	8	20%	6	14%	6	14%
50-54	7	21%	8	22%	6	15%	7	17%	7	16%
55-64	4	12%	3	8%	6	15%	7	17%	8	18%
65+	-	-	2	5%	2	5%	2	5%	3	7%
<b>Race/Ethnicity</b>										
White, not Hispanic	23	70%	25	68%	27	66%	28	67%	29	66%
Black/African American, not Hispanic	4	12%	4	11%	5	12%	5	12%	6	14%
Hispanic/Latino	5	15%	6	16%	7	17%	7	17%	7	16%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	2	5%	2	5%	2	5%	2	5%
<b>Race/Sex</b>										
White, not Hispanic Males	21	64%	23	62%	25	61%	26	62%	27	61%
White, not Hispanic Females	2	6%	2	5%	2	5%	2	5%	2	5%
Black/African American, not Hispanic Males	3	9%	3	8%	4	10%	4	10%	5	11%
Black/African American, not Hispanic Females	1	3%	1	3%	1	2%	1	2%	1	2%
Hispanic/Latino Males	3	9%	4	11%	4	10%	4	10%	4	9%
Hispanic/Latina Females	2	6%	2	5%	3	7%	3	7%	3	7%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	2	5%	2	5%	2	5%	2	5%
<b>Total</b>	<b>33</b>		<b>37</b>		<b>41</b>		<b>42</b>		<b>44</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Sandusky

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	20	74%	23	74%	25	74%	26	74%	28	76%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	15%	4	13%	4	12%	4	11%	4	11%
Heterosexual contact	3	11%	3	10%	3	9%	3	9%	3	8%
Other/unknown	-	-	1	3%	2	6%	2	6%	2	5%
<b>Subtotal</b>	<b>27</b>	<b>100%</b>	<b>31</b>	<b>100%</b>	<b>34</b>	<b>100%</b>	<b>35</b>	<b>100%</b>	<b>37</b>	<b>100%</b>
<b>Female adult or adolescent</b>										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	5	100%	5	100%	6	100%	6	100%	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
<b>Subtotal</b>	<b>5</b>	<b>100%</b>	<b>5</b>	<b>100%</b>	<b>6</b>	<b>100%</b>	<b>6</b>	<b>100%</b>	<b>6</b>	<b>100%</b>
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	1	100%	1	100%	1	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
<b>Subtotal</b>	<b>1</b>	<b>100%</b>								
<b>Total</b>	<b>33</b>		<b>37</b>		<b>41</b>		<b>42</b>		<b>44</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Sandusky

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	18	55%	21	57%	23	56%	24	57%	26	59%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	8	24%	8	22%	9	22%	9	21%	9	20%
MSM & IDU	4	12%	4	11%	4	10%	4	10%	4	9%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	2	6%	2	5%	2	5%	2	5%	2	5%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	3%	1	3%	1	2%	1	2%	1	2%
Other/unknown	-	-	1	3%	2	5%	2	5%	2	5%
<b>Total</b>	<b>33</b>		<b>37</b>		<b>41</b>		<b>42</b>		<b>44</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.