

## Guidance for Users of Ohio HIV/AIDS Surveillance Data

**\*\*Please note the HIV/AIDS surveillance data presented herein is the most currently available from the Ohio Department of Health (ODH) HIV/AIDS Surveillance Program. Please consider the following in your use of Ohio's HIV/AIDS surveillance data.\*\***

### *Explanation of Terms*

Ohio's HIV/AIDS surveillance data tables provide useful information on new diagnoses of HIV infection, persons living with an HIV infection, and deaths among person with an HIV infection reported to the ODH from physicians, hospitals, outpatient facilities and labs. A *diagnosis of HIV infection* refers to persons newly diagnosed and reported with an HIV infection in a particular year, regardless of the stage of disease at initial diagnosis. This includes persons newly diagnosed with HIV (not AIDS), persons previously diagnosed with HIV who are now newly diagnosed with AIDS, and persons concurrently diagnosed with HIV and AIDS at initial diagnosis. *Persons living with an HIV infection* represent all persons ever diagnosed and reported with an HIV infection since the beginning of the epidemic living in Ohio who have not been reported as having died. *Deaths* among persons with an HIV infection represent deaths attributed to any cause among persons reported with a diagnosis of HIV infection.

### *Reporting Delays/Lags*

A key consideration for the analysis of HIV/AIDS surveillance data requires the factoring in of reporting delays. A reporting delay – the time between diagnosis of HIV and/or AIDS and report of the case to the public health authority – can differ among exposure, geographic, racial/ethnic, age, sex and vital status categories. In some instances, reporting delays for mode of exposure (i.e. risk factor) can be several years, even with active follow-up to obtain this key information.

All population-based disease surveillance systems experience reporting lags and incomplete reporting. The reporting lags for Ohio's HIV/AIDS surveillance data range from 9-18 months. This is an acceptable range and is recognized as meeting the CDC national quality standards for HIV/AIDS reporting. This allows states time to collect more complete and accurate information so that the data can be used to accurately identify trends in HIV transmission and populations at risk.

### *Representativeness and Limitations of Data*

HIV infection surveillance data represents confidential reports of HIV infection and AIDS diagnoses; not all persons with an HIV infection. The distinction here is that HIV diagnosis data represent the earliest date of diagnosis reported to the ODH HIV/AIDS Surveillance Program. The earliest date reported may not be the earliest date an individual became aware of their HIV infection. Individuals may have previously tested anonymously or were diagnosed out-of-state prior to being confidentially tested and reported to Ohio. HIV infection surveillance data may underestimate the level of recently infected persons because some infected persons do not know they are infected as they have not sought testing or have sought testing but did not respond to learn their test results. Reporting of behavioral risk information may not be complete as some persons diagnosed with an HIV infection may be reluctant to disclose their sexual and drug use history.

Due to the long and variable period from initial HIV infection to the development of AIDS; trends in AIDS surveillance do not represent recent HIV infections. Asymptomatic HIV-infected persons are not represented by AIDS case data. In addition, incomplete HIV or CD4+ t-cell testing may interfere with the representativeness of reporting. Widespread use of HAART complicates the interpretation of AIDS case surveillance data and estimation of the HIV/AIDS epidemic in an area. Newly reported AIDS cases may reflect treatment failures or the failure of the health care system to halt progression of HIV infection to AIDS. AIDS cases represent late-stage HIV infection.

# Diagnoses of HIV and/or AIDS Reported in Ohio

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
<b>Sex</b>									
Males	14.3	813	86%	608	86%	151	84%	54	87%
Females	2.3	137	14%	101	14%	28	16%	8	13%
<b>Age at diagnosis (yr)</b>									
<13	*	3	<1%	3	<1%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	6.9	53	6%	50	7%	1	1%	2	3%
20-24	27.4	217	23%	187	26%	22	12%	8	13%
25-29	24.2	181	19%	150	21%	22	12%	9	15%
30-34	16.8	121	13%	80	11%	29	16%	12	19%
35-39	14.1	96	10%	74	10%	16	9%	6	10%
40-44	11.8	86	9%	60	8%	22	12%	4	6%
45-49	11.5	87	9%	50	7%	27	15%	10	16%
50-54	4.7	40	4%	23	3%	16	9%	1	2%
55-64	3.5	55	6%	27	4%	20	11%	8	13%
65+	0.6	11	1%	5	1%	4	2%	2	3%
<b>Race/Ethnicity</b>									
White, not Hispanic	3.8	355	37%	257	36%	77	43%	21	34%
Black/African American, not Hispanic	34.6	493	52%	371	52%	85	47%	37	60%
Hispanic/Latino	16.9	68	7%	52	7%	13	7%	3	5%
Asian/Pacific Islander	3.4	8	1%	6	1%	1	1%	1	2%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-	-	-
Unknown	*	25	3%	22	3%	3	2%	-	-
<b>Race/Sex</b>									
White, not Hispanic Males	6.9	314	33%	227	32%	68	38%	19	31%
White, not Hispanic Females	0.9	41	4%	30	4%	9	5%	2	3%
Black/African American, not Hispanic Males	60.5	412	43%	315	44%	66	37%	31	50%
Black/African American, not Hispanic Females	10.9	81	9%	56	8%	19	11%	6	10%
Hispanic/Latino Males	28.5	59	6%	43	6%	13	7%	3	5%
Hispanic/Latina Females	4.6	9	1%	9	1%	-	-	-	-
Asian/Pacific Islander Males	5.3	6	1%	4	1%	1	1%	1	2%
Asian/Pacific Islander Females	*	2	<1%	2	<1%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-	-	-
Unknown	*	25	3%	22	3%	3	2%	-	-
<b>Total</b>	<b>8.2</b>	<b>950</b>		<b>709</b>		<b>179</b>		<b>62</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	553	68%	424	70%	99	66%	30	56%
Injection drug use (IDU)	15	2%	11	2%	4	3%	-	-
Male-to-male sexual contact and IDU	19	2%	19	3%	-	-	-	-
Heterosexual contact	38	5%	25	4%	5	3%	8	15%
Other/unknown	187	23%	128	21%	43	28%	16	30%
Subtotal	812	100%	607	100%	151	100%	54	100%
<b>Female adult or adolescent</b>								
Injection drug use	13	10%	11	11%	2	7%	-	-
Heterosexual contact	87	65%	67	68%	16	57%	4	50%
Other/unknown	34	25%	20	20%	10	36%	4	50%
Subtotal	134	100%	98	100%	28	100%	8	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	2	50%	2	50%	-	-	-	-
Other/unknown	2	50%	2	50%	-	-	-	-
Subtotal	4	100%	4	100%	-	-	-	-
<b>Total</b>	<b>950</b>		<b>709</b>		<b>179</b>		<b>62</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	537	57%	410	58%	97	54%	30	48%
Injection drug use (IDU) only	12	1%	9	1%	3	2%	-	-
Heterosexual contact only	125	13%	92	13%	21	12%	12	19%
MSM & IDU	15	2%	15	2%	-	-	-	-
IDU & Heterosexual contact	16	2%	13	2%	3	2%	-	-
MSM & Heterosexual contact	16	2%	14	2%	2	1%	-	-
MSM & IDU & Heterosexual contact	4	<1%	4	1%	-	-	-	-
Perinatal exposure	3	<1%	3	<1%	-	-	-	-
Other/unknown	222	23%	149	21%	43	24%	20	32%
<b>Total</b>	<b>950</b>		<b>709</b>		<b>179</b>		<b>62</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, Ohio

Transmission Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	214	68%	286	70%	36	61%	4	67%	-	-	13	59%
Injection drug use (IDU)	11	4%	4	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	14	4%	1	0%	2	3%	-	-	-	-	2	9%
Heterosexual contact	10	3%	25	6%	1	2%	-	-	-	-	2	9%
Other/unknown	65	21%	95	23%	20	34%	2	33%	-	-	5	23%
Subtotal	314	100%	411	100%	59	100%	6	100%	-	-	22	100%
<b>Female adult or adolescent</b>												
Injection drug use	11	27%	-	-	2	25%	-	-	-	-	-	-
Heterosexual contact	24	59%	54	68%	4	50%	1	100%	1	100%	3	100%
Other/unknown	6	15%	26	33%	2	25%	-	-	-	-	-	-
Subtotal	41	100%	80	100%	8	100%	1	100%	1	100%	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	1	50%	-	-	1	100%	-	-	-	-
Other/unknown	-	-	1	50%	1	100%	-	-	-	-	-	-
Subtotal	-	-	2	100%	1	100%	1	100%	-	-	-	-
<b>Total</b>	<b>355</b>		<b>493</b>		<b>68</b>		<b>8</b>		<b>1</b>		<b>25</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, Ohio

Exposure Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	209	59%	277	56%	35	51%	3	38%	-	-	13	52%
Injection drug use (IDU) only	9	3%	2	0%	1	1%	-	-	-	-	-	-
Heterosexual contact only	34	10%	79	16%	5	7%	1	13%	1	100%	5	20%
MSM & IDU	11	3%	1	0%	2	3%	-	-	-	-	1	4%
IDU & Heterosexual contact	13	4%	2	0%	1	1%	-	-	-	-	-	-
MSM & Heterosexual contact	5	1%	9	2%	1	1%	1	13%	-	-	-	-
MSM & IDU & Heterosexual contact	3	1%	-	-	-	-	-	-	-	-	1	4%
Perinatal exposure	-	-	1	0%	1	1%	1	13%	-	-	-	-
Other/unknown	71	20%	122	25%	22	32%	2	25%	-	-	5	20%
<b>Total</b>	<b>355</b>		<b>493</b>		<b>68</b>		<b>8</b>		<b>1</b>		<b>25</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county**

County <sup>a</sup>	2014 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Adams	*	3	3	-	-
Allen	5.7	6	5	1	-
Ashland	*	1	1	-	-
Ashtabula	*	3	3	-	-
Athens	*	2	2	-	-
Auglaize	*	-	-	-	-
Belmont	*	2	1	-	1
Brown	*	2	1	-	1
Butler	5.1	19	13	6	-
Carroll	*	-	-	-	-
Champaign	*	3	1	1	1
Clark	*	3	2	1	-
Clermont	5.0	10	6	3	1
Clinton	*	1	1	-	-
Columbiana	*	4	2	1	1
Coshocton	*	2	-	2	-
Crawford	*	-	-	-	-
Cuyahoga	16.7	210	171	26	13
Darke	*	3	3	-	-
Defiance	*	2	2	-	-
Delaware	3.2	6	5	1	-
Erie	*	4	4	-	-
Fairfield	4.7	7	5	2	-
Fayette	*	1	1	-	-
Franklin	17.5	215	146	49	20
Fulton	*	-	-	-	-
Gallia	*	2	1	1	-
Geauga	*	2	2	-	-
Greene		7	7	-	-
Guernsey	*	2	2	-	-
<b>Ohio</b>	<b>8.2</b>	<b>950</b>	<b>709</b>	<b>179</b>	<b>62</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2014 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Hamilton	18.3	148	111	29	8
Hancock	*	-	-	-	-
Hardin	*	1	-	1	-
Harrison	*	-	-	-	-
Henry	*	-	-	-	-
Highland	*	-	-	-	-
Hocking	*	1	1	-	-
Holmes	*	-	-	-	-
Huron	*	-	-	-	-
Jackson	*	2	2	-	-
Jefferson	*	1	-	1	-
Knox	*	-	-	-	-
Lake	2.6	6	4	2	-
Lawrence	*	-	-	-	-
Licking	*	4	2	2	-
Logan	*	-	-	-	-
Lorain	2.6	8	7	-	1
Lucas	9.9	43	38	5	-
Madison	*	-	-	-	-
Mahoning	9.4	22	10	7	5
Marion	*	1	-	-	1
Medina	*	2	2	-	-
Meigs	*	-	-	-	-
Mercer	*	1	1	-	-
Miami	*	2	1	1	-
Monroe	*	-	-	-	-
Montgomery	11.1	59	45	11	3
Morgan	*	1	-	1	-
Morrow	*	1	-	-	1
Muskingum	*	2	1	1	-
<b>Ohio</b>	<b>8.2</b>	<b>950</b>	<b>709</b>	<b>179</b>	<b>62</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2014 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Noble	*	-	-	-	-
Ottawa	*	-	-	-	-
Paulding	*	-	-	-	-
Perry	*	2	1	1	-
Pickaway	*	1	1	-	-
Pike	*	-	-	-	-
Portage	5.6	9	9	-	-
Preble	*	3	3	-	-
Putnam	*	-	-	-	-
Richland	*	-	-	-	-
Ross	*	1	1	-	-
Sandusky	*	2	1	1	-
Scioto	*	1	-	-	1
Seneca	*	-	-	-	-
Shelby	*	4	2	-	2
Stark	5.3	20	15	4	1
Summit	7.0	38	34	4	-
Trumbull	3.4	7	3	3	1
Tuscarawas	*	1	1	-	-
Union	*	1	1	-	-
Van Wert	*	-	-	-	-
Vinton	*	-	-	-	-
Warren	3.2	7	6	1	-
Washington	*	-	-	-	-
Wayne	*	2	2	-	-
Williams	*	3	2	1	-
Wood	*	2	1	1	-
Wyandot	*	-	-	-	-
No County	*	19	11	8	-
<b>Ohio</b>	<b>8.2</b>	<b>950</b>	<b>709</b>	<b>179</b>	<b>62</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio

Characteristic	2013 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
<b>Sex</b>										
Males	15.3	871	82%	607	81%	203	87%	61	79%	
Females	3.2	189	18%	142	19%	31	13%	16	21%	
<b>Age at diagnosis (yr)</b>										
<13	0.5	9	1%	9	1%	-	-	-	-	
13-14	*	1	<1%	1	<1%	-	-	-	-	
15-19	8.3	64	6%	60	8%	2	1%	2	3%	
20-24	29.6	234	22%	185	25%	46	20%	3	4%	
25-29	23.1	173	16%	131	17%	36	15%	6	8%	
30-34	16.2	117	11%	90	12%	18	8%	9	12%	
35-39	15.0	102	10%	68	9%	22	9%	12	16%	
40-44	13.5	98	9%	59	8%	29	12%	10	13%	
45-49	12.8	97	9%	60	8%	23	10%	14	18%	
50-54	8.1	69	7%	35	5%	25	11%	9	12%	
55-64	4.8	76	7%	42	6%	25	11%	9	12%	
65+	1.1	20	2%	9	1%	8	3%	3	4%	
<b>Race/Ethnicity</b>										
White, not Hispanic	4.7	432	41%	297	40%	100	43%	35	45%	
Black/African American, not Hispanic	35.8	511	48%	369	49%	110	47%	32	42%	
Hispanic/Latino	15.4	62	6%	41	5%	12	5%	9	12%	
Asian/Pacific Islander	3.4	8	1%	5	1%	3	1%	-	-	
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-	-	-	
Unknown	*	46	4%	36	5%	9	4%	1	1%	
<b>Race/Sex</b>										
White, not Hispanic Males	8.2	375	35%	252	34%	93	40%	30	39%	
White, not Hispanic Females	1.2	57	5%	45	6%	7	3%	5	6%	
Black/African American, not Hispanic Males	59.9	408	38%	294	39%	91	39%	23	30%	
Black/African American, not Hispanic Females	13.8	103	10%	75	10%	19	8%	9	12%	
Hispanic/Latino Males	22.7	47	4%	29	4%	10	4%	8	10%	
Hispanic/Latina Females	7.7	15	1%	12	2%	2	1%	1	1%	
Asian/Pacific Islander Males	5.3	6	1%	4	1%	2	1%	-	-	
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%	-	-	
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	46	4%	36	5%	9	4%	1	1%	
<b>Total</b>	<b>9.1</b>	<b>1,060</b>		<b>749</b>		<b>234</b>		<b>77</b>		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	2013 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	589	68%	421	70%	132	65%	36	60%
Injection drug use (IDU)	20	2%	15	2%	4	2%	1	2%
Male-to-male sexual contact and IDU	19	2%	17	3%	2	1%	-	-
Heterosexual contact	37	4%	20	3%	14	7%	3	5%
Other/unknown	202	23%	131	22%	51	25%	20	33%
Subtotal	867	100%	604	100%	203	100%	60	100%
<b>Female adult or adolescent</b>								
Injection drug use	19	10%	17	13%	2	6%	-	-
Heterosexual contact	113	62%	81	60%	21	68%	11	69%
Other/unknown	50	27%	37	27%	8	26%	5	31%
Subtotal	182	100%	135	100%	31	100%	16	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	4	36%	4	40%	-	-	-	-
Other/unknown	7	64%	6	60%	-	-	1	100%
Subtotal	11	100%	10	100%	-	-	1	100%
<b>Total</b>	<b>1,060</b>		<b>749</b>		<b>234</b>		<b>77</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	2013 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	563	53%	399	53%	129	55%	35	45%
Injection drug use (IDU) only	17	2%	14	2%	2	1%	1	1%
Heterosexual contact only	150	14%	101	13%	35	15%	14	18%
MSM & IDU	14	1%	13	2%	1	<1%	-	-
IDU & Heterosexual contact	22	2%	18	2%	4	2%	-	-
MSM & Heterosexual contact	26	2%	22	3%	3	1%	1	1%
MSM & IDU & Heterosexual contact	5	<1%	4	1%	1	<1%	-	-
Perinatal exposure	6	1%	5	1%	-	-	1	1%
Other/unknown	257	24%	173	23%	59	25%	25	32%
<b>Total</b>	<b>1,060</b>		<b>749</b>		<b>234</b>		<b>77</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, Ohio

Transmission Category <sup>a</sup>	2013 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	281	75%	253	62%	31	66%	4	80%	1	100%	19	56%
Injection drug use (IDU)	13	3%	7	2%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	12	3%	1	0%	2	4%	-	-	-	-	4	12%
Heterosexual contact	11	3%	20	5%	2	4%	-	-	-	-	4	12%
Other/unknown	58	15%	124	31%	12	26%	1	20%	-	-	7	21%
Subtotal	375	100%	405	100%	47	100%	5	100%	1	100%	34	100%
<b>Female adult or adolescent</b>												
Injection drug use	16	29%	1	1%	-	-	-	-	-	-	2	18%
Heterosexual contact	29	52%	69	70%	7	47%	1	100%	-	-	7	64%
Other/unknown	11	20%	29	29%	8	53%	-	-	-	-	2	18%
Subtotal	56	100%	99	100%	15	100%	1	100%	-	-	11	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	1	100%	3	43%	-	-	-	-	-	-	-	-
Other/unknown	-	-	4	57%	-	-	2	100%	-	-	1	100%
Subtotal	1	100%	7	100%	-	-	2	100%	-	-	1	100%
<b>Total</b>	<b>432</b>		<b>511</b>		<b>62</b>		<b>8</b>		<b>1</b>		<b>46</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, Ohio

Exposure Category <sup>a</sup>	2013 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	271	63%	241	47%	29	47%	4	50%	1	100%	17	37%
Injection drug use (IDU) only	13	3%	4	1%	-	-	-	-	-	-	-	-
Heterosexual contact only	40	9%	89	17%	9	15%	1	13%	-	-	11	24%
MSM & IDU	8	2%	1	0%	2	3%	-	-	-	-	3	7%
IDU & Heterosexual contact	16	4%	4	1%	-	-	-	-	-	-	2	4%
MSM & Heterosexual contact	10	2%	12	2%	2	3%	-	-	-	-	2	4%
MSM & IDU & Heterosexual contact	4	1%	-	-	-	-	-	-	-	-	1	2%
Perinatal exposure	1	0%	5	1%	-	-	-	-	-	-	-	-
Other/unknown	69	16%	155	30%	20	32%	3	38%	-	-	10	22%
<b>Total</b>	<b>432</b>		<b>511</b>		<b>62</b>		<b>8</b>		<b>1</b>		<b>46</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county**

County <sup>a</sup>	2013 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Adams	*	1	-	1	-
Allen	12.4	13	9	4	-
Ashland	*	4	2	2	-
Ashtabula	5.0	5	4	1	-
Athens	*	-	-	-	-
Auglaize	*	-	-	-	-
Belmont	*	2	2	-	-
Brown	*	1	1	-	-
Butler	6.7	25	16	7	2
Carroll	*	1	-	1	-
Champaign	12.8	5	2	1	2
Clark	3.7	5	1	4	-
Clermont	4.5	9	9	-	-
Clinton	*	1	1	-	-
Columbiana	5.7	6	-	4	2
Coshocton	*	1	1	-	-
Crawford	*	2	-	-	2
Cuyahoga	18.1	228	166	44	18
Darke	*	2	2	-	-
Defiance	*	-	-	-	-
Delaware	3.7	7	4	1	2
Erie	6.6	5	3	1	1
Fairfield	*	4	2	2	-
Fayette	*	-	-	-	-
Franklin	18.4	227	155	54	18
Fulton	*	-	-	-	-
Gallia	*	1	-	1	-
Geauga	*	1	1	-	-
Greene		11	6	4	1
Guernsey	*	2	2	-	-
<b>Ohio</b>	<b>9.1</b>	<b>1,060</b>	<b>749</b>	<b>234</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2013 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Hamilton	20.3	164	110	44	10
Hancock	*	4	3	1	-
Hardin	*	3	1	2	-
Harrison	*	-	-	-	-
Henry	*	1	1	-	-
Highland	*	-	-	-	-
Hocking	*	1	1	-	-
Holmes	*	-	-	-	-
Huron	*	2	2	-	-
Jackson	*	-	-	-	-
Jefferson	*	1	1	-	-
Knox	*	1	1	-	-
Lake	3.5	8	6	2	-
Lawrence	*	2	1	1	-
Licking	3.5	6	6	-	-
Logan	*	-	-	-	-
Lorain	4.9	15	11	3	1
Lucas	6.9	30	23	5	2
Madison	*	1	1	-	-
Mahoning	7.7	18	14	2	2
Marion	*	3	2	-	1
Medina	4.0	7	4	3	-
Meigs	*	-	-	-	-
Mercer	*	1	-	1	-
Miami	7.7	8	7	1	-
Monroe	*	-	-	-	-
Montgomery	12.2	65	50	12	3
Morgan	*	-	-	-	-
Morrow	*	3	3	-	-
Muskingum	*	3	2	-	1
<b>Ohio</b>	<b>9.1</b>	<b>1,060</b>	<b>749</b>	<b>234</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2013 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Noble	*	-	-	-	-
Ottawa	*	1	1	-	-
Paulding	*	2	2	-	-
Perry	*	1	1	-	-
Pickaway	*	2	1	-	1
Pike	*	-	-	-	-
Portage	3.7	6	6	-	-
Preble	*	1	1	-	-
Putnam	*	-	-	-	-
Richland	*	4	4	-	-
Ross	*	1	-	1	-
Sandusky	*	1	-	1	-
Scioto	*	2	1	1	-
Seneca	*	1	1	-	-
Shelby	*	1	-	1	-
Stark	3.2	12	9	1	2
Summit	7.0	38	32	5	1
Trumbull	5.4	11	7	2	2
Tuscarawas	*	1	-	1	-
Union	*	-	-	-	-
Van Wert	*	-	-	-	-
Vinton	*	-	-	-	-
Warren	4.1	9	4	3	2
Washington	*	3	3	-	-
Wayne	*	1	1	-	-
Williams	*	1	-	1	-
Wood	6.2	8	6	2	-
Wyandot	*	-	-	-	-
No County	*	37	30	6	1
<b>Ohio</b>	<b>9.1</b>	<b>1,060</b>	<b>749</b>	<b>234</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
<b>Sex</b>									
Males	14.8	839	82%	550	81%	228	85%	61	79%
Females	3.2	187	18%	130	19%	41	15%	16	21%
<b>Age at diagnosis (yr)</b>									
<13	0.4	8	1%	6	1%	-	-	2	3%
13-14	*	-	-	-	-	-	-	-	-
15-19	8.2	63	6%	57	8%	5	2%	1	1%
20-24	27.1	214	21%	170	25%	37	14%	7	9%
25-29	21.6	162	16%	118	17%	36	13%	8	10%
30-34	20.2	146	14%	93	14%	39	14%	14	18%
35-39	13.7	93	9%	55	8%	28	10%	10	13%
40-44	12.5	91	9%	49	7%	34	13%	8	10%
45-49	13.0	98	10%	57	8%	29	11%	12	16%
50-54	9.1	77	8%	38	6%	32	12%	7	9%
55-64	4.0	63	6%	31	5%	24	9%	8	10%
65+	0.6	11	1%	6	1%	5	2%	-	-
<b>Race/Ethnicity</b>									
White, not Hispanic	4.5	418	41%	277	41%	112	42%	29	38%
Black/African American, not Hispanic	34.1	486	47%	331	49%	118	44%	37	48%
Hispanic/Latino	14.1	57	6%	33	5%	19	7%	5	6%
Asian/Pacific Islander	2.6	6	1%	5	1%	-	-	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	59	6%	34	5%	20	7%	5	6%
<b>Race/Sex</b>									
White, not Hispanic Males	8.1	368	36%	238	35%	103	38%	27	35%
White, not Hispanic Females	1.1	50	5%	39	6%	9	3%	2	3%
Black/African American, not Hispanic Males	53.7	366	36%	250	37%	91	34%	25	32%
Black/African American, not Hispanic Females	16.1	120	12%	81	12%	27	10%	12	16%
Hispanic/Latino Males	25.1	52	5%	29	4%	18	7%	5	6%
Hispanic/Latina Females	2.6	5	<1%	4	1%	1	<1%	-	-
Asian/Pacific Islander Males	4.5	5	<1%	4	1%	-	-	1	1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	59	6%	34	5%	20	7%	5	6%
<b>Total</b>	<b>8.8</b>	<b>1,026</b>		<b>680</b>		<b>269</b>		<b>77</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	572	68%	390	71%	151	66%	31	52%
Injection drug use (IDU)	18	2%	12	2%	6	3%	-	-
Male-to-male sexual contact and IDU	25	3%	15	3%	8	4%	2	3%
Heterosexual contact	26	3%	19	3%	6	3%	1	2%
Other/unknown	195	23%	112	20%	57	25%	26	43%
Subtotal	836	100%	548	100%	228	100%	60	100%
<b>Female adult or adolescent</b>								
Injection drug use	8	4%	6	5%	2	5%	-	-
Heterosexual contact	107	59%	75	60%	26	63%	6	40%
Other/unknown	65	36%	43	35%	13	32%	9	60%
Subtotal	180	100%	124	100%	41	100%	15	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	5	50%	3	38%	-	-	2	100%
Other/unknown	5	50%	5	63%	-	-	-	-
Subtotal	10	100%	8	100%	-	-	2	100%
<b>Total</b>	<b>1,026</b>		<b>680</b>		<b>269</b>		<b>77</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	539	53%	366	54%	143	53%	30	39%
Injection drug use (IDU) only	18	2%	12	2%	6	2%	-	-
Heterosexual contact only	133	13%	94	14%	32	12%	7	9%
MSM & IDU	17	2%	12	2%	5	2%	-	-
IDU & Heterosexual contact	8	1%	6	1%	2	1%	-	-
MSM & Heterosexual contact	33	3%	24	4%	8	3%	1	1%
MSM & IDU & Heterosexual contact	8	1%	3	<1%	3	1%	2	3%
Perinatal exposure	7	1%	5	1%	-	-	2	3%
Other/unknown	263	26%	158	23%	70	26%	35	45%
<b>Total</b>	<b>1,026</b>		<b>680</b>		<b>269</b>		<b>77</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, Ohio

Transmission Category <sup>a</sup>	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	266	72%	234	64%	38	73%	1	20%	-	-	33	69%
Injection drug use (IDU)	7	2%	6	2%	3	6%	-	-	-	-	2	4%
Male-to-male sexual contact and IDU	15	4%	7	2%	1	2%	-	-	-	-	2	4%
Heterosexual contact	7	2%	17	5%	1	2%	-	-	-	-	1	2%
Other/unknown	73	20%	99	27%	9	17%	4	80%	-	-	10	21%
Subtotal	368	100%	363	100%	52	100%	5	100%	-	-	48	100%
<b>Female adult or adolescent</b>												
Injection drug use	5	10%	2	2%	1	20%	-	-	-	-	-	-
Heterosexual contact	28	56%	69	61%	3	60%	1	100%	-	-	6	55%
Other/unknown	17	34%	42	37%	1	20%	-	-	-	-	5	45%
Subtotal	50	100%	113	100%	5	100%	1	100%	-	-	11	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	5	50%	-	-	-	-	-	-	-	-
Other/unknown	-	-	5	50%	-	-	-	-	-	-	-	-
Subtotal	-	-	10	100%	-	-	-	-	-	-	-	-
<b>Total</b>	<b>418</b>		<b>486</b>		<b>57</b>		<b>6</b>		<b>0</b>		<b>59</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, Ohio

Exposure Category <sup>a</sup>	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	251	60%	219	45%	37	65%	1	17%	-	-	31	53%
Injection drug use (IDU) only	8	2%	5	1%	4	7%	-	-	-	-	1	2%
Heterosexual contact only	35	8%	86	18%	4	7%	1	17%	-	-	7	12%
MSM & IDU	11	3%	5	1%	-	-	-	-	-	-	1	2%
IDU & Heterosexual contact	4	1%	3	1%	-	-	-	-	-	-	1	2%
MSM & Heterosexual contact	15	4%	15	3%	1	2%	-	-	-	-	2	3%
MSM & IDU & Heterosexual contact	4	1%	2	0%	1	2%	-	-	-	-	1	2%
Perinatal exposure	-	-	7	1%	-	-	-	-	-	-	-	-
Other/unknown	90	22%	144	30%	10	18%	4	67%	-	-	15	25%
<b>Total</b>	<b>418</b>		<b>486</b>		<b>57</b>		<b>6</b>		<b>0</b>		<b>59</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county**

County <sup>a</sup>	2012 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Adams	*	-	-	-	-
Allen	6.7	7	4	2	1
Ashland	*	1	-	1	-
Ashtabula	*	1	-	1	-
Athens	*	1	-	-	1
Auglaize	*	-	-	-	-
Belmont	*	3	1	1	1
Brown	*	1	-	-	1
Butler	2.7	10	6	4	-
Carroll	*	2	2	-	-
Champaign	*	2	1	1	-
Clark	*	4	3	-	1
Clermont	4.0	8	4	4	-
Clinton	*	4	3	1	-
Columbiana	*	3	1	2	-
Coshocton	*	1	-	-	1
Crawford	*	1	-	1	-
Cuyahoga	16.5	208	158	38	12
Darke	*	1	-	1	-
Defiance	*	1	-	1	-
Delaware	6.9	13	9	3	1
Erie	*	2	2	-	-
Fairfield	*	4	3	-	1
Fayette	*	1	1	-	-
Franklin	20.0	246	158	65	23
Fulton	*	3	3	-	-
Gallia	*	1	1	-	-
Geauga	*	3	3	-	-
Greene		9	4	4	1
Guernsey	*	2	1	1	-
<b>Ohio</b>	<b>8.8</b>	<b>1,026</b>	<b>680</b>	<b>269</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2012 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Hamilton	18.2	147	88	53	6
Hancock	*	-	-	-	-
Hardin	*	1	1	-	-
Harrison	*	-	-	-	-
Henry	*	2	2	-	-
Highland	*	1	1	-	-
Hocking	*	1	-	-	1
Holmes	*	-	-	-	-
Huron	*	-	-	-	-
Jackson	*	2	2	-	-
Jefferson	7.4	5	1	3	1
Knox	*	2	2	-	-
Lake	*	4	4	-	-
Lawrence	*	1	1	-	-
Licking	5.9	10	4	3	3
Logan	*	-	-	-	-
Lorain	5.6	17	7	7	3
Lucas	10.1	44	34	6	4
Madison	*	1	1	-	-
Mahoning	9.9	23	14	8	1
Marion	*	4	1	2	1
Medina	4.0	7	4	2	1
Meigs	*	-	-	-	-
Mercer	*	1	1	-	-
Miami	*	2	-	1	1
Monroe	*	-	-	-	-
Montgomery	11.8	63	39	21	3
Morgan	*	1	-	1	-
Morrow	*	2	2	-	-
Muskingum	*	2	1	1	-
<b>Ohio</b>	<b>8.8</b>	<b>1,026</b>	<b>680</b>	<b>269</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2012 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Noble	*	-	-	-	-
Ottawa	*	2	2	-	-
Paulding	*	-	-	-	-
Perry	*	-	-	-	-
Pickaway	*	3	3	-	-
Pike	*	1	-	-	1
Portage	*	1	1	-	-
Preble	*	1	-	1	-
Putnam	*	-	-	-	-
Richland	*	4	1	2	1
Ross	*	2	1	1	-
Sandusky	*	3	1	2	-
Scioto	6.5	5	1	3	1
Seneca	*	-	-	-	-
Shelby	*	-	-	-	-
Stark	3.5	13	12	-	1
Summit	7.4	40	32	7	1
Trumbull	*	2	2	-	-
Tuscarawas	*	3	1	2	-
Union	*	1	-	1	-
Van Wert	*	-	-	-	-
Vinton	*	-	-	-	-
Warren	*	3	3	-	-
Washington	*	3	3	-	-
Wayne	*	4	3	1	-
Williams	*	-	-	-	-
Wood	4.6	6	4	1	1
Wyandot	*	-	-	-	-
No County	*	43	32	9	2
<b>Ohio</b>	<b>8.8</b>	<b>1,026</b>	<b>680</b>	<b>269</b>	<b>77</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio

Characteristic	2011 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
<b>Sex</b>									
Males	14.5	823	79%	546	80%	187	75%	90	80%
Females	3.7	220	21%	136	20%	62	25%	22	20%
<b>Age at diagnosis (yr)</b>									
<13	*	3	<1%	2	<1%	-	-	1	1%
13-14	*	-	-	-	-	-	-	-	-
15-19	8.3	64	6%	57	8%	6	2%	1	1%
20-24	27.8	220	21%	166	24%	46	18%	8	7%
25-29	20.7	155	15%	112	16%	33	13%	10	9%
30-34	18.8	136	13%	90	13%	29	12%	17	15%
35-39	16.8	114	11%	62	9%	32	13%	20	18%
40-44	15.4	112	11%	70	10%	27	11%	15	13%
45-49	12.2	92	9%	41	6%	31	12%	20	18%
50-54	9.4	80	8%	47	7%	23	9%	10	9%
55-64	3.4	54	5%	26	4%	20	8%	8	7%
65+	0.7	13	1%	9	1%	2	1%	2	2%
<b>Race/Ethnicity</b>									
White, not Hispanic	4.7	436	42%	291	43%	92	37%	53	47%
Black/African American, not Hispanic	35.5	506	49%	326	48%	129	52%	51	46%
Hispanic/Latino	14.9	60	6%	38	6%	16	6%	6	5%
Asian/Pacific Islander	3.0	7	1%	6	1%	-	-	1	1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-	-	-
Unknown	*	33	3%	20	3%	12	5%	1	1%
<b>Race/Sex</b>									
White, not Hispanic Males	8.0	366	35%	244	36%	76	31%	46	41%
White, not Hispanic Females	1.5	70	7%	47	7%	16	6%	7	6%
Black/African American, not Hispanic Males	54.6	372	36%	247	36%	88	35%	37	33%
Black/African American, not Hispanic Females	18.0	134	13%	79	12%	41	16%	14	13%
Hispanic/Latino Males	24.1	50	5%	32	5%	13	5%	5	4%
Hispanic/Latina Females	5.1	10	1%	6	1%	3	1%	1	1%
Asian/Pacific Islander Males	*	4	<1%	3	<1%	-	-	1	1%
Asian/Pacific Islander Females	*	3	<1%	3	<1%	-	-	-	-
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	33	3%	20	3%	12	5%	1	1%
<b>Total</b>	<b>9.0</b>	<b>1,043</b>		<b>682</b>		<b>249</b>		<b>112</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	2011 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	555	68%	369	68%	131	70%	55	61%
Injection drug use (IDU)	14	2%	10	2%	1	1%	3	3%
Male-to-male sexual contact and IDU	18	2%	14	3%	3	2%	1	1%
Heterosexual contact	33	4%	21	4%	10	5%	2	2%
Other/unknown	201	24%	130	24%	42	22%	29	32%
Subtotal	821	100%	544	100%	187	100%	90	100%
<b>Female adult or adolescent</b>								
Injection drug use	16	7%	8	6%	6	10%	2	10%
Heterosexual contact	141	64%	89	65%	40	65%	12	57%
Other/unknown	62	28%	39	29%	16	26%	7	33%
Subtotal	219	100%	136	100%	62	100%	21	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	3	100%	2	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	-	-	1	100%
<b>Total</b>	<b>1,043</b>		<b>682</b>		<b>249</b>		<b>112</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	2011 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	539	52%	356	52%	128	51%	55	49%
Injection drug use (IDU) only	13	1%	7	1%	3	1%	3	3%
Heterosexual contact only	174	17%	110	16%	50	20%	14	13%
MSM & IDU	13	1%	9	1%	3	1%	1	1%
IDU & Heterosexual contact	17	2%	11	2%	4	2%	2	2%
MSM & Heterosexual contact	16	2%	13	2%	3	1%	-	-
MSM & IDU & Heterosexual contact	5	<1%	5	1%	-	-	-	-
Perinatal exposure	3	<1%	2	<1%	-	-	1	1%
Other/unknown	263	25%	169	25%	58	23%	36	32%
<b>Total</b>	<b>1,043</b>		<b>682</b>		<b>249</b>		<b>112</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, Ohio

Transmission Category <sup>a</sup>	2011 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	259	71%	239	65%	30	60%	2	50%	-	-	25	83%
Injection drug use (IDU)	4	1%	3	1%	5	10%	-	-	-	-	2	7%
Male-to-male sexual contact and IDU	15	4%	3	1%	-	-	-	-	-	-	-	-
Heterosexual contact	10	3%	20	5%	2	4%	-	-	-	-	1	3%
Other/unknown	78	21%	105	28%	13	26%	2	50%	1	100%	2	7%
Subtotal	366	100%	370	100%	50	100%	4	100%	1	100%	30	100%
<b>Female adult or adolescent</b>												
Injection drug use	11	16%	5	4%	-	-	-	-	-	-	-	-
Heterosexual contact	41	59%	90	67%	6	67%	2	67%	-	-	2	67%
Other/unknown	18	26%	39	29%	3	33%	1	33%	-	-	1	33%
Subtotal	70	100%	134	100%	9	100%	3	100%	-	-	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	2	100%	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	1	100%	-	-	-	-	-	-
<b>Total</b>	<b>436</b>		<b>506</b>		<b>60</b>		<b>7</b>		<b>1</b>		<b>33</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, Ohio

Exposure Category <sup>a</sup>	2011 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	253	58%	231	46%	29	48%	2	29%	-	-	24	73%
Injection drug use (IDU) only	8	2%	2	0%	2	3%	-	-	-	-	1	3%
Heterosexual contact only	51	12%	110	22%	8	13%	2	29%	-	-	3	9%
MSM & IDU	11	3%	2	0%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	7	2%	6	1%	3	5%	-	-	-	-	1	3%
MSM & Heterosexual contact	6	1%	8	2%	1	2%	-	-	-	-	1	3%
MSM & IDU & Heterosexual contact	4	1%	1	0%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	2	0%	1	2%	-	-	-	-	-	-
Other/unknown	96	22%	144	28%	16	27%	3	43%	1	100%	3	9%
<b>Total</b>	<b>436</b>		<b>506</b>		<b>60</b>		<b>7</b>		<b>1</b>		<b>33</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county**

County <sup>a</sup>	2011 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Adams	*	-	-	-	-
Allen	6.7	7	5	1	1
Ashland	*	1	1	-	-
Ashtabula	*	1	-	-	1
Athens	*	3	2	1	-
Auglaize	*	-	-	-	-
Belmont	*	3	1	-	2
Brown	*	3	3	-	-
Butler	5.6	21	8	8	5
Carroll	*	-	-	-	-
Champaign	*	2	1	1	-
Clark	5.9	8	5	2	1
Clermont	5.0	10	5	5	-
Clinton	*	2	2	-	-
Columbiana	*	3	1	2	-
Coshocton	*	1	1	-	-
Crawford	*	2	1	1	-
Cuyahoga	15.7	198	138	34	26
Darke	*	3	2	1	-
Defiance	*	3	1	2	-
Delaware	3.2	6	4	-	2
Erie	*	3	3	-	-
Fairfield	4.7	7	4	1	2
Fayette	*	-	-	-	-
Franklin	21.2	261	163	69	29
Fulton	*	-	-	-	-
Gallia	*	1	1	-	-
Geauga	*	3	3	-	-
Greene		6	4	2	-
Guernsey	*	-	-	-	-
<b>Ohio</b>	<b>9.0</b>	<b>1,043</b>	<b>682</b>	<b>249</b>	<b>112</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2011 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Hamilton	16.9	136	90	38	8
Hancock	*	1	-	1	-
Hardin	*	-	-	-	-
Harrison	*	1	-	-	1
Henry	*	1	-	-	1
Highland	*	2	-	2	-
Hocking	*	-	-	-	-
Holmes	*	-	-	-	-
Huron	*	3	2	-	1
Jackson	*	-	-	-	-
Jefferson	*	2	2	-	-
Knox	*	3	3	-	-
Lake	2.2	5	3	-	2
Lawrence	*	1	-	1	-
Licking	3.0	5	1	4	-
Logan	*	1	-	-	1
Lorain	5.3	16	10	4	2
Lucas	9.6	42	25	8	9
Madison	*	-	-	-	-
Mahoning	11.1	26	16	6	4
Marion	*	3	3	-	-
Medina	*	3	3	-	-
Meigs	*	1	1	-	-
Mercer	*	3	2	1	-
Miami	*	2	1	1	-
Monroe	*	-	-	-	-
Montgomery	14.4	77	50	24	3
Morgan	*	1	1	-	-
Morrow	*	2	1	1	-
Muskingum	*	2	1	1	-
<b>Ohio</b>	<b>9.0</b>	<b>1,043</b>	<b>682</b>	<b>249</b>	<b>112</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2011 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Noble	*	-	-	-	-
Ottawa	*	2	2	-	-
Paulding	*	1	1	-	-
Perry	*	2	1	1	-
Pickaway	*	3	2	1	-
Pike	*	1	-	1	-
Portage	3.1	5	4	-	1
Preble	*	-	-	-	-
Putnam	*	1	1	-	-
Richland	*	1	1	-	-
Ross	7.8	6	4	1	1
Sandusky	8.3	5	4	-	1
Scioto	*	1	1	-	-
Seneca	*	-	-	-	-
Shelby	*	-	-	-	-
Stark	7.7	29	20	6	3
Summit	7.9	43	36	6	1
Trumbull	2.4	5	4	1	-
Tuscarawas	*	-	-	-	-
Union	*	1	1	-	-
Van Wert	*	-	-	-	-
Vinton	*	-	-	-	-
Warren	4.1	9	4	3	2
Washington	*	-	-	-	-
Wayne	*	1	1	-	-
Williams	*	1	-	1	-
Wood	*	1	1	-	-
Wyandot	*	-	-	-	-
No County	*	28	20	6	2
<b>Ohio</b>	<b>9.0</b>	<b>1,043</b>	<b>682</b>	<b>249</b>	<b>112</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio

Characteristic	2010 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
<b>Sex</b>									
Males	14.3	809	83%	495	82%	244	85%	70	78%
Females	2.9	170	17%	106	18%	44	15%	20	22%
<b>Age at diagnosis (yr)</b>									
<13	0.3	5	1%	4	1%	-	-	1	1%
13-14	*	1	<1%	1	<1%	-	-	-	-
15-19	10.1	78	8%	62	10%	15	5%	1	1%
20-24	24.5	194	20%	146	24%	44	15%	4	4%
25-29	19.0	142	15%	93	15%	38	13%	11	12%
30-34	16.8	121	12%	64	11%	44	15%	13	14%
35-39	16.8	114	12%	72	12%	29	10%	13	14%
40-44	14.4	105	11%	50	8%	39	14%	16	18%
45-49	13.2	100	10%	48	8%	36	13%	16	18%
50-54	7.4	63	6%	31	5%	24	8%	8	9%
55-64	3.0	47	5%	25	4%	16	6%	6	7%
65+	0.5	9	1%	5	1%	3	1%	1	1%
<b>Race/Ethnicity</b>									
White, not Hispanic	4.5	415	42%	246	41%	128	44%	41	46%
Black/African American, not Hispanic	33.1	472	48%	295	49%	135	47%	42	47%
Hispanic/Latino	11.2	45	5%	29	5%	11	4%	5	6%
Asian/Pacific Islander	2.6	6	1%	5	1%	1	<1%	-	-
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%	-	-
Unknown	*	40	4%	26	4%	12	4%	2	2%
<b>Race/Sex</b>									
White, not Hispanic Males	8.0	363	37%	214	36%	113	39%	36	40%
White, not Hispanic Females	1.1	52	5%	32	5%	15	5%	5	6%
Black/African American, not Hispanic Males	54.2	369	38%	230	38%	110	38%	29	32%
Black/African American, not Hispanic Females	13.8	103	11%	65	11%	25	9%	13	14%
Hispanic/Latino Males	18.3	38	4%	26	4%	9	3%	3	3%
Hispanic/Latina Females	3.6	7	1%	3	<1%	2	1%	2	2%
Asian/Pacific Islander Males	*	4	<1%	3	<1%	1	<1%	-	-
Asian/Pacific Islander Females	*	2	<1%	2	<1%	-	-	-	-
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	40	4%	26	4%	12	4%	2	2%
<b>Total</b>	<b>8.4</b>	<b>979</b>		<b>601</b>		<b>288</b>		<b>90</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	2010 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	570	71%	353	72%	175	72%	42	61%
Injection drug use (IDU)	8	1%	3	1%	4	2%	1	1%
Male-to-male sexual contact and IDU	22	3%	16	3%	5	2%	1	1%
Heterosexual contact	41	5%	28	6%	10	4%	3	4%
Other/unknown	163	20%	91	19%	50	20%	22	32%
Subtotal	804	100%	491	100%	244	100%	69	100%
<b>Female adult or adolescent</b>								
Injection drug use	13	8%	8	8%	2	5%	3	15%
Heterosexual contact	120	71%	74	70%	36	82%	10	50%
Other/unknown	36	21%	23	22%	6	14%	7	35%
Subtotal	169	100%	105	100%	44	100%	20	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	5	83%	4	80%	-	-	1	100%
Other/unknown	1	17%	1	20%	-	-	-	-
Subtotal	6	100%	5	100%	-	-	1	100%
<b>Total</b>	<b>979</b>		<b>601</b>		<b>288</b>		<b>90</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	2010 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	555	57%	345	57%	168	58%	42	47%
Injection drug use (IDU) only	8	1%	4	1%	2	1%	2	2%
Heterosexual contact only	161	16%	102	17%	46	16%	13	14%
MSM & IDU	18	2%	13	2%	4	1%	1	1%
IDU & Heterosexual contact	13	1%	7	1%	4	1%	2	2%
MSM & Heterosexual contact	15	2%	8	1%	7	2%	-	-
MSM & IDU & Heterosexual contact	4	<1%	3	<1%	1	<1%	-	-
Perinatal exposure	5	1%	4	1%	-	-	1	1%
Other/unknown	200	20%	115	19%	56	19%	29	32%
<b>Total</b>	<b>979</b>		<b>601</b>		<b>288</b>		<b>90</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, Ohio

Transmission Category <sup>a</sup>	2010 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	265	73%	254	70%	23	61%	2	50%	-	-	26	76%
Injection drug use (IDU)	4	1%	3	1%	1	3%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	15	4%	6	2%	1	0	-	-	-	-	-	-
Heterosexual contact	9	2%	23	6%	5	13%	1	25%	-	-	3	9%
Other/unknown	70	19%	78	21%	8	21%	1	25%	1	1	5	15%
Subtotal	363	100%	364	100%	38	100%	4	100%	1	1	34	100%
<b>Female adult or adolescent</b>												
Injection drug use	9	17%	3	3%	1	0	-	-	-	-	-	-
Heterosexual contact	33	63%	76	75%	3	43%	2	100%	-	-	6	100%
Other/unknown	10	19%	23	23%	3	43%	-	-	-	-	-	-
Subtotal	52	100%	102	100%	7	100%	2	100%	-	-	6	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	5	83%	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	17%	-	-	-	-	-	-	-	-
Subtotal	-	-	6	100%	-	-	-	-	-	-	-	-
<b>Total</b>	<b>415</b>		<b>472</b>		<b>45</b>		<b>6</b>		<b>1</b>		<b>40</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, Ohio

Exposure Category <sup>a</sup>	2010 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	259	62%	245	52%	23	51%	2	33%	-	-	26	65%
Injection drug use (IDU) only	6	1%	1	0%	1	2%	-	-	-	-	-	-
Heterosexual contact only	42	10%	99	21%	8	18%	3	50%	-	-	9	23%
MSM & IDU	12	3%	5	1%	1	2%	-	-	-	-	-	-
IDU & Heterosexual contact	7	2%	5	1%	1	2%	-	-	-	-	-	-
MSM & Heterosexual contact	6	1%	9	2%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	3	1%	1	0%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	5	1%	-	-	-	-	-	-	-	-
Other/unknown	80	19%	102	22%	11	24%	1	17%	1	100%	5	13%
<b>Total</b>	<b>415</b>		<b>472</b>		<b>45</b>		<b>6</b>		<b>1</b>		<b>40</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county**

County <sup>a</sup>	2010 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Adams	*	-	-	-	-
Allen	7.6	8	6	2	-
Ashland	*	-	-	-	-
Ashtabula	*	4	3	-	1
Athens	7.7	5	3	1	1
Auglaize	*	1	-	1	-
Belmont	*	1	-	1	-
Brown	*	1	1	-	-
Butler	4.8	18	8	9	1
Carroll	*	1	-	1	-
Champaign	*	3	-	2	1
Clark	4.4	6	2	3	1
Clermont	2.5	5	2	2	1
Clinton	*	-	-	-	-
Columbiana	*	2	2	-	-
Coshocton	*	1	1	-	-
Crawford	*	-	-	-	-
Cuyahoga	13.4	169	117	35	17
Darke	*	-	-	-	-
Defiance	*	4	3	1	-
Delaware	*	4	3	1	-
Erie	*	1	-	-	1
Fairfield	8.0	12	6	4	2
Fayette	*	-	-	-	-
Franklin	20.6	254	152	84	18
Fulton	*	2	2	-	-
Gallia	*	-	-	-	-
Geauga	*	2	-	1	1
Greene		12	8	2	2
Guernsey	*	-	-	-	-
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>601</b>	<b>288</b>	<b>90</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2010 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Hamilton	19.0	153	86	45	22
Hancock	*	-	-	-	-
Hardin	*	2	1	-	1
Harrison	*	1	-	1	-
Henry	*	2	1	1	-
Highland	*	1	-	1	-
Hocking	*	1	-	1	-
Holmes	*	-	-	-	-
Huron	*	1	1	-	-
Jackson	*	1	1	-	-
Jefferson	*	4	2	2	-
Knox	*	-	-	-	-
Lake	*	3	2	1	-
Lawrence	*	3	2	1	-
Licking	6.5	11	7	2	2
Logan	*	3	3	-	-
Lorain	4.9	15	9	5	1
Lucas	6.2	27	15	9	3
Madison	*	1	-	1	-
Mahoning	7.7	18	10	7	1
Marion	*	2	2	-	-
Medina	*	2	2	-	-
Meigs	*	-	-	-	-
Mercer	*	2	1	1	-
Miami	*	3	2	1	-
Monroe	*	-	-	-	-
Montgomery	14.1	75	50	17	8
Morgan	*	-	-	-	-
Morrow	*	-	-	-	-
Muskingum	*	3	-	2	1
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>601</b>	<b>288</b>	<b>90</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, Ohio, by county (*continued*)**

County <sup>a</sup>	2010 Diagnosis of HIV Infection		Disease Status at Initial Diagnosis		
	Rate <sup>b</sup>	No.	HIV (not AIDS)	HIV & later AIDS	AIDS
			No.	No.	No.
Noble	*	-	-	-	-
Ottawa	*	-	-	-	-
Paulding	*	-	-	-	-
Perry	*	-	-	-	-
Pickaway	*	-	-	-	-
Pike	*	1	1	-	-
Portage	*	4	2	1	1
Preble	*	2	1	1	-
Putnam	*	1	-	1	-
Richland	*	4	3	1	-
Ross	*	4	3	1	-
Sandusky	*	1	1	-	-
Scioto	*	1	1	-	-
Seneca	*	3	2	1	-
Shelby	*	2	1	1	-
Stark	7.5	28	19	7	2
Summit	4.8	26	17	9	-
Trumbull	2.9	6	3	3	-
Tuscarawas	*	1	-	1	-
Union	*	1	-	1	-
Van Wert	*	1	-	1	-
Vinton	*	-	-	-	-
Warren	2.3	5	3	1	1
Washington	*	-	-	-	-
Wayne	*	3	1	2	-
Williams	*	1	1	-	-
Wood	*	3	1	2	-
Wyandot	*	-	-	-	-
No County	*	31	25	6	-
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>601</b>	<b>288</b>	<b>90</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons newly diagnosed with an HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, Ohio

Characteristic	Diagnosis of HIV Infection by Year											
	2010			2011			2012			2013		
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%
<b>Sex</b>												
Males	14.3	809	83%	14.5	823	79%	14.8	839	82%	15.3	871	82%
Females	2.9	170	17%	3.7	220	21%	3.2	187	18%	3.2	189	18%
<b>Age at diagnosis (yr)</b>												
<13	0.3	5	1%	*	3	<1%	0.4	8	1%	0.5	9	1%
13-14	*	1	<1%	*	-	-	*	-	-	*	1	<1%
15-19	10.1	78	8%	8.3	64	6%	8.2	63	6%	8.3	64	6%
20-24	24.5	194	20%	27.8	220	21%	27.1	214	21%	29.6	234	22%
25-29	19.0	142	15%	20.7	155	15%	21.6	162	16%	23.1	173	16%
30-34	16.8	121	12%	18.8	136	13%	20.2	146	14%	16.2	117	11%
35-39	16.8	114	12%	16.8	114	11%	13.7	93	9%	15.0	102	10%
40-44	14.4	105	11%	15.4	112	11%	12.5	91	9%	13.5	98	9%
45-49	13.2	100	10%	12.2	92	9%	13.0	98	10%	12.8	97	9%
50-54	7.4	63	6%	9.4	80	8%	9.1	77	8%	8.1	69	7%
55-64	3.0	47	5%	3.4	54	5%	4.0	63	6%	4.8	76	7%
65+	0.5	9	1%	0.7	13	1%	0.6	11	1%	1.1	20	2%
<b>Race/Ethnicity</b>												
White, not Hispanic	4.5	415	42%	4.7	436	42%	4.5	418	41%	4.7	432	41%
Black/African American, not Hispanic	33.1	472	48%	35.5	506	49%	34.1	486	47%	35.8	511	48%
Hispanic/Latino	11.2	45	5%	14.9	60	6%	14.1	57	6%	15.4	62	6%
Asian/Pacific Islander	2.6	6	1%	3.0	7	1%	2.6	6	1%	3.4	8	1%
American Indian/Alaska Native	*	1	<1%	*	1	<1%	*	-	-	*	1	<1%
Unknown	*	40	4%	*	33	3%	*	59	6%	*	46	4%
<b>Race/Sex</b>												
White, not Hispanic Males	8.0	363	37%	8.0	366	35%	8.1	368	36%	8.2	375	35%
White, not Hispanic Females	1.1	52	5%	1.5	70	7%	1.1	50	5%	1.2	57	5%
Black/African American, not Hispanic Males	54.2	369	38%	54.6	372	36%	53.7	366	36%	59.9	408	38%
Black/African American, not Hispanic Females	13.8	103	11%	18.0	134	13%	16.1	120	12%	13.8	103	10%
Hispanic/Latino Males	18.3	38	4%	24.1	50	5%	25.1	52	5%	22.7	47	4%
Hispanic/Latina Females	3.6	7	1%	5.1	10	1%	2.6	5	<1%	7.7	15	1%
Asian/Pacific Islander Males	*	4	<1%	*	4	<1%	4.5	5	<1%	5.3	6	1%
Asian/Pacific Islander Females	*	2	<1%	*	3	<1%	*	1	<1%	*	2	<1%
American Indian/Alaska Native Males	*	1	<1%	*	1	<1%	*	-	-	*	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	40	4%	*	33	3%	*	59	6%	*	46	4%
<b>Total</b>	<b>8.4</b>	<b>979</b>		<b>9.0</b>	<b>1,043</b>		<b>8.8</b>	<b>1,026</b>		<b>9.1</b>	<b>1,060</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, Ohio

Transmission Category <sup>a</sup>	Diagnosis of HIV Infection by Year							
	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	570	71%	555	68%	572	68%	589	68%
Injection drug use (IDU)	8	1%	14	2%	18	2%	20	2%
Male-to-male sexual contact and IDU	22	3%	18	2%	25	3%	19	2%
Heterosexual contact	41	5%	33	4%	26	3%	37	4%
Other/unknown	163	20%	201	24%	195	23%	202	23%
Subtotal	804	100%	821	100%	836	100%	867	100%
<b>Female adult or adolescent</b>								
Injection drug use	13	8%	16	7%	8	4%	19	10%
Heterosexual contact	120	71%	141	64%	107	59%	113	62%
Other/unknown	36	21%	62	28%	65	36%	50	27%
Subtotal	169	100%	219	100%	180	100%	182	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	5	83%	3	100%	5	50%	4	36%
Other/unknown	1	17%	-	-	5	50%	7	64%
Subtotal	6	100%	3	100%	10	100%	11	100%
<b>Total</b>	<b>979</b>		<b>1,043</b>		<b>1,026</b>		<b>1,060</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, Ohio

Exposure Category <sup>a</sup>	Diagnosis of HIV Infection by Year							
	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	555	57%	539	52%	539	53%	563	53%
Injection drug use (IDU) only	8	1%	13	1%	18	2%	17	2%
Heterosexual contact only	161	16%	174	17%	133	13%	150	14%
MSM & IDU	18	2%	13	1%	17	2%	14	1%
IDU & Heterosexual contact	13	1%	17	2%	8	1%	22	2%
MSM & Heterosexual contact	15	2%	16	2%	33	3%	26	2%
MSM & IDU & Heterosexual contact	4	<1%	5	<1%	8	1%	5	<1%
Perinatal exposure	5	1%	3	<1%	7	1%	6	1%
Other/unknown	200	20%	263	25%	263	26%	257	24%
<b>Total</b>	<b>979</b>		<b>1,043</b>		<b>1,026</b>		<b>1,060</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and cumulative diagnoses, Ohio, by county

### Diagnosis of HIV Infection by Year

County <sup>a</sup>	2010		2011		2012		2013	
	Rate	No.	Rate	No.	Rate	No.	Rate	No.
Adams	*	-	*	-	*	-	*	1
Allen	7.6	8	6.7	7	6.7	7	12.4	13
Ashland	*	-	*	1	*	1	*	4
Ashtabula	*	4	*	1	*	1	5.0	5
Athens	7.7	5	*	3	*	1	*	-
Auglaize	*	1	*	-	*	-	*	-
Belmont	*	1	*	3	*	3	*	2
Brown	*	1	*	3	*	1	*	1
Butler	4.8	18	5.6	21	2.7	10	6.7	25
Carroll	*	1	*	-	*	2	*	1
Champaign	*	3	*	2	*	2	12.8	5
Clark	4.4	6	5.9	8	*	4	3.7	5
Clermont	2.5	5	5.0	10	4.0	8	4.5	9
Clinton	*	-	*	2	*	4	*	1
Columbiana	*	2	*	3	*	3	5.7	6
Coshocton	*	1	*	1	*	1	*	1
Crawford	*	-	*	2	*	1	*	2
Cuyahoga	13.4	169	15.7	198	16.5	208	18.1	228
Darke	*	-	*	3	*	1	*	2
Defiance	*	4	*	3	*	1	*	-
Delaware	*	4	3.2	6	6.9	13	3.7	7
Erie	*	1	*	3	*	2	6.6	5
Fairfield	8.0	12	4.7	7	*	4	*	4
Fayette	*	-	*	-	*	1	*	-
Franklin	20.6	254	21.2	261	20.0	246	18.4	227
Fulton	*	2	*	-	*	3	*	-
Gallia	*	-	*	1	*	1	*	1
Geauga	*	2	*	3	*	3	*	1
Greene	7.3	12	3.7	6	5.5	9	6.7	11
Guernsey	*	-	*	-	*	2	*	2
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>9.0</b>	<b>1,043</b>	<b>8.8</b>	<b>1,026</b>	<b>9.1</b>	<b>1060</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and cumulative diagnoses, Ohio, by county**

**Diagnosis of HIV Infection by Year**

County <sup>a</sup>	2010		2011		2012		2013	
	Rate	No.	Rate	No.	Rate	No.	Rate	No.
Hamilton	19.0	153	16.9	136	18.2	147	20.3	164
Hancock	*	-	*	1	*	-	*	4
Hardin	*	2	*	-	*	1	*	3
Harrison	*	1	*	1	*	-	*	-
Henry	*	2	*	1	*	2	*	1
Highland	*	1	*	2	*	1	*	-
Hocking	*	1	*	-	*	1	*	1
Holmes	*	-	*	-	*	-	*	-
Huron	*	1	*	3	*	-	*	2
Jackson	*	1	*	-	*	2	*	-
Jefferson	*	4	*	2	7.4	5	*	1
Knox	*	-	*	3	*	2	*	1
Lake	*	3	2.2	5	*	4	3.5	8
Lawrence	*	3	*	1	*	1	*	2
Licking	6.5	11	3.0	5	5.9	10	3.5	6
Logan	*	3	*	1	*	-	*	-
Lorain	4.9	15	5.3	16	5.6	17	4.9	15
Lucas	6.2	27	9.6	42	10.1	44	6.9	30
Madison	*	1	*	-	*	1	*	1
Mahoning	7.7	18	11.1	26	9.9	23	7.7	18
Marion	*	2	*	3	*	4	*	3
Medina	*	2	*	3	4.0	7	4.0	7
Meigs	*	-	*	1	*	-	*	-
Mercer	*	2	*	3	*	1	*	1
Miami	*	3	*	2	*	2	7.7	8
Monroe	*	-	*	-	*	-	*	-
Montgomery	14.1	75	14.4	77	11.8	63	12.2	65
Morgan	*	-	*	1	*	1	*	-
Morrow	*	-	*	2	*	2	*	3
Muskingum	*	3	*	2	*	2	*	3
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>9.0</b>	<b>1,043</b>	<b>8.8</b>	<b>1,026</b>	<b>9.1</b>	<b>1060</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and cumulative diagnoses, Ohio, by county**

**Diagnosis of HIV Infection by Year**

County <sup>a</sup>	2010		2011		2012		2013	
	Rate	No.	Rate	No.	Rate	No.	Rate	No.
Noble	*	-	*	-	*	-	*	-
Ottawa	*	-	*	2	*	2	*	1
Paulding	*	-	*	1	*	-	*	2
Perry	*	-	*	2	*	-	*	1
Pickaway	*	-	*	3	*	3	*	2
Pike	*	1	*	1	*	1	*	-
Portage	*	4	3.1	5	*	1	3.7	6
Preble	*	2	*	-	*	1	*	1
Putnam	*	1	*	1	*	-	*	-
Richland	*	4	*	1	*	4	*	4
Ross	*	4	7.8	6	*	2	*	1
Sandusky	*	1	8.3	5	*	3	*	1
Scioto	*	1	*	1	6.5	5	*	2
Seneca	*	3	*	-	*	-	*	1
Shelby	*	2	*	-	*	-	*	1
Stark	7.5	28	7.7	29	3.5	13	3.2	12
Summit	4.8	26	7.9	43	7.4	40	7.0	38
Trumbull	2.9	6	2.4	5	*	2	5.4	11
Tuscarawas	*	1	*	-	*	3	*	1
Union	*	1	*	1	*	1	*	-
Van Wert	*	1	*	-	*	-	*	-
Vinton	*	-	*	-	*	-	*	-
Warren	2.3	5	4.1	9	*	3	4.1	9
Washington	*	-	*	-	*	3	*	3
Wayne	*	3	*	1	*	4	*	1
Williams	*	1	*	1	*	-	*	1
Wood	*	3	*	1	4.6	6	6.2	8
Wyandot	*	-	*	-	*	-	*	-
No County	*	31	*	28	*	43	*	37
<b>Ohio</b>	<b>8.4</b>	<b>979</b>	<b>9.0</b>	<b>1,043</b>	<b>8.8</b>	<b>1,026</b>	<b>9.1</b>	<b>1060</b>

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2013) represent all reported cases diagnosed in each year.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Ohio**

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Sex</b>												
Males	443	80%	435	79%	387	80%	421	85%	309	82%	16,356	85%
Females	111	20%	113	21%	96	20%	76	15%	66	18%	2,996	15%
<b>Age at AIDS diagnosis (yr)</b>												
<13	2	<1%	1	<1%	1	<1%	-	-	-	-	148	1%
13-14	-	-	-	-	-	-	-	-	-	-	18	<1%
15-19	13	2%	11	2%	6	1%	5	1%	1	<1%	172	1%
20-24	49	9%	62	11%	51	11%	59	12%	42	11%	1,027	5%
25-29	71	13%	66	12%	50	10%	79	16%	45	12%	2,549	13%
30-34	81	15%	72	13%	67	14%	45	9%	57	15%	3,802	20%
35-39	67	12%	70	13%	56	12%	51	10%	36	10%	3,868	20%
40-44	93	17%	73	13%	70	14%	69	14%	45	12%	3,179	16%
45-49	86	16%	82	15%	68	14%	65	13%	48	13%	2,159	11%
50-54	49	9%	52	9%	55	11%	58	12%	43	11%	1,216	6%
55-64	35	6%	54	10%	54	11%	53	11%	48	13%	977	5%
65+	8	1%	5	1%	5	1%	13	3%	10	3%	222	1%
<b>Race/Ethnicity</b>												
White, not Hispanic	241	44%	232	42%	201	42%	216	43%	153	41%	10,498	54%
Black/African American, not Hispanic	278	50%	266	49%	220	46%	225	45%	185	49%	7,691	40%
Hispanic/Latino	20	4%	27	5%	32	7%	28	6%	20	5%	808	4%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	2	<1%	4	1%	50	<1%
American Indian/Alaska Native	-	-	1	<1%	-	-	-	-	1	<1%	16	<1%
Unknown	14	3%	21	4%	29	6%	26	5%	12	3%	289	1%
<b>Race/Sex</b>												
White, not Hispanic Males	213	38%	200	36%	175	36%	191	38%	133	35%	9,528	49%
White, not Hispanic Females	28	5%	32	6%	26	5%	25	5%	20	5%	970	5%
Black/African American, not Hispanic Males	200	36%	196	36%	159	33%	182	37%	142	38%	5,924	31%
Black/African American, not Hispanic Females	78	14%	70	13%	61	13%	43	9%	43	11%	1,767	9%
Hispanic/Latino Males	16	3%	21	4%	29	6%	24	5%	20	5%	622	3%
Hispanic/Latina Females	4	1%	6	1%	3	1%	4	1%	-	-	186	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	2	<1%	3	1%	41	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%	9	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	1	<1%	10	<1%
American Indian/Alaska Native Females	-	-	1	<1%	-	-	-	-	-	-	6	<1%
Unknown	14	3%	21	4%	29	6%	26	5%	12	3%	289	1%
<b>Total</b>	<b>554</b>		<b>548</b>		<b>483</b>		<b>497</b>		<b>375</b>		<b>19,352</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Ohio

Transmission Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	296	67%	278	64%	245	63%	281	67%	186	60%	11,101	68%
Injection drug use (IDU)	12	3%	8	2%	8	2%	10	2%	7	2%	1,152	7%
Male-to-male sexual contact and IDU	17	4%	13	3%	17	4%	10	2%	7	2%	1,020	6%
Heterosexual contact	26	6%	29	7%	14	4%	28	7%	28	9%	940	6%
Other/unknown	90	20%	106	24%	103	27%	90	21%	81	26%	2,042	13%
Subtotal	441	100%	434	100%	387	100%	419	100%	309	100%	16,255	100%
<b>Female adult or adolescent</b>												
Injection drug use	9	8%	8	7%	8	9%	6	8%	5	8%	537	19%
Heterosexual contact	80	73%	82	74%	61	65%	52	69%	40	61%	1982	68%
Other/unknown	20	18%	21	19%	25	27%	17	23%	21	32%	382	13%
Subtotal	109	100%	111	100%	94	100%	75	100%	66	100%	2,901	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	3	75%	3	100%	2	100%	1	33%	-	-	151	77%
Other/unknown	1	25%	-	-	-	-	2	67%	-	-	45	23%
Subtotal	4	100%	3	100%	2	100%	3	100%	-	-	196	100%
<b>Total</b>	<b>554</b>		<b>548</b>		<b>483</b>		<b>497</b>		<b>375</b>		<b>19,352</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Ohio

Exposure Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	280	51%	264	48%	231	48%	274	55%	182	49%	10,553	55%
Injection drug use (IDU) only	12	2%	11	2%	9	2%	8	2%	6	2%	1,026	5%
Heterosexual contact only	106	19%	111	20%	75	16%	80	16%	68	18%	2,922	15%
MSM & IDU	13	2%	12	2%	11	2%	8	2%	5	1%	821	4%
IDU & Heterosexual contact	9	2%	5	1%	7	1%	8	2%	6	2%	661	3%
MSM & Heterosexual contact	16	3%	14	3%	14	3%	7	1%	4	1%	543	3%
MSM & IDU & Heterosexual contact	4	1%	1	<1%	6	1%	2	<1%	2	1%	198	1%
Perinatal exposure	3	1%	3	1%	2	<1%	2	<1%	-	-	153	1%
Other/unknown	111	20%	127	23%	128	27%	108	22%	102	27%	2,475	13%
<b>Total</b>	<b>554</b>		<b>548</b>		<b>483</b>		<b>497</b>		<b>375</b>		<b>19,352</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses, Ohio, by county**

County <sup>a</sup>	Year of AIDS Diagnosis					Cumulative AIDS
	2010	2011	2012	2013	2014	
	No.	No.	No.	No.	No.	No.
Adams	-	-	-	1	-	13
Allen	3	4	5	7	4	132
Ashland	-	-	1	2	-	22
Ashtabula	1	1	1	1	-	65
Athens	2	2	1	-	1	47
Auglaize	1	-	-	-	-	22
Belmont	-	3	2	-	1	44
Brown	-	-	2	-	1	22
Butler	10	20	4	10	10	290
Carroll	-	-	-	1	1	13
Champaign	3	-	2	5	2	41
Clark	6	4	2	4	5	183
Clermont	4	3	6	1	5	94
Clinton	-	1	1	1	-	33
Columbiana	1	1	1	7	3	67
Coshocton	-	-	2	1	2	12
Crawford	-	1	2	2	-	29
Cuyahoga	91	83	75	91	52	4302
Darke	-	1	1	-	1	35
Defiance	1	2	1	1	-	29
Delaware	-	3	7	3	1	69
Erie	2	-	-	3	-	86
Fairfield	7	4	2	2	3	101
Fayette	-	-	-	-	-	19
Franklin	146	147	130	132	103	3778
Fulton	1	-	-	-	-	26
Gallia	-	-	-	-	1	14
Geauga	2	-	-	-	-	38
Greene	3	2	6	7	1	137
Guernsey	-	-	2	-	-	18
<b>Ohio</b>	<b>554</b>	<b>548</b>	<b>483</b>	<b>497</b>	<b>375</b>	<b>19,352</b>

*Note:* AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses, Ohio, by county (continued)**

County <sup>a</sup>	Year of AIDS Diagnosis					Cumulative AIDS
	2010	2011	2012	2013	2014	
	No.	No.	No.	No.	No.	No.
Hamilton	86	74	69	73	60	2506
Hancock	1	-	1	1	-	44
Hardin	1	-	-	2	1	12
Harrison	-	2	-	-	-	8
Henry	1	1	-	-	-	10
Highland	2	2	-	-	-	28
Hocking	1	-	1	-	-	20
Holmes	-	-	-	1	-	11
Huron	-	2	-	-	-	25
Jackson	-	-	1	-	-	14
Jefferson	3	1	4	1	1	68
Knox	-	-	-	-	-	21
Lake	-	5	-	3	2	154
Lawrence	1	1	-	1	-	43
Licking	6	4	8	4	1	140
Logan	-	1	-	-	-	24
Lorain	6	10	8	6	4	286
Lucas	24	30	11	12	6	895
Madison	1	-	-	-	-	27
Mahoning	12	17	11	5	11	401
Marion	-	2	3	3	1	68
Medina	-	-	3	3	-	49
Meigs	-	-	-	-	-	5
Mercer	1	2	-	1	-	18
Miami	1	1	4	2	2	82
Monroe	-	-	-	1	-	4
Montgomery	32	31	36	29	30	1292
Morgan	-	-	-	2	1	8
Morrow	-	1	-	-	1	18
Muskingum	4	-	2	3	1	76
<b>Ohio</b>	<b>554</b>	<b>548</b>	<b>483</b>	<b>497</b>	<b>375</b>	<b>19,352</b>

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses, Ohio, by county (continued)**

County <sup>a</sup>	Year of AIDS Diagnosis					Cumulative AIDS No.
	2010 No.	2011 No.	2012 No.	2013 No.	2014 No.	
Noble	-	-	-	-	-	1
Ottawa	1	-	-	-	-	17
Paulding	1	-	-	-	-	10
Perry	-	-	-	-	1	17
Pickaway	-	2	-	1	-	65
Pike	-	1	1	1	-	23
Portage	4	4	-	-	-	84
Preble	1	-	2	-	-	21
Putnam	-	1	-	-	1	10
Richland	3	1	3	3	-	134
Ross	3	2	1	1	2	75
Sandusky	1	1	3	1	1	50
Scioto	-	-	3	1	2	53
Seneca	-	1	-	-	-	30
Shelby	1	-	1	1	2	32
Stark	11	14	3	5	8	367
Summit	12	13	11	12	11	766
Trumbull	1	4	1	5	4	150
Tuscarawas	-	-	2	2	-	23
Union	2	-	1	-	-	24
Van Wert	1	-	-	-	-	9
Vinton	-	-	-	-	-	5
Warren	4	5	1	5	2	91
Washington	-	-	-	1	-	56
Wayne	2	1	1	-	-	44
Williams	-	1	1	1	1	26
Wood	2	-	3	2	1	67
Wyandot	-	-	-	-	-	5
No County	37	28	27	20	20	959
<b>Ohio</b>	<b>554</b>	<b>548</b>	<b>483</b>	<b>497</b>	<b>375</b>	<b>19,352</b>

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015, by selected characteristics, Ohio

Characteristic	AIDS Diagnoses					
	2014 AIDS Diagnoses		≥ 12 Months after HIV Diagnosis		< 12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Sex</b>						
Males	309	82%	104	81%	205	83%
Females	66	18%	25	19%	41	17%
<b>Age at AIDS diagnosis (yr)</b>						
<13	-	-	-	-	-	-
13-14	-	-	-	-	-	-
15-19	1	<1%	-	-	1	<1%
20-24	42	11%	9	7%	33	13%
25-29	45	12%	16	12%	29	12%
30-34	57	15%	15	12%	42	17%
35-39	36	10%	13	10%	23	9%
40-44	45	12%	20	16%	25	10%
45-49	48	13%	12	9%	36	15%
50-54	43	11%	25	19%	18	7%
55-64	48	13%	17	13%	31	13%
65+	10	3%	2	2%	8	3%
<b>Race/Ethnicity</b>						
White, not Hispanic	153	41%	51	40%	102	41%
Black/African American, not Hispanic	185	49%	64	50%	121	49%
Hispanic/Latino	20	5%	3	2%	17	7%
Asian/Pacific Islander	4	1%	2	2%	2	1%
American Indian/Alaska Native	1	<1%	1	1%	-	-
Unknown	12	3%	8	6%	4	2%
<b>Race/Sex</b>						
White, not Hispanic Males	133	35%	44	34%	89	36%
White, not Hispanic Females	20	5%	7	5%	13	5%
Black/African American, not Hispanic Males	142	38%	48	37%	94	38%
Black/African American, not Hispanic Females	43	11%	16	12%	27	11%
Hispanic/Latino Males	20	5%	3	2%	17	7%
Hispanic/Latina Females	-	-	-	-	-	-
Asian/Pacific Islander Males	3	1%	1	1%	2	1%
Asian/Pacific Islander Females	1	<1%	1	1%	-	-
American Indian/Alaska Native Males	1	<1%	1	1%	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Unknown	12	3%	8	6%	4	2%
<b>Total</b>	<b>375</b>		<b>129</b>		<b>246</b>	

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015, by transmission category, Ohio

Transmission Category <sup>a</sup>	AIDS Diagnoses					
	2014 AIDS Diagnoses		≥ 12 Months after HIV Diagnosis		< 12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	186	60%	54	52%	132	64%
Injection drug use (IDU)	7	2%	3	3%	4	2%
Male-to-male sexual contact and IDU	7	2%	7	7%	-	-
Heterosexual contact	28	9%	15	14%	13	6%
Other/unknown	81	26%	25	24%	56	27%
Subtotal	309	100%	104	100%	205	100%
<b>Female adult or adolescent</b>						
Injection drug use	5	8%	2	8%	3	7%
Heterosexual contact	40	61%	15	60%	25	61%
Other/unknown	21	32%	8	32%	13	32%
Subtotal	66	100%	25	100%	41	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>375</b>		<b>129</b>		<b>246</b>	

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015, by exposure category, Ohio

Transmission Category	AIDS Diagnoses					
	2014 AIDS Diagnoses		≥ 12 Months after HIV Diagnosis		< 12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	182	49%	53	41%	129	52%
Injection drug use (IDU) only	6	2%	3	2%	3	1%
Heterosexual contact only	68	18%	30	23%	38	15%
MSM & IDU	5	1%	5	4%	-	-
IDU & Heterosexual contact	6	2%	2	2%	4	2%
MSM & Heterosexual contact	4	1%	1	1%	3	1%
MSM & IDU & Heterosexual contact	2	1%	2	2%	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	102	27%	33	26%	69	28%
<b>Total</b>	<b>375</b>		<b>129</b>		<b>246</b>	

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015,  
Ohio, by county**

County <sup>b</sup>	AIDS Diagnoses		
	2014 AIDS Diagnoses	>= 12 Months after Diagnosis with HIV	< 12 Months after Diagnosis with HIV <sup>a</sup>
	No.	No.	No.
Adams	-	-	-
Allen	4	1	3
Ashland	-	-	-
Ashtabula	-	-	-
Athens	1	1	-
Auglaize	-	-	-
Belmont	1	-	1
Brown	1	-	1
Butler	10	3	7
Carroll	1	1	-
Champaign	2	-	2
Clark	5	3	2
Clermont	5	1	4
Clinton	-	-	-
Columbiana	3	-	3
Coshocton	2	-	2
Crawford	-	-	-
Cuyahoga	52	16	36
Darke	1	1	-
Defiance	-	-	-
Delaware	1	-	1
Erie	-	-	-
Fairfield	3	-	3
Fayette	-	-	-
Franklin	103	34	69
Fulton	-	-	-
Gallia	1	-	1
Geauga	-	-	-
Greene	1	1	-
Guernsey	-	-	-
<b>Ohio</b>	<b>375</b>	<b>129</b>	<b>246</b>

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

<sup>b</sup>County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015,  
Ohio, by county (continued)**

County <sup>b</sup>	AIDS Diagnoses		
	2014 AIDS Diagnoses	>= 12 Months after Diagnosis with HIV	< 12 Months after Diagnosis with HIV <sup>a</sup>
	No.	No.	No.
Hamilton	60	20	40
Hancock	-	-	-
Hardin	1	-	1
Harrison	-	-	-
Henry	-	-	-
Highland	-	-	-
Hocking	-	-	-
Holmes	-	-	-
Huron	-	-	-
Jackson	-	-	-
Jefferson	1	-	1
Knox	-	-	-
Lake	2	-	2
Lawrence	-	-	-
Licking	1	-	1
Logan	-	-	-
Lorain	4	2	2
Lucas	6	1	5
Madison	-	-	-
Mahoning	11	2	9
Marion	1	-	1
Medina	-	-	-
Meigs	-	-	-
Mercer	-	-	-
Miami	2	1	1
Monroe	-	-	-
Montgomery	30	13	17
Morgan	1	-	1
Morrow	1	-	1
Muskingum	1	-	1
<b>Ohio</b>	<b>375</b>	<b>129</b>	<b>246</b>

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

<sup>b</sup>County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Time to an AIDS diagnosis after a diagnosis of HIV infection as of June 30, 2015,  
Ohio, by county (continued)**

County <sup>b</sup>	AIDS Diagnoses		
	2014 AIDS Diagnoses	>= 12 Months after Diagnosis with HIV	< 12 Months after Diagnosis with HIV <sup>a</sup>
	No.	No.	No.
Noble	-	-	-
Ottawa	-	-	-
Paulding	-	-	-
Perry	1	-	1
Pickaway	-	-	-
Pike	-	-	-
Portage	-	-	-
Preble	-	-	-
Putnam	1	1	-
Richland	-	-	-
Ross	2	2	-
Sandusky	1	-	1
Scioto	2	1	1
Seneca	-	-	-
Shelby	2	-	2
Stark	8	3	5
Summit	11	7	4
Trumbull	4	-	4
Tuscarawas	-	-	-
Union	-	-	-
Van Wert	-	-	-
Vinton	-	-	-
Warren	2	1	1
Washington	-	-	-
Wayne	-	-	-
Williams	1	-	1
Wood	1	-	1
Wyandot	-	-	-
No County	20	13	7
<b>Ohio</b>	<b>375</b>	<b>129</b>	<b>246</b>

Note: Data reflects all persons with an AIDS diagnosis in 2014 regardless of year of HIV infection diagnosis.

<sup>a</sup>Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

Dash (-) indicates no cases were reported for the given category.

<sup>b</sup>County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

# Persons Living with a Diagnosis of HIV and/or AIDS Reported in Ohio

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, Ohio**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	303.2	17,204	80%	8,572	78%	8,632	81%
Females	74.5	4,408	20%	2,358	22%	2,050	19%
<b>Age at end of year</b>							
<13	3.5	65	<1%	48	<1%	17	<1%
13-14	4.3	13	<1%	7	<1%	6	<1%
15-19	17.6	136	1%	115	1%	21	<1%
20-24	122.4	968	4%	772	7%	196	2%
25-29	225.4	1,688	8%	1,188	11%	500	5%
30-34	237.6	1,715	8%	1,085	10%	630	6%
35-39	288.3	1,959	9%	1,094	10%	865	8%
40-44	354.4	2,578	12%	1,278	12%	1,300	12%
45-49	467.0	3,531	16%	1,566	14%	1,965	18%
50-54	451.4	3,823	18%	1,589	15%	2,234	21%
55-64	258.2	4,070	19%	1,735	16%	2,335	22%
65+	59.2	1,066	5%	453	4%	613	6%
<b>Race/Ethnicity</b>							
White, not Hispanic	107.6	9,995	46%	4,993	46%	5,002	47%
Black/African American, not Hispanic	670.1	9,558	44%	4,873	45%	4,685	44%
Hispanic/Latino	310.0	1,250	6%	600	5%	650	6%
Asian/Pacific Islander	43.2	101	<1%	58	1%	43	<1%
American Indian/Alaska Native	95.5	21	<1%	11	<1%	10	<1%
Unknown	*	687	3%	395	4%	292	3%
<b>Race/Sex</b>							
White, not Hispanic Males	189.4	8,626	40%	4,224	39%	4,402	41%
White, not Hispanic Females	28.9	1,369	6%	769	7%	600	6%
Black/African American, not Hispanic Males	1,027.7	6,999	32%	3,539	32%	3,460	32%
Black/African American, not Hispanic Females	343.3	2,559	12%	1,334	12%	1,225	11%
Hispanic/Latino Males	450.9	934	4%	444	4%	490	5%
Hispanic/Latina Females	161.2	316	1%	156	1%	160	1%
Asian/Pacific Islander Males	74.8	84	<1%	48	<1%	36	<1%
Asian/Pacific Islander Females	14.0	17	<1%	10	<1%	7	<1%
American Indian/Alaska Native Males	126.9	14	<1%	8	<1%	6	<1%
American Indian/Alaska Native Females	63.9	7	<1%	3	<1%	4	<1%
Unknown	*	687	3%	395	4%	292	3%
<b>Total</b>	<b>186.4</b>	<b>21,612</b>		<b>10,930</b>		<b>10,682</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	11,389	67%	5,542	65%	5,847	68%
Injection drug use (IDU)	770	5%	288	3%	482	6%
Male-to-male sexual contact and IDU	803	5%	297	3%	506	6%
Heterosexual contact	1,145	7%	472	6%	673	8%
Other/unknown	2,983	17%	1,907	22%	1,076	13%
Subtotal	17,090	100%	8,506	100%	8,584	100%
<b>Female adult or adolescent</b>						
Injection drug use	498	12%	217	10%	281	14%
Heterosexual contact	2,873	67%	1,426	63%	1,447	73%
Other/unknown	899	21%	636	28%	263	13%
Subtotal	4,270	100%	2,279	100%	1,991	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	202	80%	115	79%	87	81%
Other/unknown	50	20%	30	21%	20	19%
Subtotal	252	100%	145	100%	107	100%
<b>Total</b>	<b>21,612</b>		<b>10,930</b>		<b>10,682</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	10,742	50%	5,281	48%	5,461	51%
Injection drug use (IDU) only	657	3%	258	2%	399	4%
Heterosexual contact only	4,018	19%	1,898	17%	2,120	20%
MSM & IDU	610	3%	234	2%	376	4%
IDU & Heterosexual contact	611	3%	247	2%	364	3%
MSM & Heterosexual contact	646	3%	260	2%	386	4%
MSM & IDU & Heterosexual contact	193	1%	63	1%	130	1%
Perinatal exposure	213	1%	122	1%	91	1%
Other/unknown	3,922	18%	2,567	23%	1,355	13%
<b>Total</b>	<b>21,612</b>		<b>10,930</b>		<b>10,682</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	6,429	75%	4,082	59%	485	53%	48	58%	9	64%	335	61%
Injection drug use (IDU)	240	3%	380	5%	136	15%	-	-	2	14%	12	2%
Male-to-male sexual contact and IDU	439	5%	286	4%	51	6%	2	2%	-	-	25	5%
Heterosexual contact	308	4%	703	10%	90	10%	6	7%	-	-	38	7%
Other/unknown	1,179	14%	1,481	21%	159	17%	27	33%	3	21%	135	25%
Subtotal	8,595	100%	6,932	100%	921	100%	83	100%	14	100%	545	100%
<b>Female adult or adolescent</b>												
Injection drug use	198	15%	238	10%	51	17%	1	7%	-	-	10	8%
Heterosexual contact	894	67%	1,691	68%	196	65%	10	67%	5	71%	77	58%
Other/unknown	249	19%	545	22%	53	18%	4	27%	2	29%	46	35%
Subtotal	1,341	100%	2,474	100%	300	100%	15	100%	7	100%	133	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	42	71%	127	84%	26	90%	1	33%	-	-	6	67%
Other/unknown	17	29%	25	16%	3	10%	2	67%	-	-	3	33%
Subtotal	59	100%	152	100%	29	100%	3	100%	-	-	9	100%
<b>Total</b>	<b>9,995</b>		<b>9,558</b>		<b>1,250</b>		<b>101</b>		<b>21</b>		<b>687</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	6,198	62%	3,739	39%	444	36%	44	44%	9	43%	308	45%
Injection drug use (IDU) only	231	2%	318	3%	98	8%	1	1%	2	10%	7	1%
Heterosexual contact only	1,202	12%	2,394	25%	286	23%	16	16%	5	24%	115	17%
MSM & IDU	343	3%	211	2%	35	3%	2	2%	-	-	19	3%
IDU & Heterosexual contact	207	2%	300	3%	89	7%	-	-	-	-	15	2%
MSM & Heterosexual contact	231	2%	343	4%	41	3%	4	4%	-	-	27	4%
MSM & IDU & Heterosexual contact	96	1%	75	1%	16	1%	-	-	-	-	6	1%
Perinatal exposure	44	0%	133	1%	28	2%	1	1%	-	-	7	1%
Other/unknown	1,443	14%	2,045	21%	213	17%	33	33%	5	24%	183	27%
<b>Total</b>	<b>9,995</b>		<b>9,558</b>		<b>1,250</b>		<b>101</b>		<b>21</b>		<b>687</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Adams	92.4	26	12	14
Allen	152.3	160	78	82
Ashland	39.6	21	7	14
Ashtabula	99.8	99	43	56
Athens	72.6	47	20	27
Auglaize	48.0	22	6	16
Belmont	64.8	45	18	27
Brown	65.7	29	11	18
Butler	91.1	341	145	196
Carroll	31.9	9	3	6
Champaign	94.6	37	12	25
Clark	93.0	127	55	72
Clermont	64.0	129	62	67
Clinton	110.0	46	20	26
Columbiana	68.1	72	33	39
Coshocton	57.5	21	8	13
Crawford	77.7	33	14	19
Cuyahoga	347.7	4381	2255	2126
Darke	61.3	32	16	16
Defiance	64.9	25	9	16
Delaware	76.7	145	82	63
Erie	102.9	78	37	41
Fairfield	97.8	147	69	78
Fayette	90.3	26	10	16
Franklin	377.0	4642	2498	2144
Fulton	54.0	23	10	13
Gallia	69.1	21	8	13
Geauga	36.1	34	13	21
Greene	92.8	152	76	76
Guernsey	58.1	23	10	13
<b>Ohio</b>	<b>186.4</b>	<b>21,612</b>	<b>10,930</b>	<b>10,682</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Hamilton	343.0	2767	1365	1402
Hancock	53.1	40	18	22
Hardin	47.2	15	9	6
Harrison	38.6	6	3	3
Henry	39.4	11	7	4
Highland	55.8	24	8	16
Hocking	73.1	21	10	11
Holmes	*	4	2	2
Huron	44.3	26	14	12
Jackson	67.2	22	11	11
Jefferson	85.7	58	27	31
Knox	49.0	30	14	16
Lake	65.4	150	67	83
Lawrence	92.5	57	28	29
Licking	93.9	159	71	88
Logan	57.1	26	11	15
Lorain	113.7	346	164	182
Lucas	222.4	968	505	463
Madison	102.5	45	18	27
Mahoning	205.8	480	240	240
Marion	105.0	69	24	45
Medina	43.2	76	34	42
Meigs	60.0	14	9	5
Mercer	19.6	8	2	6
Miami	70.3	73	32	41
Monroe	48.4	7	2	5
Montgomery	244.8	1305	622	683
Morgan	121.3	18	6	12
Morrow	42.7	15	7	8
Muskingum	85.1	73	28	45
<b>Ohio</b>	<b>186.4</b>	<b>21,612</b>	<b>10,930</b>	<b>10,682</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Noble	69.6	10	9	1
Ottawa	36.4	15	10	5
Paulding	63.2	12	6	6
Perry	81.0	29	13	16
Pickaway	112.5	64	32	32
Pike	67.2	19	9	10
Portage	59.9	97	58	39
Preble	43.3	18	9	9
Putnam	14.6	5	3	2
Richland	88.6	108	50	58
Ross	90.7	70	30	40
Sandusky	73.1	44	15	29
Scioto	106.1	82	37	45
Seneca	37.7	21	7	14
Shelby	42.9	21	5	16
Stark	108.3	407	226	181
Summit	151.3	820	450	370
Trumbull	97.5	200	109	91
Tuscarawas	20.5	19	9	10
Union	94.8	51	26	25
Van Wert	38.6	11	2	9
Vinton	45.3	6	1	5
Warren	64.5	143	67	76
Washington	67.0	41	20	21
Wayne	61.5	71	38	33
Williams	61.7	23	11	12
Wood	50.9	66	32	34
Wyandot	31.3	7	4	3
No County	*	1226	644	582
<b>Ohio</b>	<b>186.4</b>	<b>21,612</b>	<b>10,930</b>	<b>10,682</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, Ohio**

Characteristic	Living with a diagnosis of HIV infection in 2013			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	289.1	16,407	79%	7,963	78%	8,444	81%
Females	72.2	4,272	21%	2,260	22%	2,012	19%
<b>Age at end of year</b>							
<13	3.8	72	<1%	52	1%	20	<1%
13-14	2.6	8	<1%	3	<1%	5	<1%
15-19	18.6	144	1%	116	1%	28	<1%
20-24	128.0	1,012	5%	761	7%	251	2%
25-29	202.6	1,517	7%	1,026	10%	491	5%
30-34	228.3	1,648	8%	1,002	10%	646	6%
35-39	279.7	1,901	9%	1,030	10%	871	8%
40-44	367.2	2,671	13%	1,274	12%	1,397	13%
45-49	484.9	3,666	18%	1,571	15%	2,095	20%
50-54	421.3	3,568	17%	1,494	15%	2,074	20%
55-64	228.1	3,595	17%	1,508	15%	2,087	20%
65+	48.7	877	4%	386	4%	491	5%
<b>Race/Ethnicity</b>							
White, not Hispanic	103.8	9,643	47%	4,733	46%	4,910	47%
Black/African American, not Hispanic	636.3	9,076	44%	4,506	44%	4,570	44%
Hispanic/Latino	293.9	1,185	6%	550	5%	635	6%
Asian/Pacific Islander	39.8	93	<1%	52	1%	41	<1%
American Indian/Alaska Native	91.0	20	<1%	10	<1%	10	<1%
Unknown	*	662	3%	372	4%	290	3%
<b>Race/Sex</b>							
White, not Hispanic Males	182.5	8,314	40%	3,992	39%	4,322	41%
White, not Hispanic Females	28.1	1,329	6%	741	7%	588	6%
Black/African American, not Hispanic Males	968.6	6,597	32%	3,226	32%	3,371	32%
Black/African American, not Hispanic Females	332.6	2,479	12%	1,280	13%	1,199	11%
Hispanic/Latino Males	424.3	879	4%	404	4%	475	5%
Hispanic/Latina Females	156.1	306	1%	146	1%	160	2%
Asian/Pacific Islander Males	69.5	78	<1%	44	<1%	34	<1%
Asian/Pacific Islander Females	12.4	15	<1%	8	<1%	7	<1%
American Indian/Alaska Native Males	126.9	14	<1%	8	<1%	6	<1%
American Indian/Alaska Native Females	54.8	6	<1%	2	<1%	4	<1%
Unknown	*	662	3%	372	4%	290	3%
<b>Total</b>	<b>178.4</b>	<b>20,679</b>		<b>10,223</b>		<b>10,456</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2013		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	10,840	67%	5,116	65%	5,724	68%
Injection drug use (IDU)	757	5%	277	4%	480	6%
Male-to-male sexual contact and IDU	783	5%	277	4%	506	6%
Heterosexual contact	1,109	7%	448	6%	661	8%
Other/unknown	2,805	17%	1,780	23%	1,025	12%
Subtotal	16,294	100%	7,898	100%	8,396	100%
<b>Female adult or adolescent</b>						
Injection drug use	483	12%	204	9%	279	14%
Heterosexual contact	2,791	67%	1,365	63%	1,426	73%
Other/unknown	863	21%	615	28%	248	13%
Subtotal	4,137	100%	2,184	100%	1,953	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	200	81%	113	80%	87	81%
Other/unknown	48	19%	28	20%	20	19%
Subtotal	248	100%	141	100%	107	100%
<b>Total</b>	<b>20,679</b>		<b>10,223</b>		<b>10,456</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2013		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	10,210	49%	4,870	48%	5,340	51%
Injection drug use (IDU) only	647	3%	249	2%	398	4%
Heterosexual contact only	3,900	19%	1,813	18%	2,087	20%
MSM & IDU	594	3%	218	2%	376	4%
IDU & Heterosexual contact	593	3%	232	2%	361	3%
MSM & Heterosexual contact	629	3%	245	2%	384	4%
MSM & IDU & Heterosexual contact	189	1%	59	1%	130	1%
Perinatal exposure	210	1%	119	1%	91	1%
Other/unknown	3,707	18%	2,418	24%	1,289	12%
<b>Total</b>	<b>20,679</b>		<b>10,223</b>		<b>10,456</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2013											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	6,214	75%	3,800	58%	451	52%	44	57%	9	64%	321	61%
Injection drug use (IDU)	230	3%	377	6%	136	16%	-	-	2	14%	12	2%
Male-to-male sexual contact and IDU	424	5%	285	4%	49	6%	2	3%	-	-	23	4%
Heterosexual contact	299	4%	679	10%	89	10%	6	8%	-	-	36	7%
Other/unknown	1,116	13%	1,390	21%	141	16%	25	32%	3	21%	131	25%
Subtotal	8,283	100%	6,531	100%	866	100%	77	100%	14	100%	523	100%
<b>Female adult or adolescent</b>												
Injection drug use	186	14%	238	10%	48	16%	1	7%	-	-	10	8%
Heterosexual contact	872	67%	1,640	68%	192	66%	9	64%	4	67%	74	57%
Other/unknown	243	19%	517	22%	51	18%	4	29%	2	33%	46	35%
Subtotal	1,301	100%	2,395	100%	291	100%	14	100%	6	100%	130	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	42	71%	126	84%	26	93%	-	-	-	-	6	67%
Other/unknown	17	29%	24	16%	2	7%	2	100%	-	-	3	33%
Subtotal	59	100%	150	100%	28	100%	2	100%	-	-	9	100%
<b>Total</b>	<b>9,643</b>		<b>9,076</b>		<b>1,185</b>		<b>93</b>		<b>20</b>		<b>662</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2013											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	5,988	62%	3,466	38%	412	35%	41	44%	9	45%	294	44%
Injection drug use (IDU) only	223	2%	317	3%	97	8%	1	1%	2	10%	7	1%
Heterosexual contact only	1,171	12%	2,319	26%	281	24%	15	16%	4	20%	110	17%
MSM & IDU	331	3%	210	2%	33	3%	2	2%	-	-	18	3%
IDU & Heterosexual contact	193	2%	298	3%	87	7%	-	-	-	-	15	2%
MSM & Heterosexual contact	226	2%	334	4%	39	3%	3	3%	-	-	27	4%
MSM & IDU & Heterosexual contact	93	1%	75	1%	16	1%	-	-	-	-	5	1%
Perinatal exposure	44	0%	132	1%	27	2%	-	-	-	-	7	1%
Other/unknown	1,374	14%	1,925	21%	193	16%	31	33%	5	25%	179	27%
<b>Total</b>	<b>9,643</b>		<b>9,076</b>		<b>1,185</b>		<b>93</b>		<b>20</b>		<b>662</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2013		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Adams	81.8	23	9	14
Allen	146.6	154	74	80
Ashland	37.7	20	6	14
Ashtabula	95.8	95	39	56
Athens	69.5	45	18	27
Auglaize	45.8	21	6	15
Belmont	64.8	45	18	27
Brown	61.2	27	9	18
Butler	87.1	326	136	190
Carroll	31.9	9	3	6
Champaign	84.3	33	9	24
Clark	90.1	123	52	71
Clermont	60.0	121	58	63
Clinton	107.6	45	19	26
Columbiana	63.4	67	30	37
Coshocton	54.8	20	8	12
Crawford	73.0	31	12	19
Cuyahoga	331.7	4179	2089	2090
Darke	55.6	29	13	16
Defiance	59.7	23	7	16
Delaware	73.5	139	77	62
Erie	97.6	74	33	41
Fairfield	94.4	142	65	77
Fayette	86.8	25	9	16
Franklin	359.8	4431	2352	2079
Fulton	54.0	23	10	13
Gallia	55.9	17	7	10
Geauga	33.9	32	11	21
Greene	87.3	143	67	76
Guernsey	55.6	22	9	13
<b>Ohio</b>	<b>178.4</b>	<b>20,679</b>	<b>10,223</b>	<b>10,456</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (continued)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2013		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Hamilton	324.3	2616	1250	1366
Hancock	53.1	40	18	22
Hardin	44.0	14	9	5
Harrison	38.6	6	3	3
Henry	39.4	11	7	4
Highland	55.8	24	8	16
Hocking	66.1	19	8	11
Holmes	*	4	2	2
Huron	44.3	26	14	12
Jackson	64.1	21	10	11
Jefferson	84.2	57	27	30
Knox	49.0	30	14	16
Lake	63.3	145	63	82
Lawrence	92.5	57	28	29
Licking	90.3	153	67	86
Logan	57.1	26	11	15
Lorain	111.8	340	159	181
Lucas	212.7	926	468	458
Madison	102.5	45	18	27
Mahoning	196.8	459	228	231
Marion	105.0	69	24	45
Medina	40.9	72	31	41
Meigs	60.0	14	9	5
Mercer	19.6	8	2	6
Miami	69.3	72	31	41
Monroe	48.4	7	2	5
Montgomery	234.7	1251	583	668
Morgan	114.5	17	6	11
Morrow	37.0	13	7	6
Muskingum	83.9	72	28	44
<b>Ohio</b>	<b>178.4</b>	<b>20,679</b>	<b>10,223</b>	<b>10,456</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2013		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Noble	55.7	8	7	1
Ottawa	36.4	15	10	5
Paulding	63.2	12	6	6
Perry	75.4	27	12	15
Pickaway	109.0	62	30	32
Pike	67.2	19	9	10
Portage	53.7	87	48	39
Preble	38.5	16	7	9
Putnam	14.6	5	3	2
Richland	88.6	108	50	58
Ross	86.8	67	28	39
Sandusky	69.8	42	14	28
Scioto	104.8	81	36	45
Seneca	37.7	21	7	14
Shelby	36.8	18	4	14
Stark	103.3	388	212	176
Summit	144.8	785	417	368
Trumbull	94.1	193	105	88
Tuscarawas	18.3	17	7	10
Union	93.0	50	25	25
Van Wert	38.6	11	2	9
Vinton	45.3	6	1	5
Warren	60.9	135	60	75
Washington	65.3	40	19	21
Wayne	59.7	69	36	33
Williams	53.6	20	9	11
Wood	49.4	64	31	33
Wyandot	31.3	7	4	3
No County	*	1208	634	574
<b>Ohio</b>	<b>178.4</b>	<b>20,679</b>	<b>10,223</b>	<b>10,456</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2013, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, Ohio**

Characteristic	Living with a diagnosis of HIV infection in 2012			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	274.1	15,553	79%	7,361	78%	8,192	81%
Females	69.0	4,083	21%	2,117	22%	1,966	19%
<b>Age at end of year</b>							
<13	3.5	66	<1%	43	<1%	23	<1%
13-14	3.6	11	<1%	7	<1%	4	<1%
15-19	20.2	156	1%	116	1%	40	<1%
20-24	121.3	959	5%	701	7%	258	3%
25-29	182.2	1,364	7%	877	9%	487	5%
30-34	222.6	1,607	8%	952	10%	655	6%
35-39	276.6	1,880	10%	966	10%	914	9%
40-44	384.6	2,798	14%	1,283	14%	1,515	15%
45-49	482.5	3,648	19%	1,536	16%	2,112	21%
50-54	389.8	3,301	17%	1,375	15%	1,926	19%
55-64	199.6	3,147	16%	1,318	14%	1,829	18%
65+	38.9	699	4%	304	3%	395	4%
<b>Race/Ethnicity</b>							
White, not Hispanic	99.3	9,224	47%	4,439	47%	4,785	47%
Black/African American, not Hispanic	600.8	8,570	44%	4,138	44%	4,432	44%
Hispanic/Latino	278.3	1,122	6%	509	5%	613	6%
Asian/Pacific Islander	36.0	84	<1%	47	<1%	37	<1%
American Indian/Alaska Native	86.4	19	<1%	9	<1%	10	<1%
Unknown	*	617	3%	336	4%	281	3%
<b>Race/Sex</b>							
White, not Hispanic Males	174.6	7,951	40%	3,742	39%	4,209	41%
White, not Hispanic Females	26.9	1,273	6%	697	7%	576	6%
Black/African American, not Hispanic Males	909.8	6,196	32%	2,935	31%	3,261	32%
Black/African American, not Hispanic Females	318.5	2,374	12%	1,203	13%	1,171	12%
Hispanic/Latino Males	401.2	831	4%	375	4%	456	4%
Hispanic/Latina Females	148.4	291	1%	134	1%	157	2%
Asian/Pacific Islander Males	63.2	71	<1%	40	<1%	31	<1%
Asian/Pacific Islander Females	10.7	13	<1%	7	<1%	6	<1%
American Indian/Alaska Native Males	117.8	13	<1%	7	<1%	6	<1%
American Indian/Alaska Native Females	54.8	6	<1%	2	<1%	4	<1%
Unknown	*	617	3%	336	4%	281	3%
<b>Total</b>	<b>169.4</b>	<b>19,636</b>		<b>9,478</b>		<b>10,158</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2012		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	10,252	66%	4,694	64%	5,558	68%
Injection drug use (IDU)	740	5%	265	4%	475	6%
Male-to-male sexual contact and IDU	764	5%	260	4%	504	6%
Heterosexual contact	1,073	7%	428	6%	645	8%
Other/unknown	2,615	17%	1,652	23%	963	12%
Subtotal	15,444	100%	7,299	100%	8,145	100%
<b>Female adult or adolescent</b>						
Injection drug use	464	12%	187	9%	277	15%
Heterosexual contact	2,678	68%	1,284	63%	1,394	73%
Other/unknown	813	21%	577	28%	236	12%
Subtotal	3,955	100%	2,048	100%	1,907	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	196	83%	109	83%	87	82%
Other/unknown	41	17%	22	17%	19	18%
Subtotal	237	100%	131	100%	106	100%
<b>Total</b>	<b>19,636</b>		<b>9,478</b>		<b>10,158</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2012		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	9,649	49%	4,471	47%	5,178	51%
Injection drug use (IDU) only	633	3%	238	3%	395	4%
Heterosexual contact only	3,751	19%	1,712	18%	2,039	20%
MSM & IDU	579	3%	204	2%	375	4%
IDU & Heterosexual contact	571	3%	214	2%	357	4%
MSM & Heterosexual contact	602	3%	222	2%	380	4%
MSM & IDU & Heterosexual contact	185	1%	56	1%	129	1%
Perinatal exposure	204	1%	114	1%	90	1%
Other/unknown	3,462	18%	2,247	24%	1,215	12%
<b>Total</b>	<b>19,636</b>		<b>9,478</b>		<b>10,158</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2012											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	5,936	75%	3,547	58%	419	51%	39	55%	8	62%	302	62%
Injection drug use (IDU)	220	3%	370	6%	136	17%	-	-	2	15%	12	2%
Male-to-male sexual contact and IDU	413	5%	283	5%	47	6%	2	3%	-	-	19	4%
Heterosexual contact	289	4%	659	11%	87	11%	6	8%	-	-	32	7%
Other/unknown	1,062	13%	1,274	21%	129	16%	24	34%	3	23%	124	25%
Subtotal	7,920	100%	6,133	100%	818	100%	71	100%	13	100%	489	100%
<b>Female adult or adolescent</b>												
Injection drug use	170	14%	237	10%	48	17%	1	8%	-	-	8	7%
Heterosexual contact	843	68%	1,570	68%	185	67%	8	62%	4	67%	68	57%
Other/unknown	233	19%	487	21%	43	16%	4	31%	2	33%	44	37%
Subtotal	1,246	100%	2,294	100%	276	100%	13	100%	6	100%	120	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	41	71%	123	86%	26	93%	-	-	-	-	6	75%
Other/unknown	17	29%	20	14%	2	7%	-	-	-	-	2	25%
Subtotal	58	100%	143	100%	28	100%	-	-	-	-	8	100%
<b>Total</b>	<b>9,224</b>		<b>8,570</b>		<b>1,122</b>		<b>84</b>		<b>19</b>		<b>617</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2012											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	5,720	62%	3,225	38%	383	34%	36	43%	8	42%	277	45%
Injection drug use (IDU) only	212	2%	314	4%	97	9%	1	1%	2	11%	7	1%
Heterosexual contact only	1,132	12%	2,229	26%	272	24%	14	17%	4	21%	100	16%
MSM & IDU	323	4%	208	2%	31	3%	2	2%	-	-	15	2%
IDU & Heterosexual contact	178	2%	293	3%	87	8%	-	-	-	-	13	2%
MSM & Heterosexual contact	216	2%	322	4%	36	3%	3	4%	-	-	25	4%
MSM & IDU & Heterosexual contact	90	1%	75	1%	16	1%	-	-	-	-	4	1%
Perinatal exposure	43	0%	127	1%	27	2%	-	-	-	-	7	1%
Other/unknown	1,310	14%	1,777	21%	173	15%	28	33%	5	26%	169	27%
<b>Total</b>	<b>9,224</b>		<b>8,570</b>		<b>1,122</b>		<b>84</b>		<b>19</b>		<b>617</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2012		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Adams	74.7	21	8	13
Allen	136.1	143	66	77
Ashland	35.8	19	6	13
Ashtabula	91.8	91	36	55
Athens	69.5	45	18	27
Auglaize	45.8	21	6	15
Belmont	63.3	44	17	27
Brown	61.2	27	9	18
Butler	81.8	306	123	183
Carroll	28.4	8	3	5
Champaign	71.6	28	8	20
Clark	87.1	119	50	69
Clermont	56.1	113	51	62
Clinton	102.8	43	17	26
Columbiana	57.7	61	29	32
Coshocton	52.0	19	7	12
Crawford	70.6	30	12	18
Cuyahoga	313.9	3954	1927	2027
Darke	51.7	27	11	16
Defiance	59.7	23	7	16
Delaware	70.9	134	74	60
Erie	91.0	69	30	39
Fairfield	91.1	137	63	74
Fayette	83.3	24	8	16
Franklin	341.0	4199	2190	2009
Fulton	54.0	23	10	13
Gallia	52.6	16	7	9
Geauga	33.9	32	11	21
Greene	82.4	135	62	73
Guernsey	53.0	21	8	13
<b>Ohio</b>	<b>169.4</b>	<b>19,636</b>	<b>9,478</b>	<b>10,158</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2012		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Hamilton	304.1	2453	1139	1314
Hancock	51.8	39	17	22
Hardin	34.6	11	8	3
Harrison	32.2	5	2	3
Henry	28.6	8	4	4
Highland	55.8	24	8	16
Hocking	62.7	18	7	11
Holmes	*	4	2	2
Huron	40.9	24	12	12
Jackson	64.1	21	10	11
Jefferson	81.2	55	25	30
Knox	47.4	29	13	16
Lake	59.3	136	56	80
Lawrence	92.5	57	28	29
Licking	88.0	149	63	86
Logan	57.1	26	11	15
Lorain	106.8	325	148	177
Lucas	206.3	898	446	452
Madison	102.5	45	18	27
Mahoning	189.1	441	214	227
Marion	103.5	68	23	45
Medina	37.5	66	27	39
Meigs	55.7	13	8	5
Mercer	17.1	7	2	5
Miami	62.6	65	25	40
Monroe	41.5	6	1	5
Montgomery	221.0	1178	528	650
Morgan	114.5	17	6	11
Morrow	34.1	12	6	6
Muskingum	80.4	69	26	43
<b>Ohio</b>	<b>169.4</b>	<b>19,636</b>	<b>9,478</b>	<b>10,158</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2012		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Noble	55.7	8	7	1
Ottawa	34.0	14	9	5
Paulding	52.7	10	4	6
Perry	72.6	26	11	15
Pickaway	107.3	61	29	32
Pike	67.2	19	9	10
Portage	50.7	82	43	39
Preble	36.1	15	6	9
Putnam	14.6	5	3	2
Richland	81.2	99	45	54
Ross	84.2	65	27	38
Sandusky	68.1	41	14	27
Scioto	98.4	76	32	44
Seneca	37.7	21	7	14
Shelby	34.7	17	4	13
Stark	100.1	376	203	173
Summit	138.0	748	385	363
Trumbull	89.7	184	99	85
Tuscarawas	18.3	17	7	10
Union	91.1	49	25	24
Van Wert	38.6	11	2	9
Vinton	45.3	6	1	5
Warren	56.4	125	56	69
Washington	60.4	37	16	21
Wayne	57.1	66	34	32
Williams	51.0	19	9	10
Wood	44.8	58	25	33
Wyandot	31.3	7	4	3
No County	*	1173	605	568
<b>Ohio</b>	<b>169.4</b>	<b>19,636</b>	<b>9,478</b>	<b>10,158</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2012. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2012, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, Ohio**

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	259.3	14,716	79%	6,800	77%	7,916	81%
Females	65.8	3,897	21%	1,985	23%	1,912	19%
<b>Age at end of year</b>							
<13	3.2	59	<1%	35	<1%	24	<1%
13-14	6.9	21	<1%	14	<1%	7	<1%
15-19	20.6	159	1%	115	1%	44	<1%
20-24	111.7	883	5%	609	7%	274	3%
25-29	166.0	1,243	7%	773	9%	470	5%
30-34	210.7	1,521	8%	853	10%	668	7%
35-39	276.0	1,876	10%	940	11%	936	10%
40-44	403.2	2,933	16%	1,329	15%	1,604	16%
45-49	484.9	3,666	20%	1,509	17%	2,157	22%
50-54	346.9	2,938	16%	1,205	14%	1,733	18%
55-64	174.5	2,751	15%	1,153	13%	1,598	16%
65+	31.3	563	3%	250	3%	313	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	94.9	8,818	47%	4,164	47%	4,654	47%
Black/African American, not Hispanic	566.6	8,083	43%	3,802	43%	4,281	44%
Hispanic/Latino	261.9	1,056	6%	468	5%	588	6%
Asian/Pacific Islander	33.8	79	<1%	43	<1%	36	<1%
American Indian/Alaska Native	81.9	18	<1%	8	<1%	10	<1%
Unknown	*	559	3%	300	3%	259	3%
<b>Race/Sex</b>							
White, not Hispanic Males	166.8	7,598	41%	3,511	40%	4,087	42%
White, not Hispanic Females	25.8	1,220	7%	653	7%	567	6%
Black/African American, not Hispanic Males	855.3	5,825	31%	2,676	30%	3,149	32%
Black/African American, not Hispanic Females	302.9	2,258	12%	1,126	13%	1,132	12%
Hispanic/Latino Males	372.2	771	4%	339	4%	432	4%
Hispanic/Latina Females	145.4	285	2%	129	1%	156	2%
Asian/Pacific Islander Males	59.7	67	<1%	37	<1%	30	<1%
Asian/Pacific Islander Females	9.9	12	<1%	6	<1%	6	<1%
American Indian/Alaska Native Males	108.8	12	<1%	6	<1%	6	<1%
American Indian/Alaska Native Females	54.8	6	<1%	2	<1%	4	<1%
Unknown	*	559	3%	300	3%	259	3%
<b>Total</b>	<b>160.5</b>	<b>18,613</b>		<b>8,785</b>		<b>9,828</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	9,669	66%	4,294	64%	5,375	68%
Injection drug use (IDU)	722	5%	253	4%	469	6%
Male-to-male sexual contact and IDU	738	5%	245	4%	493	6%
Heterosexual contact	1,044	7%	407	6%	637	8%
Other/unknown	2,440	17%	1,545	23%	895	11%
Subtotal	14,613	100%	6,744	100%	7,869	100%
<b>Female adult or adolescent</b>						
Injection drug use	453	12%	178	9%	275	15%
Heterosexual contact	2,572	68%	1,210	63%	1,362	73%
Other/unknown	751	20%	534	28%	217	12%
Subtotal	3,776	100%	1,922	100%	1,854	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	187	83%	101	85%	86	82%
Other/unknown	37	17%	18	15%	19	18%
Subtotal	224	100%	119	100%	105	100%
<b>Total</b>	<b>18,613</b>		<b>8,785</b>		<b>9,828</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	9,099	49%	4,097	47%	5,002	51%
Injection drug use (IDU) only	613	3%	224	3%	389	4%
Heterosexual contact only	3,616	19%	1,617	18%	1,999	20%
MSM & IDU	561	3%	192	2%	369	4%
IDU & Heterosexual contact	562	3%	207	2%	355	4%
MSM & Heterosexual contact	569	3%	196	2%	373	4%
MSM & IDU & Heterosexual contact	177	1%	53	1%	124	1%
Perinatal exposure	194	1%	105	1%	89	1%
Other/unknown	3,222	17%	2,094	24%	1,128	11%
<b>Total</b>	<b>18,613</b>		<b>8,785</b>		<b>9,828</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	5,675	75%	3,303	57%	375	49%	39	58%	7	58%	269	61%
Injection drug use (IDU)	212	3%	364	6%	133	18%	-	-	2	17%	11	2%
Male-to-male sexual contact and IDU	398	5%	276	5%	45	6%	2	3%	-	-	17	4%
Heterosexual contact	280	4%	641	11%	85	11%	6	9%	-	-	32	7%
Other/unknown	1,003	13%	1,182	20%	121	16%	20	30%	3	25%	112	25%
Subtotal	7,568	100%	5,766	100%	759	100%	67	100%	12	100%	441	100%
<b>Female adult or adolescent</b>												
Injection drug use	162	14%	235	11%	47	17%	1	8%	-	-	8	7%
Heterosexual contact	817	68%	1,500	69%	182	67%	7	58%	4	67%	62	56%
Other/unknown	215	18%	449	21%	41	15%	4	33%	2	33%	40	36%
Subtotal	1,194	100%	2,184	100%	270	100%	12	100%	6	100%	110	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	39	70%	117	88%	25	93%	-	-	-	-	6	75%
Other/unknown	17	30%	16	12%	2	7%	-	-	-	-	2	25%
Subtotal	56	100%	133	100%	27	100%	-	-	-	-	8	100%
<b>Total</b>	<b>8,818</b>		<b>8,083</b>		<b>1,056</b>		<b>79</b>		<b>18</b>		<b>559</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	5,474	62%	2,996	37%	340	32%	36	46%	7	39%	246	44%
Injection drug use (IDU) only	201	2%	309	4%	93	9%	1	1%	2	11%	7	1%
Heterosexual contact only	1,097	12%	2,141	26%	267	25%	13	16%	4	22%	94	17%
MSM & IDU	312	4%	203	3%	30	3%	2	3%	-	-	14	3%
IDU & Heterosexual contact	173	2%	290	4%	87	8%	-	-	-	-	12	2%
MSM & Heterosexual contact	201	2%	307	4%	35	3%	3	4%	-	-	23	4%
MSM & IDU & Heterosexual contact	86	1%	73	1%	15	1%	-	-	-	-	3	1%
Perinatal exposure	41	0%	120	1%	26	2%	-	-	-	-	7	1%
Other/unknown	1,233	14%	1,644	20%	163	15%	24	30%	5	28%	153	27%
<b>Total</b>	<b>8,818</b>		<b>8,083</b>		<b>1,056</b>		<b>79</b>		<b>18</b>		<b>559</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2011		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Adams	71.1	20	7	13
Allen	130.4	137	61	76
Ashland	33.9	18	6	12
Ashtabula	89.7	89	35	54
Athens	68.0	44	18	26
Auglaize	45.8	21	6	15
Belmont	57.6	40	15	25
Brown	58.9	26	9	17
Butler	78.6	294	117	177
Carroll	24.8	7	2	5
Champaign	66.4	26	6	20
Clark	84.9	116	49	67
Clermont	52.1	105	45	60
Clinton	88.4	37	13	24
Columbiana	55.8	59	27	32
Coshocton	49.3	18	7	11
Crawford	68.3	29	11	18
Cuyahoga	297.6	3749	1771	1978
Darke	51.7	27	11	16
Defiance	57.1	22	7	15
Delaware	67.2	127	69	58
Erie	87.0	66	27	39
Fairfield	86.4	130	58	72
Fayette	76.4	22	6	16
Franklin	322.1	3966	2034	1932
Fulton	49.3	21	8	13
Gallia	52.6	16	7	9
Geauga	30.8	29	8	21
Greene	79.4	130	61	69
Guernsey	48.0	19	7	12
<b>Ohio</b>	<b>160.5</b>	<b>18,613</b>	<b>8,785</b>	<b>9,828</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (continued)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2011		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Hamilton	285.3	2301	1046	1255
Hancock	50.4	38	17	21
Hardin	31.5	10	7	3
Harrison	32.2	5	2	3
Henry	21.5	6	2	4
Highland	55.8	24	8	16
Hocking	55.7	16	7	9
Holmes	*	4	2	2
Huron	39.2	23	11	12
Jackson	61.1	20	9	11
Jefferson	76.8	52	24	28
Knox	42.5	26	11	15
Lake	57.6	132	52	80
Lawrence	90.9	56	27	29
Licking	80.3	136	55	81
Logan	52.7	24	10	14
Lorain	100.6	306	137	169
Lucas	196.4	855	414	441
Madison	97.9	43	17	26
Mahoning	177.1	413	199	214
Marion	98.9	65	22	43
Medina	35.2	62	24	38
Meigs	55.7	13	8	5
Mercer	17.1	7	2	5
Miami	62.6	65	25	40
Monroe	41.5	6	1	5
Montgomery	209.9	1119	491	628
Morgan	107.8	16	6	10
Morrow	28.4	10	4	6
Muskingum	80.4	69	26	43
<b>Ohio</b>	<b>160.5</b>	<b>18,613</b>	<b>8,785</b>	<b>9,828</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2011		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Noble	55.7	8	7	1
Ottawa	31.6	13	8	5
Paulding	52.7	10	4	6
Perry	67.0	24	11	13
Pickaway	98.5	56	25	31
Pike	63.7	18	9	9
Portage	50.0	81	42	39
Preble	36.1	15	6	9
Putnam	14.6	5	3	2
Richland	76.3	93	42	51
Ross	80.4	62	25	37
Sandusky	61.5	37	12	25
Scioto	90.6	70	31	39
Seneca	37.7	21	7	14
Shelby	34.7	17	4	13
Stark	93.9	353	185	168
Summit	131.2	711	356	355
Trumbull	88.7	182	97	85
Tuscarawas	16.2	15	6	9
Union	85.5	46	23	23
Van Wert	38.6	11	2	9
Vinton	45.3	6	1	5
Warren	55.0	122	53	69
Washington	55.5	34	13	21
Wayne	50.2	58	30	28
Williams	51.0	19	9	10
Wood	42.4	55	22	33
Wyandot	31.3	7	4	3
No County	*	1132	574	558
<b>Ohio</b>	<b>160.5</b>	<b>18,613</b>	<b>8,785</b>	<b>9,828</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2011, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, Ohio**

Characteristic	Living with a diagnosis of HIV infection in 2010			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	245.0	13,906	79%	6,265	77%	7,641	81%
Females	62.3	3,687	21%	1,850	23%	1,837	19%
<b>Age at end of year</b>							
<13	3.3	61	<1%	36	<1%	25	<1%
13-14	8.3	25	<1%	19	<1%	6	<1%
15-19	22.8	176	1%	119	1%	57	1%
20-24	98.6	780	4%	507	6%	273	3%
25-29	149.6	1,120	6%	679	8%	441	5%
30-34	202.0	1,458	8%	785	10%	673	7%
35-39	288.7	1,962	11%	955	12%	1,007	11%
40-44	413.5	3,008	17%	1,296	16%	1,712	18%
45-49	471.4	3,564	20%	1,460	18%	2,104	22%
50-54	314.3	2,662	15%	1,101	14%	1,561	16%
55-64	147.1	2,319	13%	956	12%	1,363	14%
65+	25.5	458	3%	202	2%	256	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	90.3	8,390	48%	3,874	48%	4,516	48%
Black/African American, not Hispanic	532.2	7,592	43%	3,487	43%	4,105	43%
Hispanic/Latino	248.3	1,001	6%	433	5%	568	6%
Asian/Pacific Islander	31.3	73	<1%	38	<1%	35	<1%
American Indian/Alaska Native	77.3	17	<1%	7	<1%	10	<1%
Unknown	*	520	3%	276	3%	244	3%
<b>Race/Sex</b>							
White, not Hispanic Males	159.0	7,240	41%	3,271	40%	3,969	42%
White, not Hispanic Females	24.3	1,150	7%	603	7%	547	6%
Black/African American, not Hispanic Males	801.4	5,458	31%	2,436	30%	3,022	32%
Black/African American, not Hispanic Females	286.3	2,134	12%	1,051	13%	1,083	11%
Hispanic/Latino Males	350.5	726	4%	310	4%	416	4%
Hispanic/Latina Females	140.3	275	2%	123	2%	152	2%
Asian/Pacific Islander Males	57.0	64	<1%	35	<1%	29	<1%
Asian/Pacific Islander Females	7.4	9	<1%	3	<1%	6	<1%
American Indian/Alaska Native Males	99.7	11	<1%	5	<1%	6	<1%
American Indian/Alaska Native Females	54.8	6	<1%	2	<1%	4	<1%
Unknown	*	520	3%	276	3%	244	3%
<b>Total</b>	<b>151.7</b>	<b>17,593</b>		<b>8,115</b>		<b>9,478</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2010		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	9,107	66%	3,920	63%	5,187	68%
Injection drug use (IDU)	709	5%	244	4%	465	6%
Male-to-male sexual contact and IDU	719	5%	232	4%	487	6%
Heterosexual contact	1,013	7%	388	6%	625	8%
Other/unknown	2,257	16%	1,427	23%	830	11%
Subtotal	13,805	100%	6,211	100%	7,594	100%
<b>Female adult or adolescent</b>						
Injection drug use	440	12%	171	10%	269	15%
Heterosexual contact	2,436	68%	1,123	63%	1,313	74%
Other/unknown	692	19%	494	28%	198	11%
Subtotal	3,568	100%	1,788	100%	1,780	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	183	83%	98	84%	85	82%
Other/unknown	37	17%	18	16%	19	18%
Subtotal	220	100%	116	100%	104	100%
<b>Total</b>	<b>17,593</b>		<b>8,115</b>		<b>9,478</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2010		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	8,552	49%	3,735	46%	4,817	51%
Injection drug use (IDU) only	600	3%	216	3%	384	4%
Heterosexual contact only	3,449	20%	1,511	19%	1,938	20%
MSM & IDU	548	3%	185	2%	363	4%
IDU & Heterosexual contact	549	3%	199	2%	350	4%
MSM & Heterosexual contact	554	3%	184	2%	370	4%
MSM & IDU & Heterosexual contact	171	1%	47	1%	124	1%
Perinatal exposure	190	1%	102	1%	88	1%
Other/unknown	2,980	17%	1,936	24%	1,044	11%
<b>Total</b>	<b>17,593</b>		<b>8,115</b>		<b>9,478</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2010											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	5,412	75%	3,062	57%	348	49%	38	59%	7	64%	239	59%
Injection drug use (IDU)	207	3%	361	7%	129	18%	-	-	2	18%	10	2%
Male-to-male sexual contact and IDU	382	5%	273	5%	45	6%	2	3%	-	-	17	4%
Heterosexual contact	271	4%	622	12%	83	12%	6	9%	-	-	31	8%
Other/unknown	938	13%	1,083	20%	109	15%	18	28%	2	18%	108	27%
Subtotal	7,210	100%	5,401	100%	714	100%	64	100%	11	100%	405	100%
<b>Female adult or adolescent</b>												
Injection drug use	153	14%	231	11%	47	18%	1	11%	-	-	8	7%
Heterosexual contact	773	69%	1,417	69%	176	67%	6	67%	4	67%	60	56%
Other/unknown	198	18%	413	20%	38	15%	2	22%	2	33%	39	36%
Subtotal	1,124	100%	2,061	100%	261	100%	9	100%	6	100%	107	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	39	70%	114	88%	24	92%	-	-	-	-	6	75%
Other/unknown	17	30%	16	12%	2	8%	-	-	-	-	2	25%
Subtotal	56	100%	130	100%	26	100%	-	-	-	-	8	100%
<b>Total</b>	<b>8,390</b>		<b>7,592</b>		<b>1,001</b>		<b>73</b>		<b>17</b>		<b>520</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2010											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	5,215	62%	2,764	36%	314	31%	35	48%	7	41%	217	42%
Injection drug use (IDU) only	193	2%	307	4%	91	9%	1	1%	2	12%	6	1%
Heterosexual contact only	1,044	12%	2,039	27%	259	26%	12	16%	4	24%	91	18%
MSM & IDU	300	4%	202	3%	30	3%	2	3%	-	-	14	3%
IDU & Heterosexual contact	167	2%	285	4%	85	8%	-	-	-	-	12	2%
MSM & Heterosexual contact	197	2%	298	4%	34	3%	3	4%	-	-	22	4%
MSM & IDU & Heterosexual contact	82	1%	71	1%	15	1%	-	-	-	-	3	1%
Perinatal exposure	41	0%	117	2%	25	2%	-	-	-	-	7	1%
Other/unknown	1,151	14%	1,509	20%	148	15%	20	27%	4	24%	148	28%
<b>Total</b>	<b>8,390</b>		<b>7,592</b>		<b>1,001</b>		<b>73</b>		<b>17</b>		<b>520</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2010		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS)	AIDS
			No.	No.
Adams	71.1	20	7	13
Allen	122.8	129	56	73
Ashland	32.1	17	6	11
Ashtabula	87.7	87	35	52
Athens	64.9	42	18	24
Auglaize	39.3	18	4	14
Belmont	53.3	37	15	22
Brown	47.6	21	5	16
Butler	74.0	277	111	166
Carroll	24.8	7	2	5
Champaign	63.9	25	6	19
Clark	79.8	109	45	64
Clermont	48.6	98	42	56
Clinton	86.1	36	12	24
Columbiana	53.0	56	26	30
Coshocton	46.6	17	6	11
Crawford	65.9	28	11	17
Cuyahoga	282.9	3564	1634	1930
Darke	46.0	24	9	15
Defiance	51.9	20	7	13
Delaware	63.5	120	64	56
Erie	83.1	63	24	39
Fairfield	79.8	120	52	68
Fayette	69.4	20	5	15
Franklin	301.5	3713	1870	1843
Fulton	49.3	21	8	13
Gallia	49.3	15	6	9
Geauga	29.7	28	7	21
Greene	75.1	123	57	66
Guernsey	48.0	19	7	12
<b>Ohio</b>	<b>151.7</b>	<b>17,593</b>	<b>8,115</b>	<b>9,478</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2010		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Hamilton	267.8	2160	954	1206
Hancock	50.4	38	17	21
Hardin	31.5	10	7	3
Harrison	*	4	2	2
Henry	17.9	5	2	3
Highland	48.8	21	7	14
Hocking	52.2	15	6	9
Holmes	*	4	2	2
Huron	32.4	19	8	11
Jackson	61.1	20	9	11
Jefferson	73.9	50	23	27
Knox	42.5	26	11	15
Lake	55.4	127	49	78
Lawrence	87.6	54	26	28
Licking	77.9	132	54	78
Logan	48.3	22	9	13
Lorain	97.3	296	130	166
Lucas	187.2	815	390	425
Madison	95.6	42	17	25
Mahoning	165.9	387	181	206
Marion	91.3	60	18	42
Medina	33.5	59	21	38
Meigs	47.1	11	7	4
Mercer	17.1	7	2	5
Miami	57.7	60	22	38
Monroe	*	4	0	4
Montgomery	196.6	1048	448	600
Morgan	101.1	15	6	9
Morrow	22.8	8	4	4
Muskingum	79.2	68	26	42
<b>Ohio</b>	<b>151.7</b>	<b>17,593</b>	<b>8,115</b>	<b>9,478</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status, Ohio, by county (*continued*)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2010		Current Disease Status	
	Rate <sup>b</sup>	No.	HIV (not AIDS) No.	AIDS No.
Noble	41.8	6	6	0
Ottawa	26.7	11	6	5
Paulding	47.4	9	3	6
Perry	64.2	23	10	13
Pickaway	96.7	55	24	31
Pike	60.2	17	9	8
Portage	47.6	77	40	37
Preble	36.1	15	6	9
Putnam	14.6	5	3	2
Richland	73.0	89	39	50
Ross	73.9	57	22	35
Sandusky	54.8	33	9	24
Scioto	85.4	66	27	39
Seneca	37.7	21	7	14
Shelby	34.7	17	4	13
Stark	87.0	327	167	160
Summit	124.2	673	325	348
Trumbull	83.3	171	87	84
Tuscarawas	16.2	15	6	9
Union	83.7	45	22	23
Van Wert	38.6	11	2	9
Vinton	45.3	6	1	5
Warren	50.1	111	46	65
Washington	53.9	33	13	20
Wayne	47.6	55	27	28
Williams	48.3	18	9	9
Wood	41.7	54	22	32
Wyandot	26.8	6	3	3
No County	*	1106	555	551
<b>Ohio</b>	<b>151.7</b>	<b>17,593</b>	<b>8,115</b>	<b>9,478</b>

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2010. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2010, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

<sup>b</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, Ohio

### Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	13,906	79%	14,716	79%	15,553	79%	16,407	79%	17,204	80%
Females	3,687	21%	3,897	21%	4,083	21%	4,272	21%	4,408	20%
<b>Age at end of year</b>										
<13	61	<1%	59	<1%	66	<1%	72	<1%	65	<1%
13-14	25	<1%	21	<1%	11	<1%	8	<1%	13	<1%
15-19	176	1%	159	1%	156	1%	144	1%	136	1%
20-24	780	4%	883	5%	959	5%	1,012	5%	968	4%
25-29	1,120	6%	1,243	7%	1,364	7%	1,517	7%	1,688	8%
30-34	1,458	8%	1,521	8%	1,607	8%	1,648	8%	1,715	8%
35-39	1,962	11%	1,876	10%	1,880	10%	1,901	9%	1,959	9%
40-44	3,008	17%	2,933	16%	2,798	14%	2,671	13%	2,578	12%
45-49	3,564	20%	3,666	20%	3,648	19%	3,666	18%	3,531	16%
50-54	2,662	15%	2,938	16%	3,301	17%	3,568	17%	3,823	18%
55-64	2,319	13%	2,751	15%	3,147	16%	3,595	17%	4,070	19%
65+	458	3%	563	3%	699	4%	877	4%	1,066	5%
<b>Race/Ethnicity</b>										
White, not Hispanic	8,390	48%	8,818	47%	9,224	47%	9,643	47%	9,995	46%
Black/African American, not Hispanic	7,592	43%	8,083	43%	8,570	44%	9,076	44%	9,558	44%
Hispanic/Latino	1,001	6%	1,056	6%	1,122	6%	1,185	6%	1,250	6%
Asian/Pacific Islander	73	<1%	79	<1%	84	<1%	93	<1%	101	<1%
American Indian/Alaska Native	17	<1%	18	<1%	19	<1%	20	<1%	21	<1%
Unknown	520	3%	559	3%	617	3%	662	3%	687	3%
<b>Race/Sex</b>										
White, not Hispanic Males	7,240	41%	7,598	41%	7,951	40%	8,314	40%	8,626	40%
White, not Hispanic Females	1,150	7%	1,220	7%	1,273	6%	1,329	6%	1,369	6%
Black/African American, not Hispanic Males	5,458	31%	5,825	31%	6,196	32%	6,597	32%	6,999	32%
Black/African American, not Hispanic Females	2,134	12%	2,258	12%	2,374	12%	2,479	12%	2,559	12%
Hispanic/Latino Males	726	4%	771	4%	831	4%	879	4%	934	4%
Hispanic/Latina Females	275	2%	285	2%	291	1%	306	1%	316	1%
Asian/Pacific Islander Males	64	<1%	67	<1%	71	<1%	78	<1%	84	<1%
Asian/Pacific Islander Females	9	<1%	12	<1%	13	<1%	15	<1%	17	<1%
American Indian/Alaska Native Males	11	<1%	12	<1%	13	<1%	14	<1%	14	<1%
American Indian/Alaska Native Females	6	<1%	6	<1%	6	<1%	6	<1%	7	<1%
Unknown	520	3%	559	3%	617	3%	662	3%	687	3%
<b>Total</b>	<b>17,593</b>		<b>18,613</b>		<b>19,636</b>		<b>20,679</b>		<b>21,612</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, Ohio

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	9,107	66%	9,669	66%	10,252	66%	10,840	67%	11,389	67%
Injection drug use (IDU)	709	5%	722	5%	740	5%	757	5%	770	5%
Male-to-male sexual contact and IDU	719	5%	738	5%	764	5%	783	5%	803	5%
Heterosexual contact	1,013	7%	1,044	7%	1,073	7%	1,109	7%	1,145	7%
Other/unknown	2,257	16%	2,440	17%	2,615	17%	2,805	17%	2,983	17%
Subtotal	13,805	100%	14,613	100%	15,444	100%	16,294	100%	17,090	100%
<b>Female adult or adolescent</b>										
Injection drug use	440	12%	453	12%	464	12%	483	12%	498	12%
Heterosexual contact	2,436	68%	2,572	68%	2,678	68%	2,791	67%	2,873	67%
Other/unknown	692	19%	751	20%	813	21%	863	21%	899	21%
Subtotal	3,568	100%	3,776	100%	3,955	100%	4,137	100%	4,270	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	183	83%	187	83%	196	83%	200	81%	202	80%
Other/unknown	37	17%	37	17%	41	17%	48	19%	50	20%
Subtotal	220	100%	224	100%	237	100%	248	100%	252	100%
<b>Total</b>	<b>17,593</b>		<b>18,613</b>		<b>19,636</b>		<b>20,679</b>		<b>21,612</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, Ohio

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	8,552	49%	9,099	49%	9,649	49%	10,210	49%	10,742	50%
Injection drug use (IDU) only	600	3%	613	3%	633	3%	647	3%	657	3%
Heterosexual contact only	3,449	20%	3,616	19%	3,751	19%	3,900	19%	4,018	19%
MSM & IDU	548	3%	561	3%	579	3%	594	3%	610	3%
IDU & Heterosexual contact	549	3%	562	3%	571	3%	593	3%	611	3%
MSM & Heterosexual contact	554	3%	569	3%	602	3%	629	3%	646	3%
MSM & IDU & Heterosexual contact	171	1%	177	1%	185	1%	189	1%	193	1%
Perinatal exposure	190	1%	194	1%	204	1%	210	1%	213	1%
Other/unknown	2,980	17%	3,222	17%	3,462	18%	3,707	18%	3,922	18%
<b>Total</b>	<b>17,593</b>		<b>18,613</b>		<b>19,636</b>		<b>20,679</b>		<b>21,612</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year, Ohio, by county**

<b>Living with a Diagnosis of HIV Infection by Year</b>					
<b>County<sup>a</sup></b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>
	<b>No.</b>	<b>No.</b>	<b>No.</b>	<b>No.</b>	<b>No.</b>
Adams	20	20	21	23	26
Allen	129	137	143	154	160
Ashland	17	18	19	20	21
Ashtabula	87	89	91	95	99
Athens	42	44	45	45	47
Auglaize	18	21	21	21	22
Belmont	37	40	44	45	45
Brown	21	26	27	27	29
Butler	277	294	306	326	341
Carroll	7	7	8	9	9
Champaign	25	26	28	33	37
Clark	109	116	119	123	127
Clermont	98	105	113	121	129
Clinton	36	37	43	45	46
Columbiana	56	59	61	67	72
Coshocton	17	18	19	20	21
Crawford	28	29	30	31	33
Cuyahoga	3564	3749	3954	4179	4381
Darke	24	27	27	29	32
Defiance	20	22	23	23	25
Delaware	120	127	134	139	145
Erie	63	66	69	74	78
Fairfield	120	130	137	142	147
Fayette	20	22	24	25	26
Franklin	3713	3966	4199	4431	4642
Fulton	21	21	23	23	23
Gallia	15	16	16	17	21
Geauga	28	29	32	32	34
Greene	123	130	135	143	152
Guernsey	19	19	21	22	23
<b>Ohio</b>	<b>17,593</b>	<b>18,613</b>	<b>19,636</b>	<b>20,679</b>	<b>21,612</b>

*Note:* Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year, Ohio, by county (continued)**

<b>Living with a Diagnosis of HIV Infection by Year</b>					
<b>County<sup>a</sup></b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>
	<b>No.</b>	<b>No.</b>	<b>No.</b>	<b>No.</b>	<b>No.</b>
Hamilton	2160	2301	2453	2616	2767
Hancock	38	38	39	40	40
Hardin	10	10	11	14	15
Harrison	4	5	5	6	6
Henry	5	6	8	11	11
Highland	21	24	24	24	24
Hocking	15	16	18	19	21
Holmes	4	4	4	4	4
Huron	19	23	24	26	26
Jackson	20	20	21	21	22
Jefferson	50	52	55	57	58
Knox	26	26	29	30	30
Lake	127	132	136	145	150
Lawrence	54	56	57	57	57
Licking	132	136	149	153	159
Logan	22	24	26	26	26
Lorain	296	306	325	340	346
Lucas	815	855	898	926	968
Madison	42	43	45	45	45
Mahoning	387	413	441	459	480
Marion	60	65	68	69	69
Medina	59	62	66	72	76
Meigs	11	13	13	14	14
Mercer	7	7	7	8	8
Miami	60	65	65	72	73
Monroe	4	6	6	7	7
Montgomery	1048	1119	1178	1251	1305
Morgan	15	16	17	17	18
Morrow	8	10	12	13	15
Muskingum	68	69	69	72	73
<b>Ohio</b>	<b>17,593</b>	<b>18,613</b>	<b>19,636</b>	<b>20,679</b>	<b>21,612</b>

*Note:* Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year, Ohio, by county (continued)**

County <sup>a</sup>	Living with a Diagnosis of HIV Infection by Year				
	2010 No.	2011 No.	2012 No.	2013 No.	2014 No.
Noble	6	8	8	8	10
Ottawa	11	13	14	15	15
Paulding	9	10	10	12	12
Perry	23	24	26	27	29
Pickaway	55	56	61	62	64
Pike	17	18	19	19	19
Portage	77	81	82	87	97
Preble	15	15	15	16	18
Putnam	5	5	5	5	5
Richland	89	93	99	108	108
Ross	57	62	65	67	70
Sandusky	33	37	41	42	44
Scioto	66	70	76	81	82
Seneca	21	21	21	21	21
Shelby	17	17	17	18	21
Stark	327	353	376	388	407
Summit	673	711	748	785	820
Trumbull	171	182	184	193	200
Tuscarawas	15	15	17	17	19
Union	45	46	49	50	51
Van Wert	11	11	11	11	11
Vinton	6	6	6	6	6
Warren	111	122	125	135	143
Washington	33	34	37	40	41
Wayne	55	58	66	69	71
Williams	18	19	19	20	23
Wood	54	55	58	64	66
Wyandot	6	7	7	7	7
No County	1106	1132	1173	1208	1226
<b>Ohio</b>	<b>17,593</b>	<b>18,613</b>	<b>19,636</b>	<b>20,679</b>	<b>21,612</b>

*Note:* Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Deaths among Persons Reported  
with a Diagnosis of HIV and/or AIDS  
in Ohio

## Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths by selected characteristics, Ohio

### Year of Death among Persons with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014		Cumulative Deaths	
	No.	%	No.	%								
<b>Sex</b>												
Males	223	82%	222	80%	232	78%	195	82%	147	83%	10,476	87%
Females	48	18%	55	20%	66	22%	44	18%	31	17%	1,525	13%
<b>Age at death (yr)</b>												
<13	-	-	-	-	-	-	-	-	-	-	81	1%
13-14	-	-	-	-	-	-	-	-	-	-	3	<1%
15-19	-	-	1	<1%	1	<1%	-	-	-	-	33	<1%
20-24	11	4%	7	3%	1	<1%	1	<1%	4	2%	200	2%
25-29	7	3%	7	3%	9	3%	10	4%	11	6%	1,000	8%
30-34	16	6%	15	5%	12	4%	12	5%	9	5%	1,884	16%
35-39	20	7%	14	5%	20	7%	10	4%	7	4%	2,297	19%
40-44	40	15%	40	14%	32	11%	30	13%	19	11%	2,151	18%
45-49	65	24%	57	21%	52	17%	52	22%	25	14%	1,688	14%
50-54	50	18%	49	18%	48	16%	40	17%	33	19%	1,144	10%
55-64	50	18%	62	22%	85	29%	59	25%	57	32%	1,111	9%
65+	12	4%	25	9%	38	13%	25	10%	13	7%	409	3%
<b>Race/Ethnicity</b>												
White, not Hispanic	159	59%	133	48%	154	52%	118	49%	93	52%	7,154	60%
Black/African American, not Hispanic	94	35%	126	45%	115	39%	99	41%	73	41%	4,347	36%
Hispanic/Latino	12	4%	12	4%	9	3%	12	5%	6	3%	407	3%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	-	-	-	-	20	<1%
American Indian/Alaska Native	2	1%	-	-	1	<1%	-	-	-	-	13	<1%
Unknown	3	1%	5	2%	18	6%	10	4%	6	3%	60	<1%
<b>Race/Sex</b>												
White, not Hispanic Males	134	49%	116	42%	124	42%	103	43%	83	47%	6,588	55%
White, not Hispanic Females	25	9%	17	6%	30	10%	15	6%	10	6%	566	5%
Black/African American, not Hispanic Males	76	28%	94	34%	82	28%	74	31%	54	30%	3,494	29%
Black/African American, not Hispanic Females	18	7%	32	12%	33	11%	25	10%	19	11%	853	7%
Hispanic/Latino Males	9	3%	8	3%	7	2%	10	4%	6	3%	316	3%
Hispanic/Latina Females	3	1%	4	1%	2	1%	2	1%	-	-	91	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	-	-	-	-	20	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	<1%	-	-	-	-	-	-	-	-	9	<1%
American Indian/Alaska Native Females	1	<1%	-	-	1	<1%	-	-	-	-	4	<1%
Unknown	3	1%	5	2%	18	6%	10	4%	6	3%	60	<1%
<b>Total</b>	<b>271</b>		<b>277</b>		<b>298</b>		<b>239</b>		<b>178</b>		<b>12,001</b>	

Notes: Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths by transmission category, Ohio

Transmission Category <sup>a</sup>	Year of Death among Persons with a Diagnosis of HIV Infection										Cumulative Deaths	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	120	54%	112	50%	118	51%	113	58%	77	53%	6,740	65%
Injection drug use (IDU)	21	10%	19	9%	16	7%	15	8%	15	10%	964	9%
Male-to-male sexual contact and IDU	15	7%	19	9%	18	8%	11	6%	12	8%	766	7%
Heterosexual contact	16	7%	26	12%	13	6%	15	8%	13	9%	408	4%
Other/unknown	49	22%	46	21%	67	29%	41	21%	29	20%	1,534	15%
Subtotal	221	100%	222	100%	232	100%	195	100%	146	100%	10,412	100%
<b>Female adult or adolescent</b>												
Injection drug use	10	22%	10	19%	11	17%	8	18%	6	19%	390	26%
Heterosexual contact	27	59%	33	61%	39	59%	29	66%	20	65%	826	56%
Other/unknown	9	20%	11	20%	16	24%	7	16%	5	16%	259	18%
Subtotal	46	100%	54	100%	66	100%	44	100%	31	100%	1,475	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	4	100%	1	100%	-	-	-	-	1	100%	85	75%
Other/unknown	-	-	-	-	-	-	-	-	-	-	29	25%
Subtotal	4	100%	1	100%	-	-	-	-	1	100%	114	100%
<b>Total</b>	<b>271</b>		<b>277</b>		<b>298</b>		<b>239</b>		<b>178</b>		<b>12,001</b>	

Notes: Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths by exposure category, Ohio

Exposure Category <sup>a</sup>	Year of Death among Persons with a Diagnosis of HIV Infection										Cumulative Deaths	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	113	42%	109	39%	108	36%	105	44%	73	41%	6,506	54%
Injection drug use (IDU) only	19	7%	15	5%	19	6%	8	3%	8	4%	876	7%
Heterosexual contact only	43	16%	59	21%	52	17%	44	18%	33	19%	1,234	10%
MSM & IDU	13	5%	12	4%	15	5%	10	4%	10	6%	636	5%
IDU & Heterosexual contact	12	4%	14	5%	8	3%	15	6%	13	7%	476	4%
MSM & Heterosexual contact	7	3%	3	1%	10	3%	8	3%	4	2%	227	2%
MSM & IDU & Heterosexual contact	2	1%	7	3%	3	1%	1	<1%	2	1%	129	1%
Perinatal exposure	4	1%	1	<1%	-	-	-	-	1	1%	85	1%
Other/unknown	58	21%	57	21%	83	28%	48	20%	34	19%	1,832	15%
<b>Total</b>	<b>271</b>		<b>277</b>		<b>298</b>		<b>239</b>		<b>178</b>		<b>12,001</b>	

Notes: Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths, Ohio, by county**

**Year of Death among Persons with a Diagnosis of HIV Infection**

County <sup>a</sup>	2010	2011	2012	2013	2014	Cumulative Deaths
	No.	No.	No.	No.	No.	No.
Adams	-	-	-	-	-	11
Allen	4	1	4	1	4	89
Ashland	-	1	-	-	-	11
Ashtabula	2	1	1	1	-	39
Athens	-	-	1	-	-	30
Auglaize	1	-	-	-	-	14
Belmont	1	-	1	-	-	29
Brown	-	-	3	1	-	13
Butler	4	3	4	5	1	155
Carroll	-	-	1	-	-	8
Champaign	3	1	1	1	-	25
Clark	4	1	2	3	2	124
Clermont	-	1	-	-	2	54
Clinton	-	-	1	1	-	21
Columbiana	4	-	-	1	3	48
Coshocton	-	-	1	-	1	9
Crawford	-	-	1	-	-	20
Cuyahoga	54	62	65	55	21	2733
Darke	-	-	-	-	-	21
Defiance	2	1	-	-	1	20
Delaware	2	1	1	1	2	38
Erie	2	1	3	2	1	60
Fairfield	-	1	3	2	3	52
Fayette	-	-	-	1	-	14
Franklin	51	56	59	44	25	2235
Fulton	2	-	-	1	1	17
Gallia	-	-	1	-	-	9
Geauga	-	-	-	-	1	26
Greene	1	2	-	2	1	78
Guernsey	2	1	-	-	-	11
<b>Ohio</b>	<b>271</b>	<b>277</b>	<b>298</b>	<b>239</b>	<b>178</b>	<b>12,001</b>

Notes: Deaths among persons with a diagnosis of HIV infection represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths, Ohio, by county (continued)**

**Year of Death among Persons with a Diagnosis of HIV Infection**

County <sup>a</sup>	2010	2011	2012	2013	2014	Cumulative Deaths
	No.	No.	No.	No.	No.	No.
Hamilton	29	27	34	37	23	1483
Hancock	1	2	-	2	-	32
Hardin	-	-	-	-	1	9
Harrison	-	-	-	-	-	5
Henry	-	2	-	-	-	9
Highland	-	1	1	-	-	15
Hocking	1	-	-	-	-	11
Holmes	-	-	1	-	-	5
Huron	1	-	-	-	-	14
Jackson	2	-	2	-	-	15
Jefferson	3	1	2	1	-	57
Knox	-	-	-	-	-	17
Lake	1	-	-	-	-	89
Lawrence	-	-	-	-	1	29
Licking	2	-	-	2	-	73
Logan	-	-	-	-	-	19
Lorain	7	1	4	3	3	185
Lucas	16	16	14	7	4	583
Madison	-	-	-	-	-	20
Mahoning	6	9	9	5	3	271
Marion	-	1	1	1	-	39
Medina	-	-	-	-	2	23
Meigs	-	-	-	-	-	6
Mercer	-	-	-	-	2	12
Miami	1	-	2	1	-	46
Monroe	-	-	-	-	-	4
Montgomery	10	14	15	13	18	776
Morgan	-	-	-	1	-	2
Morrow	-	-	-	1	-	16
Muskingum	4	-	1	-	2	41
<b>Ohio</b>	<b>271</b>	<b>277</b>	<b>298</b>	<b>239</b>	<b>178</b>	<b>12,001</b>

Notes: Deaths among persons with a diagnosis of HIV infection represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported deaths among persons with a diagnosis of HIV infection as of June 30, 2015, by year of death and cumulative deaths, Ohio, by county (continued)**

**Year of Death among Persons with a Diagnosis of HIV Infection**

County <sup>a</sup>	2010	2011	2012	2013	2014	Cumulative Deaths
	No.	No.	No.	No.	No.	No.
Noble	-	-	-	-	-	1
Ottawa	1	-	-	-	-	15
Paulding	-	1	1	-	-	10
Perry	1	-	-	1	-	12
Pickaway	-	-	1	1	1	54
Pike	-	-	1	-	-	13
Portage	3	1	1	1	1	56
Preble	-	-	-	-	-	12
Putnam	-	-	1	1	-	9
Richland	5	1	3	-	2	100
Ross	2	1	3	-	-	49
Sandusky	2	-	1	-	1	27
Scioto	-	2	-	1	1	36
Seneca	-	-	-	1	-	24
Shelby	-	-	-	1	-	19
Stark	5	10	7	4	3	270
Summit	7	16	11	7	20	504
Trumbull	1	6	2	4	2	109
Tuscarawas	-	-	1	1	-	17
Union	-	-	-	-	-	17
Van Wert	-	-	-	1	-	4
Vinton	-	-	-	-	-	3
Warren	1	3	4	-	1	43
Washington	1	-	1	-	2	49
Wayne	2	-	1	1	1	30
Williams	-	2	-	-	-	19
Wood	-	1	2	1	1	46
Wyandot	-	-	-	-	-	5
No County	17	25	18	17	14	528
<b>Ohio</b>	<b>271</b>	<b>277</b>	<b>298</b>	<b>239</b>	<b>178</b>	<b>12,001</b>

Notes: Deaths among persons with a diagnosis of HIV infection represent deaths attributed to any cause among persons reported with a diagnosis of HIV or AIDS. Deaths by year (2010-2014) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> County reflects county of residence at time of earliest diagnosis. Cases diagnosed while in a correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.