

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Allen

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Allen

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	213.5	113	71%	51	65%	62	76%
Females	90.2	47	29%	27	35%	20	24%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	2	1%	1	1%	1	1%
20-24	*	4	3%	2	3%	2	2%
25-29	205.4	13	8%	9	12%	4	5%
30-34	226.6	14	9%	9	12%	5	6%
35-39	309.1	19	12%	9	12%	10	12%
40-44	429.8	26	16%	10	13%	16	20%
45-49	385.1	24	15%	9	12%	15	18%
50-54	351.3	26	16%	14	18%	12	15%
55-64	174.5	25	16%	11	14%	14	17%
65+	41.9	7	4%	4	5%	3	4%
Race/Ethnicity							
White, not Hispanic	94.6	81	51%	40	51%	41	50%
Black/African American, not Hispanic	462.0	59	37%	27	35%	32	39%
Hispanic/Latino	379.6	11	7%	6	8%	5	6%
Asian/Pacific Islander	*	2	1%	1	1%	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	7	4%	4	5%	3	4%
Race/Sex							
White, not Hispanic Males	149.5	64	40%	29	37%	35	43%
White, not Hispanic Females	39.7	17	11%	11	14%	6	7%
Black/African American, not Hispanic Males	496.5	33	21%	14	18%	19	23%
Black/African American, not Hispanic Females	424.5	26	16%	13	17%	13	16%
Hispanic/Latino Males	489.6	8	5%	4	5%	4	5%
Hispanic/Latina Females	*	3	2%	2	3%	1	1%
Asian/Pacific Islander Males	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	7	4%	4	5%	3	4%
Total	152.3	160		78		82	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Allen

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	64	58%	26	51%	38	63%
Injection drug use (IDU)	8	7%	3	6%	5	8%
Male-to-male sexual contact and IDU	8	7%	3	6%	5	8%
Heterosexual contact	12	11%	5	10%	7	12%
Other/unknown	19	17%	14	27%	5	8%
Subtotal	111	100%	51	100%	60	100%
Female adult or adolescent						
Injection drug use	7	15%	3	11%	4	21%
Heterosexual contact	36	78%	22	81%	14	74%
Other/unknown	3	7%	2	7%	1	5%
Subtotal	46	100%	27	100%	19	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	100%	-	-	3	100%
Other/unknown	-	-	-	-	-	-
Subtotal	3	100%	-	-	3	100%
Total	160		78		82	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Allen

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	58	36%	25	32%	33	40%
Injection drug use (IDU) only	6	4%	3	4%	3	4%
Heterosexual contact only	48	30%	27	35%	21	26%
MSM & IDU	5	3%	1	1%	4	5%
IDU & Heterosexual contact	9	6%	3	4%	6	7%
MSM & Heterosexual contact	6	4%	1	1%	5	6%
MSM & IDU & Heterosexual contact	3	2%	2	3%	1	1%
Perinatal exposure	3	2%	-	-	3	4%
Other/unknown	22	14%	16	21%	6	7%
Total	160		78		82	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Allen

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	41	64%	17	53%	-	-	-	-	-	-	<10	*
Injection drug use (IDU)	2	3%	5	16%	1	14%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	5	8%	1	3%	2	29%	-	-	-	-	-	-
Heterosexual contact	4	6%	4	13%	3	43%	<10	*	-	-	-	-
Other/unknown	12	19%	5	16%	1	14%	<10	*	-	-	-	-
Subtotal	64	100%	32	100%	7	100%	<10	*	-	-	<10	*
Female adult or adolescent												
Injection drug use	3	19%	4	15%	-	-	-	-	-	-	-	-
Heterosexual contact	13	81%	19	73%	3	100%	-	-	-	-	<10	*
Other/unknown	-	-	3	12%	-	-	-	-	-	-	-	-
Subtotal	16	100%	26	100%	3	100%	-	-	-	-	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	1	100%	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	1	100%	-	-	-	-	-	-
Total	81		59		11		<10		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Allen

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	39	48%	14	24%	-	-	-	-	-	-	<10	*
Injection drug use (IDU) only	2	2%	4	7%	-	-	-	-	-	-	-	-
Heterosexual contact only	17	21%	23	39%	6	55%	<10	*	-	-	<10	*
MSM & IDU	2	2%	1	2%	2	18%	-	-	-	-	-	-
IDU & Heterosexual contact	3	4%	5	8%	1	9%	-	-	-	-	-	-
MSM & Heterosexual contact	2	2%	3	5%	-	-	-	-	-	-	<10	*
MSM & IDU & Heterosexual contact	3	4%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	1	2%	1	9%	-	-	-	-	-	-
Other/unknown	12	15%	8	14%	1	9%	<10	*	-	-	-	-
Total	81		59		11		<10		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Allen

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	91	71%	98	72%	102	71%	108	70%	113	71%
Females	38	29%	39	28%	41	29%	46	30%	47	29%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	1	1%	1	1%	-	-	-	-	-	-
15-19	-	-	-	-	1	1%	2	1%	2	1%
20-24	5	4%	8	6%	7	5%	6	4%	4	3%
25-29	10	8%	7	5%	8	6%	9	6%	13	8%
30-34	16	12%	17	12%	14	10%	16	10%	14	9%
35-39	17	13%	20	15%	23	16%	19	12%	19	12%
40-44	26	20%	26	19%	24	17%	26	17%	26	16%
45-49	21	16%	22	16%	22	15%	25	16%	24	15%
50-54	20	16%	19	14%	23	16%	21	14%	26	16%
55-64	10	8%	14	10%	16	11%	25	16%	25	16%
65+	3	2%	3	2%	5	3%	5	3%	7	4%
Race/Ethnicity										
White, not Hispanic	69	53%	72	53%	74	52%	78	51%	81	51%
Black/African American, not Hispanic	45	35%	49	36%	51	36%	56	36%	59	37%
Hispanic/Latino	10	8%	10	7%	11	8%	11	7%	11	7%
Asian/Pacific Islander	1	1%	2	1%	2	1%	2	1%	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	4	3%	4	3%	5	3%	7	5%	7	4%
Race/Sex										
White, not Hispanic Males	54	42%	57	42%	59	41%	61	40%	64	40%
White, not Hispanic Females	15	12%	15	11%	15	10%	17	11%	17	11%
Black/African American, not Hispanic Males	25	19%	28	20%	28	20%	31	20%	33	21%
Black/African American, not Hispanic Females	20	16%	21	15%	23	16%	25	16%	26	16%
Hispanic/Latino Males	7	5%	7	5%	8	6%	8	5%	8	5%
Hispanic/Latina Females	3	2%	3	2%	3	2%	3	2%	3	2%
Asian/Pacific Islander Males	1	1%	2	1%	2	1%	2	1%	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	4	3%	4	3%	5	3%	7	5%	7	4%
Total	129		137		143		154		160	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Allen

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	52	58%	55	57%	57	57%	60	57%	64	58%
Injection drug use (IDU)	7	8%	7	7%	7	7%	8	8%	8	7%
Male-to-male sexual contact and IDU	7	8%	8	8%	8	8%	8	8%	8	7%
Heterosexual contact	9	10%	10	10%	11	11%	12	11%	12	11%
Other/unknown	14	16%	16	17%	17	17%	18	17%	19	17%
Subtotal	89	100%	96	100%	100	100%	106	100%	111	100%
Female adult or adolescent										
Injection drug use	7	19%	7	18%	7	18%	7	16%	7	15%
Heterosexual contact	28	76%	29	76%	30	75%	35	78%	36	78%
Other/unknown	2	5%	2	5%	3	8%	3	7%	3	7%
Subtotal	37	100%	38	100%	40	100%	45	100%	46	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	3	100%	3	100%	3	100%	3	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	3	100%	3	100%	3	100%
Total	129		137		143		154		160	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Allen

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	47	36%	50	36%	52	36%	55	36%	58	36%
Injection drug use (IDU) only	6	5%	6	4%	6	4%	6	4%	6	4%
Heterosexual contact only	37	29%	39	28%	41	29%	47	31%	48	30%
MSM & IDU	5	4%	5	4%	5	3%	5	3%	5	3%
IDU & Heterosexual contact	8	6%	8	6%	8	6%	9	6%	9	6%
MSM & Heterosexual contact	5	4%	5	4%	5	3%	5	3%	6	4%
MSM & IDU & Heterosexual contact	2	2%	3	2%	3	2%	3	2%	3	2%
Perinatal exposure	3	2%	3	2%	3	2%	3	2%	3	2%
Other/unknown	16	12%	18	13%	20	14%	21	14%	22	14%
Total	129		137		143		154		160	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.