

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Ashtabula

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Ashtabula

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	130.3	65	66%	26	60%	39	70%
Females	69.0	34	34%	17	40%	17	30%
Age at end of year							
<13	*	1	1%	1	2%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	3	3%	3	7%	-	-
20-24	*	1	1%	1	2%	-	-
25-29	*	3	3%	3	7%	-	-
30-34	*	4	4%	3	7%	1	2%
35-39	128.5	7	7%	5	12%	2	4%
40-44	225.1	14	14%	5	12%	9	16%
45-49	217.1	15	15%	7	16%	8	14%
50-54	330.1	25	25%	6	14%	19	34%
55-64	145.3	21	21%	7	16%	14	25%
65+	29.0	5	5%	2	5%	3	5%
Race/Ethnicity							
White, not Hispanic	64.9	58	59%	26	60%	32	57%
Black/African American, not Hispanic	595.7	21	21%	9	21%	12	21%
Hispanic/Latino	472.9	18	18%	7	16%	11	20%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	2%	1	2%	1	2%
Race/Sex							
White, not Hispanic Males	92.9	41	41%	16	37%	25	45%
White, not Hispanic Females	37.6	17	17%	10	23%	7	13%
Black/African American, not Hispanic Males	503.4	12	12%	5	12%	7	13%
Black/African American, not Hispanic Females	788.8	9	9%	4	9%	5	9%
Hispanic/Latino Males	582.5	12	12%	5	12%	7	13%
Hispanic/Latina Females	343.6	6	6%	2	5%	4	7%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	2%	1	2%	1	2%
Total	99.8	99		43		56	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Ashtabula

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	36	56%	13	52%	23	59%
Injection drug use (IDU)	3	5%	1	4%	2	5%
Male-to-male sexual contact and IDU	8	13%	2	8%	6	15%
Heterosexual contact	10	16%	3	12%	7	18%
Other/unknown	7	11%	6	24%	1	3%
Subtotal	64	100%	25	100%	39	100%
Female adult or adolescent						
Injection drug use	1	3%	-	-	1	6%
Heterosexual contact	27	90%	11	85%	16	94%
Other/unknown	2	7%	2	15%	-	-
Subtotal	30	100%	13	100%	17	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	80%	4	80%	-	-
Other/unknown	1	20%	1	20%	-	-
Subtotal	5	100%	5	100%	-	-
Total	99		43		56	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Ashtabula

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	32	32%	13	30%	19	34%
Injection drug use (IDU) only	2	2%	-	-	2	4%
Heterosexual contact only	37	37%	14	33%	23	41%
MSM & IDU	6	6%	2	5%	4	7%
IDU & Heterosexual contact	2	2%	1	2%	1	2%
MSM & Heterosexual contact	4	4%	-	-	4	7%
MSM & IDU & Heterosexual contact	2	2%	-	-	2	4%
Perinatal exposure	4	4%	4	9%	-	-
Other/unknown	10	10%	9	21%	1	2%
Total	99		43		56	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Ashtabula

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	28	68%	7	58%	1	9%	-	-	-	-	-	-
Injection drug use (IDU)	2	5%	1	8%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	10%	2	17%	2	18%	-	-	-	-	-	-
Heterosexual contact	4	10%	-	-	6	55%	-	-	-	-	-	-
Other/unknown	3	7%	2	17%	2	18%	-	-	-	-	-	-
Subtotal	41	100%	12	100%	11	100%	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	-	-	1	14%	-	-	-	-	-	-	-	-
Heterosexual contact	15	94%	6	86%	4	80%	-	-	-	-	2	100%
Other/unknown	1	6%	-	-	1	20%	-	-	-	-	-	-
Subtotal	16	100%	7	100%	5	100%	-	-	-	-	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	2	100%	1	50%	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	50%	-	-	-	-	-	-
Subtotal	1	100%	2	100%	2	100%	-	-	-	-	-	-
Total	58		21		18		0		0		2	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Ashtabula

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	27	47%	4	19%	1	6%	-	-	-	-	-	-
Injection drug use (IDU) only	1	2%	1	5%	-	-	-	-	-	-	-	-
Heterosexual contact only	19	33%	6	29%	10	56%	-	-	-	-	2	100%
MSM & IDU	3	5%	1	5%	2	11%	-	-	-	-	-	-
IDU & Heterosexual contact	1	2%	1	5%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	2%	3	14%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	2%	1	5%	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	2	10%	1	6%	-	-	-	-	-	-
Other/unknown	4	7%	2	10%	4	22%	-	-	-	-	-	-
Total	58		21		18		0		0		2	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Ashtabula

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	55	63%	57	64%	58	64%	61	64%	65	66%
Females	32	37%	32	36%	33	36%	34	36%	34	34%
Age at end of year										
<13	1	1%	1	1%	1	1%	1	1%	1	1%
13-14	1	1%	1	1%	-	-	-	-	-	-
15-19	2	2%	2	2%	3	3%	3	3%	3	3%
20-24	1	1%	1	1%	1	1%	1	1%	1	1%
25-29	5	6%	4	4%	3	3%	3	3%	3	3%
30-34	3	3%	3	3%	2	2%	3	3%	4	4%
35-39	13	15%	12	13%	13	14%	11	12%	7	7%
40-44	8	9%	8	9%	8	9%	9	9%	14	14%
45-49	29	33%	26	29%	21	23%	21	22%	15	15%
50-54	7	8%	14	16%	18	20%	19	20%	25	25%
55-64	14	16%	14	16%	18	20%	20	21%	21	21%
65+	3	3%	3	3%	3	3%	4	4%	5	5%
Race/Ethnicity										
White, not Hispanic	51	59%	53	60%	54	59%	56	59%	58	59%
Black/African American, not Hispanic	18	21%	18	20%	19	21%	19	20%	21	21%
Hispanic/Latino	16	18%	16	18%	16	18%	18	19%	18	18%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	2	2%	2	2%	2	2%	2	2%
Race/Sex										
White, not Hispanic Males	35	40%	37	42%	38	42%	39	41%	41	41%
White, not Hispanic Females	16	18%	16	18%	16	18%	17	18%	17	17%
Black/African American, not Hispanic Males	10	11%	10	11%	10	11%	10	11%	12	12%
Black/African American, not Hispanic Females	8	9%	8	9%	9	10%	9	9%	9	9%
Hispanic/Latino Males	10	11%	10	11%	10	11%	12	13%	12	12%
Hispanic/Latina Females	6	7%	6	7%	6	7%	6	6%	6	6%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	2	2%	2	2%	2	2%	2	2%
Total	87		89		91		95		99	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Ashtabula

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	31	57%	33	59%	33	58%	33	55%	36	56%
Injection drug use (IDU)	3	6%	3	5%	3	5%	3	5%	3	5%
Male-to-male sexual contact and IDU	8	15%	8	14%	8	14%	8	13%	8	13%
Heterosexual contact	9	17%	9	16%	10	18%	10	17%	10	16%
Other/unknown	3	6%	3	5%	3	5%	6	10%	7	11%
Subtotal	54	100%	56	100%	57	100%	60	100%	64	100%
Female adult or adolescent										
Injection drug use	1	4%	1	4%	1	3%	1	3%	1	3%
Heterosexual contact	25	89%	25	89%	26	90%	27	90%	27	90%
Other/unknown	2	7%	2	7%	2	7%	2	7%	2	7%
Subtotal	28	100%	28	100%	29	100%	30	100%	30	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	4	80%	4	80%	4	80%
Other/unknown	1	20%	1	20%	1	20%	1	20%	1	20%
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	87		89		91		95		99	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Ashtabula

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	27	31%	29	33%	29	32%	29	31%	32	32%
Injection drug use (IDU) only	2	2%	2	2%	2	2%	2	2%	2	2%
Heterosexual contact only	34	39%	34	38%	36	40%	37	39%	37	37%
MSM & IDU	6	7%	6	7%	6	7%	6	6%	6	6%
IDU & Heterosexual contact	2	2%	2	2%	2	2%	2	2%	2	2%
MSM & Heterosexual contact	4	5%	4	4%	4	4%	4	4%	4	4%
MSM & IDU & Heterosexual contact	2	2%	2	2%	2	2%	2	2%	2	2%
Perinatal exposure	4	5%	4	4%	4	4%	4	4%	4	4%
Other/unknown	6	7%	6	7%	6	7%	9	9%	10	10%
Total	87		89		91		95		99	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.