

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Clark

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Clark**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	143.4	95	75%	43	78%	52	72%
Females	45.5	32	25%	12	22%	20	28%
<b>Age at end of year</b>							
<13	*	1	1%	-	-	1	1%
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	81.5	7	6%	5	9%	2	3%
25-29	124.2	10	8%	8	15%	2	3%
30-34	66.3	5	4%	3	5%	2	3%
35-39	161.9	12	9%	7	13%	5	7%
40-44	208.2	17	13%	8	15%	9	13%
45-49	214.9	19	15%	2	4%	17	24%
50-54	225.8	22	17%	6	11%	16	22%
55-64	135.9	26	20%	12	22%	14	19%
65+	32.7	8	6%	4	7%	4	6%
<b>Race/Ethnicity</b>							
White, not Hispanic	69.3	80	63%	32	58%	48	67%
Black/African American, not Hispanic	349.1	42	33%	19	35%	23	32%
Hispanic/Latino	*	2	2%	2	4%	-	-
Asian/Pacific Islander	*	1	1%	-	-	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	2%	2	4%	-	-
<b>Race/Sex</b>							
White, not Hispanic Males	116.5	65	51%	28	51%	37	51%
White, not Hispanic Females	25.1	15	12%	4	7%	11	15%
Black/African American, not Hispanic Males	444.4	26	20%	12	22%	14	19%
Black/African American, not Hispanic Females	258.9	16	13%	7	13%	9	13%
Hispanic/Latino Males	*	2	2%	2	4%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	-	-	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	2%	2	4%	-	-
<b>Total</b>	<b>93.0</b>	<b>127</b>		<b>55</b>		<b>72</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Clark**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	66	70%	30	70%	36	71%
Injection drug use (IDU)	3	3%	2	5%	1	2%
Male-to-male sexual contact and IDU	2	2%	1	2%	1	2%
Heterosexual contact	9	10%	3	7%	6	12%
Other/unknown	14	15%	7	16%	7	14%
Subtotal	94	100%	43	100%	51	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	9%	2	17%	1	5%
Heterosexual contact	23	72%	7	58%	16	80%
Other/unknown	6	19%	3	25%	3	15%
Subtotal	32	100%	12	100%	20	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	1	100%	-	-	1	100%
Subtotal	1	100%	-	-	1	100%
<b>Total</b>	<b>127</b>		<b>55</b>		<b>72</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Clark**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	65	51%	30	55%	35	49%
Injection drug use (IDU) only	2	2%	2	4%	-	-
Heterosexual contact only	32	25%	10	18%	22	31%
MSM & IDU	2	2%	1	2%	1	1%
IDU & Heterosexual contact	4	3%	2	4%	2	3%
MSM & Heterosexual contact	1	1%	-	-	1	1%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	21	17%	10	18%	11	15%
<b>Total</b>	<b>127</b>		<b>55</b>		<b>72</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Clark**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	49	75%	14	56%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	1	2%	2	8%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	2%	1	4%	-	-	-	-	-	-	-	-
Heterosexual contact	5	8%	4	16%	-	-	-	-	-	-	-	-
Other/unknown	9	14%	4	16%	-	-	<10	*	-	-	-	-
Subtotal	65	100%	25	100%	<10	*	<10	*	-	-	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	1	7%	2	13%	-	-	-	-	-	-	-	-
Heterosexual contact	12	80%	11	69%	-	-	-	-	-	-	-	-
Other/unknown	2	13%	3	19%	-	-	-	-	-	-	<10	*
Subtotal	15	100%	16	100%	-	-	-	-	-	-	<10	*
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
<b>Total</b>	<b>80</b>		<b>42</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Clark**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	48	60%	14	33%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	1	1%	1	2%	-	-	-	-	-	-	-	-
Heterosexual contact only	17	21%	15	36%	-	-	-	-	-	-	-	-
MSM & IDU	1	1%	1	2%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	1%	3	7%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	1%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	11	14%	8	19%	-	-	<10	*	-	-	<10	*
<b>Total</b>	<b>80</b>		<b>42</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Clark**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	83	76%	87	75%	90	76%	92	75%	95	75%
Females	26	24%	29	25%	29	24%	31	25%	32	25%
<b>Age at end of year</b>										
<13	1	1%	1	1%	1	1%	1	1%	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	1%	-	-	-	-	1	1%	-	-
20-24	7	6%	9	8%	9	8%	9	7%	7	6%
25-29	4	4%	5	4%	6	5%	7	6%	10	8%
30-34	9	8%	8	7%	9	8%	6	5%	5	4%
35-39	9	8%	11	9%	8	7%	9	7%	12	9%
40-44	19	17%	19	16%	18	15%	18	15%	17	13%
45-49	20	18%	21	18%	23	19%	19	15%	19	15%
50-54	16	15%	19	16%	20	17%	23	19%	22	17%
55-64	16	15%	15	13%	17	14%	22	18%	26	20%
65+	7	6%	8	7%	8	7%	8	7%	8	6%
<b>Race/Ethnicity</b>										
White, not Hispanic	71	65%	74	64%	75	63%	77	63%	80	63%
Black/African American, not Hispanic	34	31%	38	33%	39	33%	41	33%	42	33%
Hispanic/Latino	2	2%	2	2%	2	2%	2	2%	2	2%
Asian/Pacific Islander	-	-	-	-	1	1%	1	1%	1	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	2	2%	2	2%	2	2%	2	2%
<b>Race/Sex</b>										
White, not Hispanic Males	59	54%	61	53%	62	52%	63	51%	65	51%
White, not Hispanic Females	12	11%	13	11%	13	11%	14	11%	15	12%
Black/African American, not Hispanic Males	21	19%	23	20%	24	20%	25	20%	26	20%
Black/African American, not Hispanic Females	13	12%	15	13%	15	13%	16	13%	16	13%
Hispanic/Latino Males	2	2%	2	2%	2	2%	2	2%	2	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	1	1%	1	1%	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	2	2%	2	2%	2	2%	2	2%
<b>Total</b>	<b>109</b>		<b>116</b>		<b>119</b>		<b>123</b>		<b>127</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Clark

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	60	73%	62	72%	64	72%	66	73%	66	70%
Injection drug use (IDU)	3	4%	3	3%	3	3%	3	3%	3	3%
Male-to-male sexual contact and IDU	2	2%	2	2%	2	2%	2	2%	2	2%
Heterosexual contact	8	10%	9	10%	9	10%	9	10%	9	10%
Other/unknown	9	11%	10	12%	11	12%	11	12%	14	15%
Subtotal	82	100%	86	100%	89	100%	91	100%	94	100%
<b>Female adult or adolescent</b>										
Injection drug use	2	8%	2	7%	2	7%	2	6%	3	9%
Heterosexual contact	19	73%	21	72%	21	72%	23	74%	23	72%
Other/unknown	5	19%	6	21%	6	21%	6	19%	6	19%
Subtotal	26	100%	29	100%	29	100%	31	100%	32	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	1	100%	1	100%	1	100%
Subtotal	1	100%	1	100%	1	100%	1	100%	1	100%
<b>Total</b>	<b>109</b>		<b>116</b>		<b>119</b>		<b>123</b>		<b>127</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Clark

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	59	54%	61	53%	63	53%	65	53%	65	51%
Injection drug use (IDU) only	2	2%	2	2%	2	2%	2	2%	2	2%
Heterosexual contact only	27	25%	30	26%	30	25%	32	26%	32	25%
MSM & IDU	2	2%	2	2%	2	2%	2	2%	2	2%
IDU & Heterosexual contact	3	3%	3	3%	3	3%	3	2%	4	3%
MSM & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	15	14%	17	15%	18	15%	18	15%	21	17%
<b>Total</b>	<b>109</b>		<b>116</b>		<b>119</b>		<b>123</b>		<b>127</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.