

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Coshocton

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Coshocton

| Characteristic | Living with a diagnosis of HIV infection in 2014 | | | Current Disease Status | | | |
|--|--|-----------|-----|------------------------|---|-----------|------|
| | Rate ^a | No. | % | HIV (not AIDS) | | AIDS | |
| | | | | No. | % | No. | % |
| Sex | | | | | | | |
| Males | 88.5 | 16 | 76% | <10 | * | <10 | * |
| Females | 27.1 | 5 | 24% | <10 | * | <10 | * |
| Age at end of year | | | | | | | |
| <13 | * | - | - | - | - | - | - |
| 13-14 | * | - | - | - | - | - | - |
| 15-19 | * | - | - | - | - | - | - |
| 20-24 | * | 1 | 5% | <10 | * | - | - |
| 25-29 | * | - | - | - | - | - | - |
| 30-34 | * | 1 | 5% | <10 | * | - | - |
| 35-39 | * | 2 | 10% | <10 | * | <10 | * |
| 40-44 | * | 3 | 14% | <10 | * | <10 | * |
| 45-49 | 259.0 | 6 | 29% | <10 | * | <10 | * |
| 50-54 | * | 3 | 14% | <10 | * | <10 | * |
| 55-64 | * | 4 | 19% | <10 | * | <10 | * |
| 65+ | * | 1 | 5% | - | - | <10 | * |
| Race/Ethnicity | | | | | | | |
| White, not Hispanic | 54.1 | 19 | 90% | <10 | * | 13 | 100% |
| Black/African American, not Hispanic | * | 1 | 5% | <10 | * | - | - |
| Hispanic/Latino | * | 1 | 5% | <10 | * | - | - |
| Asian/Pacific Islander | * | - | - | - | - | - | - |
| American Indian/Alaska Native | * | - | - | - | - | - | - |
| Unknown | * | - | - | - | - | - | - |
| Race/Sex | | | | | | | |
| White, not Hispanic Males | 86.1 | 15 | 71% | <10 | * | <10 | * |
| White, not Hispanic Females | * | 4 | 19% | <10 | * | <10 | * |
| Black/African American, not Hispanic Males | * | 1 | 5% | <10 | * | - | - |
| Black/African American, not Hispanic Females | * | - | - | - | - | - | - |
| Hispanic/Latino Males | * | - | - | - | - | - | - |
| Hispanic/Latina Females | * | 1 | 5% | <10 | * | - | - |
| Asian/Pacific Islander Males | * | - | - | - | - | - | - |
| Asian/Pacific Islander Females | * | - | - | - | - | - | - |
| American Indian/Alaska Native Males | * | - | - | - | - | - | - |
| American Indian/Alaska Native Females | * | - | - | - | - | - | - |
| Unknown | * | - | - | - | - | - | - |
| Total | 57.5 | 21 | | <10 | | 13 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Coshocton

| Transmission Category ^a | Living with a diagnosis of HIV infection in 2014 | | Current Disease Status | | | |
|--|--|------|------------------------|---|-----------|---|
| | No. | % | HIV (not AIDS) | | AIDS | |
| | | | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 16 | 100% | <10 | * | <10 | * |
| Injection drug use (IDU) | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | - | - | - | - |
| Heterosexual contact | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - |
| Subtotal | 16 | 100% | <10 | * | <10 | * |
| Female adult or adolescent | | | | | | |
| Injection drug use | - | - | - | - | - | - |
| Heterosexual contact | 5 | 100% | <10 | * | <10 | * |
| Other/unknown | - | - | - | - | - | - |
| Subtotal | 5 | 100% | <10 | * | <10 | * |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - |
| Total | 21 | | <10 | | 13 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Coshocton

| Exposure Category ^a | Living with a diagnosis of HIV infection in 2014 | | Current Disease Status | | | |
|----------------------------------|--|-----|------------------------|---|-----------|---|
| | No. | % | HIV (not AIDS) | | AIDS | |
| | | | No. | % | No. | % |
| Male-to-male sex (MSM) only | 16 | 76% | <10 | * | <10 | * |
| Injection drug use (IDU) only | - | - | - | - | - | - |
| Heterosexual contact only | 5 | 24% | <10 | * | <10 | * |
| MSM & IDU | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - |
| Total | 21 | | <10 | | 13 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Coshocton

| Transmission Category ^a | Living with a Diagnosis of HIV Infection in 2014 | | | | | | | | | | | |
|--|--|------|------------------------|---|-----------------|---|------------------------|---|-------------------------------|---|---------------|---|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 15 | 100% | <10 | * | - | - | - | - | - | - | <10 | * |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 15 | 100% | <10 | * | - | - | - | - | - | - | <10 | * |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | 4 | 100% | - | - | <10 | * | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 4 | 100% | - | - | <10 | * | - | - | - | - | - | - |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - | - | - | - | - |
| Total | 19 | | <10 | | <10 | | 0 | | 0 | | <10 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Coshocton

| Exposure Category ^a | Living with a Diagnosis of HIV Infection in 2014 | | | | | | | | | | | |
|----------------------------------|--|-----|------------------------|---|-----------------|---|------------------------|---|-------------------------------|---|---------------|---|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 15 | 79% | <10 | * | - | - | - | - | - | - | <10 | * |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 4 | 21% | - | - | <10 | * | - | - | - | - | - | - |
| MSM & IDU | - | - | - | - | - | - | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Total | 19 | | <10 | | <10 | | 0 | | 0 | | <10 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Coshocton

Living with a Diagnosis of HIV Infection

| Characteristic | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|--|-----------|-----|-----------|-----|-----------|-----|-----------|-----|-----------|-----|
| | No. | % |
| Sex | | | | | | | | | | |
| Males | 14 | 82% | 14 | 78% | 15 | 79% | 16 | 80% | 16 | 76% |
| Females | 3 | 18% | 4 | 22% | 4 | 21% | 4 | 20% | 5 | 24% |
| Age at end of year | | | | | | | | | | |
| <13 | - | - | - | - | - | - | - | - | - | - |
| 13-14 | - | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | - | - | - | - | - | - | - | - |
| 20-24 | - | - | - | - | - | - | 1 | 5% | 1 | 5% |
| 25-29 | 1 | 6% | 1 | 6% | - | - | - | - | - | - |
| 30-34 | 2 | 12% | 1 | 6% | 1 | 5% | 1 | 5% | 1 | 5% |
| 35-39 | 2 | 12% | 3 | 17% | 2 | 11% | 2 | 10% | 2 | 10% |
| 40-44 | 6 | 35% | 4 | 22% | 5 | 26% | 4 | 20% | 3 | 14% |
| 45-49 | 3 | 18% | 5 | 28% | 5 | 26% | 5 | 25% | 6 | 29% |
| 50-54 | 2 | 12% | 3 | 17% | 3 | 16% | 4 | 20% | 3 | 14% |
| 55-64 | 1 | 6% | 1 | 6% | 3 | 16% | 3 | 15% | 4 | 19% |
| 65+ | - | - | - | - | - | - | - | - | 1 | 5% |
| Race/Ethnicity | | | | | | | | | | |
| White, not Hispanic | 16 | 94% | 17 | 94% | 18 | 95% | 18 | 90% | 19 | 90% |
| Black/African American, not Hispanic | - | - | - | - | - | - | 1 | 5% | 1 | 5% |
| Hispanic/Latino | 1 | 6% | 1 | 6% | 1 | 5% | 1 | 5% | 1 | 5% |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - |
| Race/Sex | | | | | | | | | | |
| White, not Hispanic Males | 14 | 82% | 14 | 78% | 15 | 79% | 15 | 75% | 15 | 71% |
| White, not Hispanic Females | 2 | 12% | 3 | 17% | 3 | 16% | 3 | 15% | 4 | 19% |
| Black/African American, not Hispanic Males | - | - | - | - | - | - | 1 | 5% | 1 | 5% |
| Black/African American, not Hispanic Females | - | - | - | - | - | - | - | - | - | - |
| Hispanic/Latino Males | - | - | - | - | - | - | - | - | - | - |
| Hispanic/Latina Females | 1 | 6% | 1 | 6% | 1 | 5% | 1 | 5% | 1 | 5% |
| Asian/Pacific Islander Males | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Females | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Males | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Females | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - |
| Total | 17 | | 18 | | 19 | | 20 | | 21 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Coshocton

Living with a Diagnosis of HIV Infection

| Transmission Category ^a | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|--|-----------|------|-----------|------|-----------|------|-----------|------|-----------|------|
| | No. | % |
| Male adult or adolescent | | | | | | | | | | |
| Male-to-male sexual contact | 14 | 100% | 14 | 100% | 15 | 100% | 16 | 100% | 16 | 100% |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 14 | 100% | 14 | 100% | 15 | 100% | 16 | 100% | 16 | 100% |
| Female adult or adolescent | | | | | | | | | | |
| Injection drug use | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | 3 | 100% | 4 | 100% | 4 | 100% | 4 | 100% | 5 | 100% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 3 | 100% | 4 | 100% | 4 | 100% | 4 | 100% | 5 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - | - | - |
| Total | 17 | | 18 | | 19 | | 20 | | 21 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Coshocton

Living with a Diagnosis of HIV Infection

| Exposure Category ^a | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|----------------------------------|-----------|-----|-----------|-----|-----------|-----|-----------|-----|-----------|-----|
| | No. | % |
| Male-to-male sex (MSM) only | 14 | 82% | 14 | 78% | 15 | 79% | 16 | 80% | 16 | 76% |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 3 | 18% | 4 | 22% | 4 | 21% | 4 | 20% | 5 | 24% |
| MSM & IDU | - | - | - | - | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Total | 17 | | 18 | | 19 | | 20 | | 21 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.