

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Crawford

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Crawford

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status				
	Rate ^a	No.	%	HIV (not AIDS)		AIDS		
				No.	%	No.	%	
Sex								
Males	120.9	25	76%	10	71%	15	79%	
Females	36.7	8	24%	4	29%	4	21%	
Age at end of year								
<13	*	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	
20-24	*	1	3%	1	7%	-	-	
25-29	*	3	9%	3	21%	-	-	
30-34	*	3	9%	1	7%	2	11%	
35-39	*	3	9%	3	21%	-	-	
40-44	187.0	5	15%	1	7%	4	21%	
45-49	*	3	9%	-	-	3	16%	
50-54	163.8	5	15%	-	-	5	26%	
55-64	114.9	7	21%	4	29%	3	16%	
65+	*	3	9%	1	7%	2	11%	
Race/Ethnicity								
White, not Hispanic	49.1	20	61%	5	36%	15	79%	
Black/African American, not Hispanic	2,919.7	12	36%	8	57%	4	21%	
Hispanic/Latino	*	1	3%	1	7%	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	
Unknown	*	-	-	-	-	-	-	
Race/Sex								
White, not Hispanic Males	65.9	13	39%	2	14%	11	58%	
White, not Hispanic Females	33.4	7	21%	3	21%	4	21%	
Black/African American, not Hispanic Males	4,280.2	11	33%	7	50%	4	21%	
Black/African American, not Hispanic Females	*	1	3%	1	7%	-	-	
Hispanic/Latino Males	*	1	3%	1	7%	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	
Unknown	*	-	-	-	-	-	-	
Total	77.7	33		14		19		

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Crawford

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	15	60%	6	60%	9	60%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	8%	-	-	2	13%
Heterosexual contact	4	16%	1	10%	3	20%
Other/unknown	4	16%	3	30%	1	7%
Subtotal	25	100%	10	100%	15	100%
Female adult or adolescent						
Injection drug use	1	13%	-	-	1	25%
Heterosexual contact	6	75%	4	100%	2	50%
Other/unknown	1	13%	-	-	1	25%
Subtotal	8	100%	4	100%	4	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	33		14		19	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Crawford

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	14	42%	5	36%	9	47%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	10	30%	5	36%	5	26%
MSM & IDU	2	6%	-	-	2	11%
IDU & Heterosexual contact	1	3%	-	-	1	5%
MSM & Heterosexual contact	1	3%	1	7%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	5	15%	3	21%	2	11%
Total	33		14		19	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Crawford

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	10	77%	4	36%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	15%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	8%	3	27%	-	-	-	-	-	-	-	-
Other/unknown	-	-	4	36%	-	-	-	-	-	-	-	-
Subtotal	13	100%	11	100%	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	1	14%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	5	71%	1	100%	-	-	-	-	-	-	-	-
Other/unknown	1	14%	-	-	-	-	-	-	-	-	-	-
Subtotal	7	100%	1	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	20		12		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Crawford

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	50%	4	33%	-	-	-	-	-	-	<10	*
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	6	30%	4	33%	-	-	-	-	-	-	-	-
MSM & IDU	2	10%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	<10	*	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	5%	4	33%	-	-	-	-	-	-	-	-
Total	20		12		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Crawford

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	22	79%	22	76%	23	77%	23	74%	25	76%
Females	6	21%	7	24%	7	23%	8	26%	8	24%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	1	3%	-	-	-	-
20-24	2	7%	2	7%	1	3%	2	6%	1	3%
25-29	2	7%	2	7%	2	7%	1	3%	3	9%
30-34	1	4%	1	3%	2	7%	3	10%	3	9%
35-39	5	18%	4	14%	4	13%	2	6%	3	9%
40-44	4	14%	5	17%	4	13%	6	19%	5	15%
45-49	4	14%	5	17%	5	17%	2	6%	3	9%
50-54	3	11%	2	7%	3	10%	6	19%	5	15%
55-64	6	21%	7	24%	6	20%	6	19%	7	21%
65+	1	4%	1	3%	2	7%	3	10%	3	9%
Race/Ethnicity										
White, not Hispanic	18	64%	19	66%	19	63%	20	65%	20	61%
Black/African American, not Hispanic	10	36%	10	34%	10	33%	10	32%	12	36%
Hispanic/Latino	-	-	-	-	1	3%	1	3%	1	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	-	-
Race/Sex										
White, not Hispanic Males	13	46%	13	45%	13	43%	13	42%	13	39%
White, not Hispanic Females	5	18%	6	21%	6	20%	7	23%	7	21%
Black/African American, not Hispanic Males	9	32%	9	31%	9	30%	9	29%	11	33%
Black/African American, not Hispanic Females	1	4%	1	3%	1	3%	1	3%	1	3%
Hispanic/Latino Males	-	-	-	-	1	3%	1	3%	1	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	-	-
Total	28		29		30		31		33	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Crawford

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	13	59%	13	59%	14	61%	14	61%	15	60%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	9%	2	9%	2	9%	2	9%	2	8%
Heterosexual contact	4	18%	4	18%	4	17%	4	17%	4	16%
Other/unknown	3	14%	3	14%	3	13%	3	13%	4	16%
Subtotal	22	100%	22	100%	23	100%	23	100%	25	100%
Female adult or adolescent										
Injection drug use	-	-	1	14%	1	14%	1	13%	1	13%
Heterosexual contact	6	100%	6	86%	6	86%	6	75%	6	75%
Other/unknown	-	-	-	-	-	-	1	13%	1	13%
Subtotal	6	100%	7	100%	7	100%	8	100%	8	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	28		29		30		31		33	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Crawford

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	13	46%	13	45%	13	43%	13	42%	14	42%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	10	36%	10	34%	10	33%	10	32%	10	30%
MSM & IDU	2	7%	2	7%	2	7%	2	6%	2	6%
IDU & Heterosexual contact	-	-	1	3%	1	3%	1	3%	1	3%
MSM & Heterosexual contact	-	-	-	-	1	3%	1	3%	1	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	11%	3	10%	3	10%	4	13%	5	15%
Total	28		29		30		31		33	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.