

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Cuyahoga

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Cuyahoga

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	31.0	186	89%	154	90%	21	81%	11	85%	
Females	3.6	24	11%	17	10%	5	19%	2	15%	
Age at diagnosis (yr)										
<13	*	1	<1%	1	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	18.7	15	7%	13	8%	1	4%	1	8%	
20-24	70.8	60	29%	56	33%	3	12%	1	8%	
25-29	40.2	34	16%	30	18%	2	8%	2	15%	
30-34	34.6	28	13%	22	13%	5	19%	1	8%	
35-39	32.2	23	11%	19	11%	3	12%	1	8%	
40-44	22.6	17	8%	11	6%	5	19%	1	8%	
45-49	16.0	13	6%	10	6%	1	4%	2	15%	
50-54	7.5	7	3%	4	2%	3	12%	-	-	
55-64	5.6	10	5%	5	3%	3	12%	2	15%	
65+	*	2	1%	-	-	-	-	2	15%	
Race/Ethnicity										
White, not Hispanic	4.5	34	16%	29	17%	4	15%	1	8%	
Black/African American, not Hispanic	41.7	155	74%	125	73%	19	73%	11	85%	
Hispanic/Latino	19.2	13	6%	9	5%	3	12%	1	8%	
Asian/Pacific Islander	*	2	1%	2	1%	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	6	3%	6	4%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	7.6	28	13%	24	14%	3	12%	1	8%	
White, not Hispanic Females	1.5	6	3%	5	3%	1	4%	-	-	
Black/African American, not Hispanic Males	81.8	138	66%	114	67%	15	58%	9	69%	
Black/African American, not Hispanic Females	8.4	17	8%	11	6%	4	15%	2	15%	
Hispanic/Latino Males	35.8	12	6%	8	5%	3	12%	1	8%	
Hispanic/Latina Females	*	1	<1%	1	1%	-	-	-	-	
Asian/Pacific Islander Males	*	2	1%	2	1%	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	6	3%	6	4%	-	-	-	-	
Total	16.7	210		171		26		13		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	122	66%	104	68%	12	57%	6	55%
Injection drug use (IDU)	1	1%	-	-	1	5%	-	-
Male-to-male sexual contact and IDU	2	1%	2	1%	-	-	-	-
Heterosexual contact	11	6%	9	6%	1	5%	1	9%
Other/unknown	49	26%	38	25%	7	33%	4	36%
Subtotal	185	100%	153	100%	21	100%	11	100%
Female adult or adolescent								
Injection drug use	3	13%	2	12%	1	20%	-	-
Heterosexual contact	16	67%	13	76%	3	60%	-	-
Other/unknown	5	21%	2	12%	1	20%	2	100%
Subtotal	24	100%	17	100%	5	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	210		171		26		13	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	121	58%	103	60%	12	46%	6	46%
Injection drug use (IDU) only	2	1%	1	1%	1	4%	-	-
Heterosexual contact only	27	13%	22	13%	4	15%	1	8%
MSM & IDU	2	1%	2	1%	-	-	-	-
IDU & Heterosexual contact	2	1%	1	1%	1	4%	-	-
MSM & Heterosexual contact	1	<1%	1	1%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	55	26%	41	24%	8	31%	6	46%
Total	210		171		26		13	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	16	57%	94	69%	9	75%	1	50%	-	-	2	33%
Injection drug use (IDU)	-	-	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	4%	-	-	1	8%	-	-	-	-	-	-
Heterosexual contact	2	7%	7	5%	-	-	-	-	-	-	2	33%
Other/unknown	9	32%	35	26%	2	17%	1	50%	-	-	2	33%
Subtotal	28	100%	137	100%	12	100%	2	100%	-	-	6	100%
Female adult or adolescent												
Injection drug use	3	50%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	33%	13	76%	1	100%	-	-	-	-	-	-
Other/unknown	1	17%	4	24%	-	-	-	-	-	-	-	-
Subtotal	6	100%	17	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	34		155		13		2		0		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	47%	93	60%	9	69%	1	50%	-	-	2	33%
Injection drug use (IDU) only	1	3%	1	1%	-	-	-	-	-	-	-	-
Heterosexual contact only	4	12%	20	13%	1	8%	-	-	-	-	2	33%
MSM & IDU	1	3%	-	-	1	8%	-	-	-	-	-	-
IDU & Heterosexual contact	2	6%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	1	1%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	10	29%	40	26%	2	15%	1	50%	-	-	2	33%
Total	34		155		13		2		0		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Cuyahoga

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	23.2	139	82%	26.2	157	79%	28.3	170	82%	31.5	189	83%
Females	4.5	30	18%	6.2	41	21%	5.8	38	18%	5.9	39	17%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	1	<1%
13-14	*	1	1%	*	0	<1%	*	-	-	*	1	<1%
15-19	23.7	19	11%	15.0	12	6%	16.2	13	6%	21.2	17	7%
20-24	38.9	33	20%	44.8	38	19%	48.3	41	20%	57.8	49	21%
25-29	24.8	21	12%	31.9	27	14%	36.7	31	15%	41.4	35	15%
30-34	28.4	23	14%	33.4	27	14%	39.5	32	15%	39.5	32	14%
35-39	23.8	17	10%	14.0	10	5%	29.4	21	10%	29.4	21	9%
40-44	18.6	14	8%	34.6	26	13%	22.6	17	8%	25.3	19	8%
45-49	19.7	16	9%	32.0	26	13%	22.2	18	9%	28.3	23	10%
50-54	12.8	12	7%	16.0	15	8%	21.3	20	10%	17.0	16	7%
55-64	7.3	13	8%	8.4	15	8%	6.7	12	6%	6.7	12	5%
65+	*	-	-	*	2	1%	*	3	1%	*	2	1%
Race/Ethnicity												
White, not Hispanic	5.4	41	24%	6.3	48	24%	6.1	46	22%	7.5	57	25%
Black/African American, not Hispanic	28.8	107	63%	35.0	130	66%	33.4	124	60%	37.1	138	61%
Hispanic/Latino	19.2	13	8%	19.2	13	7%	35.5	24	12%	31.0	21	9%
Asian/Pacific Islander	*	1	1%	*	1	1%	*	1	<1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	6	3%	*	13	6%	*	12	5%
Race/Sex												
White, not Hispanic Males	10.3	38	22%	11.2	41	21%	11.2	41	20%	13.6	50	22%
White, not Hispanic Females	*	3	2%	1.8	7	4%	1.3	5	2%	1.8	7	3%
Black/African American, not Hispanic Males	49.2	83	49%	59.3	100	51%	57.5	97	47%	67.0	113	50%
Black/African American, not Hispanic Females	11.8	24	14%	14.8	30	15%	13.3	27	13%	12.3	25	11%
Hispanic/Latino Males	29.8	10	6%	29.8	10	5%	62.7	21	10%	50.7	17	7%
Hispanic/Latina Females	*	3	2%	*	3	2%	*	3	1%	*	4	2%
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	<1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	6	3%	*	13	6%	*	12	5%
Total	13.4	169		15.7	198		16.5	208		18.1	228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Cuyahoga

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	96	70%	118	75%	116	68%	120	63%
Injection drug use (IDU)	1	1%	3	2%	4	2%	3	2%
Male-to-male sexual contact and IDU	1	1%	1	1%	10	6%	2	1%
Heterosexual contact	3	2%	5	3%	5	3%	9	5%
Other/unknown	37	27%	30	19%	35	21%	55	29%
Subtotal	138	100%	157	100%	170	100%	189	100%
Female adult or adolescent								
Injection drug use	4	13%	2	5%	2	5%	3	8%
Heterosexual contact	14	47%	23	56%	17	46%	20	54%
Other/unknown	12	40%	16	39%	18	49%	14	38%
Subtotal	30	100%	41	100%	37	100%	37	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	2	100%
Subtotal	1	100%	-	-	1	100%	2	100%
Total	169		198		208		228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Cuyahoga

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	96	57%	118	60%	113	54%	117	51%
Injection drug use (IDU) only	1	1%	3	2%	5	2%	5	2%
Heterosexual contact only	17	10%	28	14%	22	11%	29	13%
MSM & IDU	1	1%	1	1%	6	3%	2	1%
IDU & Heterosexual contact	4	2%	2	1%	1	<1%	1	<1%
MSM & Heterosexual contact	-	-	-	-	3	1%	3	1%
MSM & IDU & Heterosexual contact	-	-	-	-	4	2%	-	-
Perinatal exposure	1	1%	-	-	1	<1%	1	<1%
Other/unknown	49	29%	46	23%	53	25%	70	31%
Total	169		198		208		228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Cuyahoga

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	94	80%	114	83%	131	83%	137	83%	154	90%
Females	23	20%	24	17%	27	17%	29	17%	17	10%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	1	1%	1	1%
13-14	1	1%	-	-	-	-	1	1%	-	-
15-19	15	13%	11	8%	13	8%	17	10%	13	8%
20-24	24	21%	34	25%	34	22%	38	23%	56	33%
25-29	16	14%	20	14%	29	18%	25	15%	30	18%
30-34	15	13%	19	14%	25	16%	25	15%	22	13%
35-39	13	11%	8	6%	14	9%	15	9%	19	11%
40-44	8	7%	17	12%	11	7%	12	7%	11	6%
45-49	10	9%	8	6%	11	7%	15	9%	10	6%
50-54	6	5%	12	9%	11	7%	10	6%	4	2%
55-64	9	8%	8	6%	9	6%	6	4%	5	3%
65+	-	-	1	1%	1	1%	1	1%	-	-
Race/Ethnicity										
White, not Hispanic	26	22%	30	22%	37	23%	41	25%	29	17%
Black/African American, not Hispanic	78	67%	91	66%	94	59%	101	61%	125	73%
Hispanic/Latino	7	6%	12	9%	14	9%	13	8%	9	5%
Asian/Pacific Islander	1	1%	1	1%	1	1%	-	-	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	5	4%	4	3%	12	8%	11	7%	6	4%
Race/Sex										
White, not Hispanic Males	24	21%	26	19%	33	21%	38	23%	24	14%
White, not Hispanic Females	2	2%	4	3%	4	3%	3	2%	5	3%
Black/African American, not Hispanic Males	58	50%	75	54%	76	48%	81	49%	114	67%
Black/African American, not Hispanic Females	20	17%	16	12%	18	11%	20	12%	11	6%
Hispanic/Latino Males	6	5%	9	7%	11	7%	10	6%	8	5%
Hispanic/Latina Females	1	1%	3	2%	3	2%	3	2%	1	1%
Asian/Pacific Islander Males	1	1%	1	1%	1	1%	-	-	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	5	4%	4	3%	12	8%	11	7%	6	4%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	68	73%	87	76%	94	72%	87	64%	104	68%
Injection drug use (IDU)	-	-	3	3%	1	1%	2	1%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	7	5%	2	1%	2	1%
Heterosexual contact	3	3%	5	4%	3	2%	4	3%	9	6%
Other/unknown	22	24%	19	17%	26	20%	42	31%	38	25%
Subtotal	93	100%	114	100%	131	100%	137	100%	153	100%
Female adult or adolescent										
Injection drug use	2	9%	1	4%	1	4%	2	7%	2	12%
Heterosexual contact	13	57%	14	58%	13	50%	15	56%	13	76%
Other/unknown	8	35%	9	38%	12	46%	10	37%	2	12%
Subtotal	23	100%	24	100%	26	100%	27	100%	17	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	2	100%	1	100%
Subtotal	1	100%	-	-	1	100%	2	100%	1	100%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	68	58%	87	63%	91	58%	84	51%	103	60%
Injection drug use (IDU) only	-	-	3	2%	2	1%	4	2%	1	1%
Heterosexual contact only	16	14%	19	14%	16	10%	19	11%	22	13%
MSM & IDU	-	-	-	-	5	3%	2	1%	2	1%
IDU & Heterosexual contact	2	2%	1	1%	-	-	-	-	1	1%
MSM & Heterosexual contact	-	-	-	-	3	2%	3	2%	1	1%
MSM & IDU & Heterosexual contact	-	-	-	-	2	1%	-	-	-	-
Perinatal exposure	1	1%	-	-	1	1%	1	1%	-	-
Other/unknown	30	26%	28	20%	38	24%	53	32%	41	24%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Cuyahoga

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	67	74%	65	78%	54	72%	76	84%	43	83%	3,531	82%
Females	24	26%	18	22%	21	28%	15	16%	9	17%	771	18%
Age at AIDS diagnosis (yr)												
<13	2	2%	1	1%	1	1%	-	-	-	-	51	1%
13-14	1	1%	-	-	-	-	-	-	-	-	6	<1%
15-19	5	5%	4	5%	3	4%	2	2%	2	4%	91	2%
20-24	14	15%	10	12%	9	12%	13	14%	8	15%	363	8%
25-29	12	13%	10	12%	4	5%	18	20%	8	15%	719	17%
30-34	18	20%	10	12%	8	11%	12	13%	7	13%	878	20%
35-39	10	11%	5	6%	11	15%	10	11%	5	10%	827	19%
40-44	8	9%	11	13%	11	15%	9	10%	7	13%	557	13%
45-49	10	11%	21	25%	10	13%	10	11%	3	6%	373	9%
50-54	6	7%	4	5%	11	15%	7	8%	5	10%	203	5%
55-64	5	5%	6	7%	5	7%	8	9%	5	10%	183	4%
65+	-	-	1	1%	2	3%	2	2%	2	4%	51	1%
Race/Ethnicity												
White, not Hispanic	24	26%	20	24%	12	16%	24	26%	12	23%	1,545	36%
Black/African American, not Hispanic	59	65%	59	71%	52	69%	48	53%	34	65%	2,362	55%
Hispanic/Latino	7	8%	3	4%	10	13%	12	13%	4	8%	329	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	5	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	5	<1%
Unknown	1	1%	1	1%	1	1%	7	8%	2	4%	56	1%
Race/Sex												
White, not Hispanic Males	20	22%	18	22%	10	13%	18	20%	11	21%	1,425	33%
White, not Hispanic Females	4	4%	2	2%	2	3%	6	7%	1	2%	120	3%
Black/African American, not Hispanic Males	42	46%	43	52%	34	45%	42	46%	26	50%	1,816	42%
Black/African American, not Hispanic Females	17	19%	16	19%	18	24%	6	7%	8	15%	546	13%
Hispanic/Latino Males	4	4%	3	4%	9	12%	11	12%	4	8%	240	6%
Hispanic/Latina Females	3	3%	-	-	1	1%	1	1%	-	-	89	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	5	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	1	1%	1	1%	1	1%	7	8%	2	4%	56	1%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	44	66%	41	64%	32	59%	52	68%	26	60%	2,391	68%
Injection drug use (IDU)	2	3%	1	2%	3	6%	4	5%	1	2%	346	10%
Male-to-male sexual contact and IDU	3	4%	2	3%	4	7%	-	-	2	5%	268	8%
Heterosexual contact	2	3%	1	2%	2	4%	6	8%	4	9%	190	5%
Other/unknown	16	24%	19	30%	13	24%	14	18%	10	23%	316	9%
Subtotal	67	100%	64	100%	54	100%	76	100%	43	100%	3,511	100%
Female adult or adolescent												
Injection drug use	5	23%	1	6%	1	5%	2	13%	1	11%	182	25%
Heterosexual contact	9	41%	12	67%	10	50%	9	60%	3	33%	472	64%
Other/unknown	8	36%	5	28%	9	45%	4	27%	5	56%	86	12%
Subtotal	22	100%	18	100%	20	100%	15	100%	9	100%	740	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	1	100%	1	100%	-	-	-	-	46	90%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	10%
Subtotal	2	100%	1	100%	1	100%	-	-	-	-	51	100%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	41	45%	39	47%	32	43%	52	57%	26	50%	2,251	52%
Injection drug use (IDU) only	5	5%	1	1%	3	4%	3	3%	1	2%	318	7%
Heterosexual contact only	11	12%	13	16%	12	16%	15	16%	7	13%	662	15%
MSM & IDU	3	3%	2	2%	2	3%	-	-	2	4%	220	5%
IDU & Heterosexual contact	2	2%	1	1%	1	1%	3	3%	1	2%	210	5%
MSM & Heterosexual contact	3	3%	2	2%	-	-	-	-	-	-	138	3%
MSM & IDU & Heterosexual contact	-	-	-	-	2	3%	-	-	-	-	48	1%
Perinatal exposure	2	2%	1	1%	1	1%	-	-	-	-	46	1%
Other/unknown	24	26%	24	29%	22	29%	18	20%	15	29%	409	10%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Cuyahoga

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Cuyahoga

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	572.1	3,432	78%	1,766	78%	1,666	78%
Females	143.8	949	22%	489	22%	460	22%
Age at end of year							
<13	6.3	12	<1%	9	<1%	3	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	46.1	37	1%	32	1%	5	<1%
20-24	260.6	221	5%	178	8%	43	2%
25-29	350.1	296	7%	223	10%	73	3%
30-34	436.1	353	8%	241	11%	112	5%
35-39	478.6	342	8%	209	9%	133	6%
40-44	648.5	488	11%	259	11%	229	11%
45-49	851.5	691	16%	312	14%	379	18%
50-54	851.8	800	18%	334	15%	466	22%
55-64	511.4	911	21%	367	16%	544	26%
65+	110.6	229	5%	91	4%	138	6%
Race/Ethnicity							
White, not Hispanic	172.2	1,306	30%	655	29%	651	31%
Black/African American, not Hispanic	674.1	2,506	57%	1,304	58%	1,202	57%
Hispanic/Latino	572.0	387	9%	174	8%	213	10%
Asian/Pacific Islander	39.8	15	<1%	11	<1%	4	<1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	163	4%	109	5%	54	3%
Race/Sex							
White, not Hispanic Males	316.9	1,164	27%	579	26%	585	28%
White, not Hispanic Females	36.3	142	3%	76	3%	66	3%
Black/African American, not Hispanic Males	1,102.0	1,859	42%	968	43%	891	42%
Black/African American, not Hispanic Females	318.6	647	15%	336	15%	311	15%
Hispanic/Latino Males	790.6	265	6%	122	5%	143	7%
Hispanic/Latina Females	357.3	122	3%	52	2%	70	3%
Asian/Pacific Islander Males	64.9	12	<1%	9	<1%	3	<1%
Asian/Pacific Islander Females	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Males	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	163	4%	109	5%	54	3%
Total	347.7	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,317	68%	1,170	67%	1,147	69%
Injection drug use (IDU)	159	5%	54	3%	105	6%
Male-to-male sexual contact and IDU	163	5%	53	3%	110	7%
Heterosexual contact	231	7%	111	6%	120	7%
Other/unknown	537	16%	361	21%	176	11%
Subtotal	3,407	100%	1,749	100%	1,658	100%
Female adult or adolescent						
Injection drug use	124	13%	40	8%	84	19%
Heterosexual contact	593	64%	289	61%	304	68%
Other/unknown	203	22%	146	31%	57	13%
Subtotal	920	100%	475	100%	445	100%
Child (<13 yrs at diagnosis)						
Perinatal	44	81%	22	71%	22	96%
Other/unknown	10	19%	9	29%	1	4%
Subtotal	54	100%	31	100%	23	100%
Total	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,197	50%	1,128	50%	1,069	50%
Injection drug use (IDU) only	130	3%	50	2%	80	4%
Heterosexual contact only	824	19%	400	18%	424	20%
MSM & IDU	125	3%	48	2%	77	4%
IDU & Heterosexual contact	153	3%	44	2%	109	5%
MSM & Heterosexual contact	120	3%	42	2%	78	4%
MSM & IDU & Heterosexual contact	38	1%	5	<1%	33	2%
Perinatal exposure	45	1%	23	1%	22	1%
Other/unknown	749	17%	515	23%	234	11%
Total	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Cuyahoga

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	914	79%	1,194	65%	125	48%	6	50%	2	67%	76	59%
Injection drug use (IDU)	30	3%	71	4%	54	21%	-	-	-	-	4	3%
Male-to-male sexual contact and IDU	67	6%	71	4%	21	8%	-	-	-	-	4	3%
Heterosexual contact	33	3%	163	9%	23	9%	-	-	-	-	12	9%
Other/unknown	116	10%	344	19%	37	14%	6	50%	1	33%	33	26%
Subtotal	1,160	100%	1,843	100%	260	100%	12	100%	3	100%	129	100%
Female adult or adolescent												
Injection drug use	26	18%	68	11%	28	24%	-	-	-	-	2	6%
Heterosexual contact	86	61%	414	66%	72	61%	2	67%	1	100%	18	53%
Other/unknown	29	21%	141	23%	18	15%	1	33%	-	-	14	41%
Subtotal	141	100%	623	100%	118	100%	3	100%	1	100%	34	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	40%	33	83%	9	100%	-	-	-	-	-	-
Other/unknown	3	60%	7	18%	-	-	-	-	-	-	-	-
Subtotal	5	100%	40	100%	9	100%	-	-	-	-	-	-
Total	1,306		2,506		387		15		4		163	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Cuyahoga

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	895	69%	1106	44%	119	31%	5	33%	2	50%	70	43%
Injection drug use (IDU) only	29	2%	57	2%	41	11%	-	-	-	-	3	2%
Heterosexual contact only	119	9%	577	23%	95	25%	2	13%	1	25%	30	18%
MSM & IDU	60	5%	48	2%	14	4%	-	-	-	-	3	2%
IDU & Heterosexual contact	27	2%	82	3%	41	11%	-	-	-	-	3	2%
MSM & Heterosexual contact	19	1%	88	4%	6	2%	1	7%	-	-	6	4%
MSM & IDU & Heterosexual contact	7	1%	23	1%	7	2%	-	-	-	-	1	1%
Perinatal exposure	2	0%	34	1%	9	2%	-	-	-	-	-	-
Other/unknown	148	11%	491	20%	55	14%	7	47%	1	25%	47	29%
Total	1,306		2,506		387		15		4		163	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Cuyahoga

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	2,753	77%	2,901	77%	3,067	78%	3,252	78%	3,432	78%
Females	811	23%	848	23%	887	22%	927	22%	949	22%
Age at end of year										
<13	15	<1%	12	<1%	11	<1%	11	<1%	12	<1%
13-14	7	<1%	7	<1%	4	<1%	2	<1%	1	<1%
15-19	41	1%	34	1%	33	1%	36	1%	37	1%
20-24	146	4%	159	4%	182	5%	208	5%	221	5%
25-29	217	6%	230	6%	245	6%	272	7%	296	7%
30-34	254	7%	276	7%	314	8%	332	8%	353	8%
35-39	352	10%	321	9%	313	8%	328	8%	342	8%
40-44	596	17%	587	16%	552	14%	496	12%	488	11%
45-49	723	20%	732	20%	705	18%	726	17%	691	16%
50-54	579	16%	631	17%	716	18%	763	18%	800	18%
55-64	529	15%	632	17%	725	18%	815	20%	911	21%
65+	105	3%	128	3%	154	4%	190	5%	229	5%
Race/Ethnicity										
White, not Hispanic	1,138	32%	1,182	32%	1,220	31%	1,273	30%	1,306	30%
Black/African American, not Hispanic	1,974	55%	2,094	56%	2,219	56%	2,357	56%	2,506	57%
Hispanic/Latino	315	9%	328	9%	355	9%	376	9%	387	9%
Asian/Pacific Islander	12	<1%	12	<1%	13	<1%	13	<1%	15	<1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	121	3%	129	3%	143	4%	156	4%	163	4%
Race/Sex										
White, not Hispanic Males	1,020	29%	1,057	28%	1,090	28%	1,136	27%	1,164	27%
White, not Hispanic Females	118	3%	125	3%	130	3%	137	3%	142	3%
Black/African American, not Hispanic Males	1,423	40%	1,517	40%	1,615	41%	1,726	41%	1,859	42%
Black/African American, not Hispanic Females	551	15%	577	15%	604	15%	631	15%	647	15%
Hispanic/Latino Males	204	6%	214	6%	237	6%	255	6%	265	6%
Hispanic/Latina Females	111	3%	114	3%	118	3%	121	3%	122	3%
Asian/Pacific Islander Males	9	<1%	9	<1%	10	<1%	10	<1%	12	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	121	3%	129	3%	143	4%	156	4%	163	4%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Cuyahoga

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,848	68%	1,958	68%	2,076	68%	2,198	68%	2,317	68%
Injection drug use (IDU)	150	5%	152	5%	155	5%	158	5%	159	5%
Male-to-male sexual contact and IDU	148	5%	149	5%	158	5%	161	5%	163	5%
Heterosexual contact	201	7%	206	7%	211	7%	220	7%	231	7%
Other/unknown	382	14%	412	14%	443	15%	491	15%	537	16%
Subtotal	2,729	100%	2,877	100%	3,043	100%	3,228	100%	3,407	100%
Female adult or adolescent										
Injection drug use	114	15%	116	14%	118	14%	121	13%	124	13%
Heterosexual contact	521	66%	542	66%	558	65%	579	64%	593	64%
Other/unknown	149	19%	163	20%	184	21%	198	22%	203	22%
Subtotal	784	100%	821	100%	860	100%	898	100%	920	100%
Child (<13 yrs at diagnosis)										
Perinatal	44	86%	44	86%	44	86%	44	83%	44	81%
Other/unknown	7	14%	7	14%	7	14%	9	17%	10	19%
Subtotal	51	100%	51	100%	51	100%	53	100%	54	100%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Cuyahoga

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	1,737	49%	1,846	49%	1,961	50%	2,079	50%	2,197	50%
Injection drug use (IDU) only	117	3%	119	3%	123	3%	128	3%	130	3%
Heterosexual contact only	722	20%	748	20%	769	19%	799	19%	824	19%
MSM & IDU	113	3%	114	3%	120	3%	123	3%	125	3%
IDU & Heterosexual contact	147	4%	149	4%	150	4%	151	4%	153	3%
MSM & Heterosexual contact	111	3%	112	3%	115	3%	119	3%	120	3%
MSM & IDU & Heterosexual contact	35	1%	35	1%	38	1%	38	1%	38	1%
Perinatal exposure	44	1%	44	1%	44	1%	45	1%	45	1%
Other/unknown	538	15%	582	16%	634	16%	697	17%	749	17%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.