

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Delaware

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Delaware

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	132.3	124	86%	71	87%	53	84%
Females	22.0	21	14%	11	13%	10	16%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	63.8	6	4%	5	6%	1	2%
25-29	88.8	7	5%	5	6%	2	3%
30-34	69.4	8	6%	6	7%	2	3%
35-39	108.4	15	10%	8	10%	7	11%
40-44	136.1	22	15%	9	11%	13	21%
45-49	134.9	20	14%	14	17%	6	10%
50-54	217.8	31	21%	14	17%	17	27%
55-64	119.6	27	19%	16	20%	11	17%
65+	41.3	9	6%	5	6%	4	6%
Race/Ethnicity							
White, not Hispanic	70.0	115	79%	64	78%	51	81%
Black/African American, not Hispanic	340.5	22	15%	13	16%	9	14%
Hispanic/Latino	*	3	2%	3	4%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	5	3%	2	2%	3	5%
Race/Sex							
White, not Hispanic Males	133.0	108	74%	62	76%	46	73%
White, not Hispanic Females	8.4	7	5%	2	2%	5	8%
Black/African American, not Hispanic Males	330.2	11	8%	6	7%	5	8%
Black/African American, not Hispanic Females	351.4	11	8%	7	9%	4	6%
Hispanic/Latino Males	*	1	1%	1	1%	-	-
Hispanic/Latina Females	*	2	1%	2	2%	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	5	3%	2	2%	3	5%
Total	76.7	145		82		63	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Delaware

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	85	69%	48	68%	37	70%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	6	5%	-	-	6	11%
Heterosexual contact	3	2%	3	4%	-	-
Other/unknown	30	24%	20	28%	10	19%
Subtotal	124	100%	71	100%	53	100%
Female adult or adolescent						
Injection drug use	1	5%	-	-	1	10%
Heterosexual contact	18	86%	10	91%	8	80%
Other/unknown	2	10%	1	9%	1	10%
Subtotal	21	100%	11	100%	10	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	145		82		63	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Delaware

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	81	56%	46	56%	35	56%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	21	14%	13	16%	8	13%
MSM & IDU	6	4%	-	-	6	10%
IDU & Heterosexual contact	1	1%	-	-	1	2%
MSM & Heterosexual contact	4	3%	2	2%	2	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	32	22%	21	26%	11	17%
Total	145		82		63	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Delaware

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	79	73%	4	36%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	4%	1	9%	-	-	-	-	-	-	<10	*
Heterosexual contact	1	1%	2	18%	-	-	-	-	-	-	-	-
Other/unknown	24	22%	4	36%	-	-	-	-	-	-	<10	*
Subtotal	108	100%	11	100%	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	1	14%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	5	71%	10	91%	<10	*	-	-	-	-	<10	*
Other/unknown	1	14%	1	9%	-	-	-	-	-	-	-	-
Subtotal	7	100%	11	100%	<10	*	-	-	-	-	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	115		22		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Delaware

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	76	66%	3	14%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	6	5%	12	55%	<10	*	-	-	-	-	<10	*
MSM & IDU	4	3%	1	5%	-	-	-	-	-	-	<10	*
IDU & Heterosexual contact	1	1%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	3	3%	1	5%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	25	22%	5	23%	-	-	-	-	-	-	<10	*
Total	115		22		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Delaware

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	103	86%	109	86%	115	86%	120	86%	124	86%
Females	17	14%	18	14%	19	14%	19	14%	21	14%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	1%	-	-	1	1%	1	1%	-	-
20-24	2	2%	3	2%	6	4%	6	4%	6	4%
25-29	2	2%	2	2%	3	2%	5	4%	7	5%
30-34	11	9%	13	10%	12	9%	8	6%	8	6%
35-39	19	16%	14	11%	13	10%	17	12%	15	10%
40-44	17	14%	22	17%	22	16%	20	14%	22	15%
45-49	29	24%	30	24%	27	20%	27	19%	20	14%
50-54	9	8%	11	9%	17	13%	20	14%	31	21%
55-64	26	22%	27	21%	27	20%	29	21%	27	19%
65+	4	3%	5	4%	6	4%	6	4%	9	6%
Race/Ethnicity										
White, not Hispanic	98	82%	102	80%	106	79%	110	79%	115	79%
Black/African American, not Hispanic	17	14%	19	15%	21	16%	21	15%	22	15%
Hispanic/Latino	2	2%	3	2%	3	2%	3	2%	3	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	3	2%	4	3%	5	4%	5	3%
Race/Sex										
White, not Hispanic Males	92	77%	96	76%	100	75%	104	75%	108	74%
White, not Hispanic Females	6	5%	6	5%	6	4%	6	4%	7	5%
Black/African American, not Hispanic Males	9	8%	10	8%	11	8%	11	8%	11	8%
Black/African American, not Hispanic Females	8	7%	9	7%	10	7%	10	7%	11	8%
Hispanic/Latino Males	-	-	1	1%	1	1%	1	1%	1	1%
Hispanic/Latina Females	2	2%	2	2%	2	1%	2	1%	2	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	3	2%	4	3%	5	4%	5	3%
Total	120		127		134		139		145	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Delaware

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	69	67%	74	68%	80	70%	82	68%	85	69%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	6	6%	6	6%	6	5%	6	5%	6	5%
Heterosexual contact	3	3%	3	3%	3	3%	3	3%	3	2%
Other/unknown	25	24%	26	24%	26	23%	29	24%	30	24%
Subtotal	103	100%	109	100%	115	100%	120	100%	124	100%
Female adult or adolescent										
Injection drug use	1	6%	1	6%	1	5%	1	5%	1	5%
Heterosexual contact	14	82%	15	83%	16	84%	16	84%	18	86%
Other/unknown	2	12%	2	11%	2	11%	2	11%	2	10%
Subtotal	17	100%	18	100%	19	100%	19	100%	21	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	120		127		134		139		145	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Delaware

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	67	56%	72	57%	77	57%	78	56%	81	56%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	17	14%	18	14%	19	14%	19	14%	21	14%
MSM & IDU	6	5%	6	5%	6	4%	6	4%	6	4%
IDU & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
MSM & Heterosexual contact	2	2%	2	2%	3	2%	4	3%	4	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	27	23%	28	22%	28	21%	31	22%	32	22%
Total	120		127		134		139		145	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.