

Diagnoses of HIV and/or AIDS  
Reported in HIV Prevention Planning  
Franklin

**Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Franklin**

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
<b>Sex</b>										
Males	31.5	189	88%	128	88%	44	90%	17	85%	
Females	4.1	26	12%	18	12%	5	10%	3	15%	
<b>Age at diagnosis (yr)</b>										
<13	*	1	<1%	1	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	7.6	6	3%	6	4%	-	-	-	-	
20-24	50.7	47	22%	33	23%	10	20%	4	20%	
25-29	38.8	46	21%	35	24%	7	14%	4	20%	
30-34	37.2	37	17%	22	15%	7	14%	8	40%	
35-39	29.7	25	12%	20	14%	5	10%	-	-	
40-44	21.2	17	8%	10	7%	7	14%	-	-	
45-49	27.4	21	10%	12	8%	7	14%	2	10%	
50-54	7.4	6	3%	2	1%	3	6%	1	5%	
55-64	5.0	7	3%	4	3%	2	4%	1	5%	
65+	*	2	1%	1	1%	1	2%	-	-	
<b>Race/Ethnicity</b>										
White, not Hispanic	10.5	85	40%	58	40%	23	47%	4	20%	
Black/African American, not Hispanic	38.6	103	48%	70	48%	20	41%	13	65%	
Hispanic/Latino	25.4	16	7%	11	8%	3	6%	2	10%	
Asian/Pacific Islander	*	4	2%	2	1%	1	2%	1	5%	
American Indian/Alaska Native	*	1	<1%	1	1%	-	-	-	-	
Unknown	*	6	3%	4	3%	2	4%	-	-	
<b>Race/Sex</b>										
White, not Hispanic Males	20.4	81	38%	56	38%	22	45%	3	15%	
White, not Hispanic Females	*	4	2%	2	1%	1	2%	1	5%	
Black/African American, not Hispanic Males	68.3	86	40%	59	40%	16	33%	11	55%	
Black/African American, not Hispanic Females	12.0	17	8%	11	8%	4	8%	2	10%	
Hispanic/Latino Males	42.5	14	7%	9	6%	3	6%	2	10%	
Hispanic/Latina Females	*	2	1%	2	1%	-	-	-	-	
Asian/Pacific Islander Males	*	3	1%	1	1%	1	2%	1	5%	
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	1	<1%	1	1%	-	-	-	-	
Unknown	*	6	3%	4	3%	2	4%	-	-	
<b>Total</b>	<b>17.5</b>	<b>215</b>		<b>146</b>		<b>49</b>		<b>20</b>		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Franklin

Transmission Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	144	76%	101	79%	33	75%	10	59%
Injection drug use (IDU)	1	1%	1	1%	-	-	-	-
Male-to-male sexual contact and IDU	6	3%	6	5%	-	-	-	-
Heterosexual contact	5	3%	-	-	2	5%	3	18%
Other/unknown	33	17%	20	16%	9	20%	4	24%
Subtotal	189	100%	128	100%	44	100%	17	100%
<b>Female adult or adolescent</b>								
Injection drug use	1	4%	1	6%	-	-	-	-
Heterosexual contact	18	72%	13	76%	2	40%	3	100%
Other/unknown	6	24%	3	18%	3	60%	-	-
Subtotal	25	100%	17	100%	5	100%	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
<b>Total</b>	<b>215</b>		<b>146</b>		<b>49</b>		<b>20</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Franklin

Exposure Category <sup>a</sup>	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	138	64%	95	65%	33	67%	10	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	23	11%	13	9%	4	8%	6	30%
MSM & IDU	3	1%	3	2%	-	-	-	-
IDU & Heterosexual contact	2	1%	2	1%	-	-	-	-
MSM & Heterosexual contact	6	3%	6	4%	-	-	-	-
MSM & IDU & Heterosexual contact	3	1%	3	2%	-	-	-	-
Perinatal exposure	1	<1%	1	1%	-	-	-	-
Other/unknown	39	18%	23	16%	12	24%	4	20%
<b>Total</b>	<b>215</b>		<b>146</b>		<b>49</b>		<b>20</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Franklin

Transmission Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	63	78%	67	78%	8	57%	2	67%	-	-	4	80%
Injection drug use (IDU)	1	1%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	3	4%	1	1%	1	7%	-	-	-	-	1	20%
Heterosexual contact	1	1%	3	3%	1	7%	-	-	-	-	-	-
Other/unknown	13	16%	15	17%	4	29%	1	33%	-	-	-	-
Subtotal	81	100%	86	100%	14	100%	3	100%	-	-	5	100%
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	1	50%	-	-	-	-	-	-
Heterosexual contact	3	75%	12	71%	1	50%	-	-	1	100%	1	100%
Other/unknown	1	25%	5	29%	-	-	-	-	-	-	-	-
Subtotal	4	100%	17	100%	2	100%	-	-	1	100%	1	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	-	-	-	-
<b>Total</b>	<b>85</b>		<b>103</b>		<b>16</b>		<b>4</b>		<b>1</b>		<b>6</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Franklin

Exposure Category <sup>a</sup>	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	61	72%	64	62%	8	50%	1	25%	-	-	4	67%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	4	5%	15	15%	2	13%	-	-	1	100%	1	17%
MSM & IDU	1	1%	1	1%	1	6%	-	-	-	-	-	-
IDU & Heterosexual contact	1	1%	-	-	1	6%	-	-	-	-	-	-
MSM & Heterosexual contact	2	2%	3	3%	-	-	1	25%	-	-	-	-
MSM & IDU & Heterosexual contact	2	2%	-	-	-	-	-	-	-	-	1	17%
Perinatal exposure	-	-	-	-	-	-	1	25%	-	-	-	-
Other/unknown	14	16%	20	19%	4	25%	1	25%	-	-	-	-
<b>Total</b>	<b>85</b>		<b>103</b>		<b>16</b>		<b>4</b>		<b>1</b>		<b>6</b>	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Franklin

### Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate <sup>a</sup>	No.	%									
<b>Sex</b>												
Males	36.2	217	85%	35.3	212	81%	34.2	205	83%	30.8	185	81%
Females	5.9	37	15%	7.8	49	19%	6.5	41	17%	6.7	42	19%
<b>Age at diagnosis (yr)</b>												
<13	*	-	-	*	1	<1%	*	1	<1%	*	1	<1%
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	15.3	12	5%	20.3	16	6%	19.1	15	6%	19.1	15	7%
20-24	56.1	52	20%	53.9	50	19%	56.1	52	21%	60.4	56	25%
25-29	32.9	39	15%	35.5	42	16%	37.1	44	18%	27.0	32	14%
30-34	36.2	36	14%	42.2	42	16%	34.1	34	14%	22.1	22	10%
35-39	41.6	35	14%	42.8	36	14%	28.5	24	10%	22.6	19	8%
40-44	38.6	31	12%	34.9	28	11%	31.2	25	10%	31.2	25	11%
45-49	31.4	24	9%	22.2	17	7%	27.4	21	9%	28.7	22	10%
50-54	12.4	10	4%	18.5	15	6%	14.8	12	5%	24.7	20	9%
55-64	10.1	14	6%	7.9	11	4%	11.5	16	7%	9.3	13	6%
65+	*	1	<1%	*	3	1%	*	2	1%	*	2	1%
<b>Race/Ethnicity</b>												
White, not Hispanic	14.9	121	48%	14.2	115	44%	12.6	102	41%	11.5	93	41%
Black/African American, not Hispanic	40.8	109	43%	41.5	111	43%	43.4	116	47%	38.2	102	45%
Hispanic/Latino	20.7	13	5%	33.4	21	8%	12.7	8	3%	20.7	13	6%
Asian/Pacific Islander	*	2	1%	*	2	1%	*	4	2%	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	9	4%	*	12	5%	*	16	7%	*	17	7%
<b>Race/Sex</b>												
White, not Hispanic Males	27.8	110	43%	25.7	102	39%	24.5	97	39%	21.2	84	37%
White, not Hispanic Females	2.7	11	4%	3.1	13	5%	1.2	5	2%	2.2	9	4%
Black/African American, not Hispanic Males	68.3	86	34%	63.5	80	31%	67.5	85	35%	61.1	77	34%
Black/African American, not Hispanic Females	16.3	23	9%	22.0	31	12%	22.0	31	13%	17.7	25	11%
Hispanic/Latino Males	36.4	12	5%	51.6	17	7%	24.3	8	3%	27.3	9	4%
Hispanic/Latina Females	*	1	<1%	*	4	2%	*	-	-	*	4	2%
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	3	1%	*	1	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	9	4%	*	12	5%	*	16	7%	*	17	7%
<b>Total</b>	<b>20.6</b>	<b>254</b>		<b>21.2</b>	<b>261</b>		<b>20.0</b>	<b>246</b>		<b>18.4</b>	<b>227</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Franklin

### Diagnosis of HIV Infection by Year

Transmission Category <sup>a</sup>	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	156	72%	150	71%	147	72%	132	71%
Injection drug use (IDU)	-	-	2	1%	1	<1%	-	-
Male-to-male sexual contact and IDU	8	4%	6	3%	2	1%	7	4%
Heterosexual contact	9	4%	3	1%	4	2%	7	4%
Other/unknown	44	20%	51	24%	50	25%	39	21%
Subtotal	217	100%	212	100%	204	100%	185	100%
<b>Female adult or adolescent</b>								
Injection drug use	3	8%	-	-	1	2%	5	12%
Heterosexual contact	27	73%	31	65%	23	56%	22	54%
Other/unknown	7	19%	17	35%	17	41%	14	34%
Subtotal	37	100%	48	100%	41	100%	41	100%
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	-	-	1	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	1	100%	1	100%
<b>Total</b>	<b>254</b>		<b>261</b>		<b>246</b>		<b>227</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Franklin

### Diagnosis of HIV Infection by Year

Exposure Category <sup>a</sup>	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	152	60%	148	57%	138	56%	123	54%
Injection drug use (IDU) only	-	-	2	1%	2	1%	1	<1%
Heterosexual contact only	36	14%	34	13%	27	11%	29	13%
MSM & IDU	8	3%	6	2%	2	1%	3	1%
IDU & Heterosexual contact	3	1%	-	-	-	-	4	2%
MSM & Heterosexual contact	4	2%	2	1%	9	4%	9	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	4	2%
Perinatal exposure	-	-	1	<1%	1	<1%	1	<1%
Other/unknown	51	20%	68	26%	67	27%	53	23%
<b>Total</b>	<b>254</b>		<b>261</b>		<b>246</b>		<b>227</b>	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Franklin**

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Sex</b>										
Males	132	87%	132	81%	128	81%	128	83%	128	88%
Females	20	13%	31	19%	30	19%	27	17%	18	12%
<b>Age at HIV diagnosis (yr)</b>										
<13	-	-	-	-	-	-	1	1%	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	9	6%	13	8%	13	8%	13	8%	6	4%
20-24	38	25%	36	22%	42	27%	43	28%	33	23%
25-29	30	20%	30	18%	26	16%	26	17%	35	24%
30-34	15	10%	28	17%	21	13%	16	10%	22	15%
35-39	21	14%	12	7%	13	8%	13	8%	20	14%
40-44	15	10%	17	10%	14	9%	15	10%	10	7%
45-49	11	7%	9	6%	14	9%	13	8%	12	8%
50-54	6	4%	10	6%	6	4%	6	4%	2	1%
55-64	6	4%	5	3%	9	6%	8	5%	4	3%
65+	1	1%	3	2%	-	-	1	1%	1	1%
<b>Race/Ethnicity</b>										
White, not Hispanic	74	49%	80	49%	68	43%	65	42%	58	40%
Black/African American, not Hispanic	64	42%	66	40%	77	49%	67	43%	70	48%
Hispanic/Latino	8	5%	10	6%	3	2%	10	6%	11	8%
Asian/Pacific Islander	2	1%	1	1%	4	3%	1	1%	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	1%
Unknown	4	3%	6	4%	6	4%	12	8%	4	3%
<b>Race/Sex</b>										
White, not Hispanic Males	67	44%	72	44%	64	41%	56	36%	56	38%
White, not Hispanic Females	7	5%	8	5%	4	3%	9	6%	2	1%
Black/African American, not Hispanic Males	53	35%	45	28%	54	34%	54	35%	59	40%
Black/African American, not Hispanic Females	11	7%	21	13%	23	15%	13	8%	11	8%
Hispanic/Latino Males	7	5%	9	6%	3	2%	7	5%	9	6%
Hispanic/Latina Females	1	1%	1	1%	-	-	3	2%	2	1%
Asian/Pacific Islander Males	1	1%	-	-	3	2%	1	1%	1	1%
Asian/Pacific Islander Females	1	1%	1	1%	1	1%	-	-	1	1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	1%
Unknown	4	3%	6	4%	6	4%	12	8%	4	3%
<b>Total</b>	<b>152</b>		<b>163</b>		<b>158</b>		<b>155</b>		<b>146</b>	

*Note:* Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Franklin**

Transmission Category <sup>a</sup>	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	99	75%	93	70%	98	77%	96	75%	101	79%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	1%
Male-to-male sexual contact and IDU	5	4%	4	3%	2	2%	7	5%	6	5%
Heterosexual contact	5	4%	1	1%	2	2%	3	2%	-	-
Other/unknown	23	17%	34	26%	26	20%	22	17%	20	16%
Subtotal	132	100%	132	100%	128	100%	128	100%	128	100%
<b>Female adult or adolescent</b>										
Injection drug use	1	5%	-	-	1	3%	5	19%	1	6%
Heterosexual contact	14	70%	22	71%	15	50%	15	58%	13	76%
Other/unknown	5	25%	9	29%	14	47%	6	23%	3	18%
Subtotal	20	100%	31	100%	30	100%	26	100%	17	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	1	100%
<b>Total</b>	<b>152</b>		<b>163</b>		<b>158</b>		<b>155</b>		<b>146</b>	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Franklin**

Exposure Category <sup>a</sup>	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	98	64%	91	56%	92	58%	89	57%	95	65%
Injection drug use (IDU) only	-	-	-	-	1	1%	1	1%	-	-
Heterosexual contact only	19	13%	23	14%	17	11%	18	12%	13	9%
MSM & IDU	5	3%	4	2%	2	1%	3	2%	3	2%
IDU & Heterosexual contact	1	1%	-	-	-	-	4	3%	2	1%
MSM & Heterosexual contact	1	1%	2	1%	6	4%	7	5%	6	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	4	3%	3	2%
Perinatal exposure	-	-	-	-	-	-	1	1%	1	1%
Other/unknown	28	18%	43	26%	40	25%	28	18%	23	16%
<b>Total</b>	<b>152</b>		<b>163</b>		<b>158</b>		<b>155</b>		<b>146</b>	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Franklin**

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Sex</b>												
Males	125	86%	117	80%	107	82%	109	83%	87	84%	3,279	87%
Females	21	14%	30	20%	23	18%	23	17%	16	16%	499	13%
<b>Age at AIDS diagnosis (yr)</b>												
<13	-	-	1	1%	-	-	-	-	-	-	23	1%
13-14	-	-	-	-	-	-	-	-	-	-	5	<1%
15-19	9	6%	8	5%	4	3%	3	2%	4	4%	75	2%
20-24	19	13%	18	12%	20	15%	32	24%	18	17%	379	10%
25-29	16	11%	16	11%	25	19%	16	12%	18	17%	644	17%
30-34	30	21%	24	16%	19	15%	16	12%	24	23%	851	23%
35-39	22	15%	34	23%	19	15%	18	14%	7	7%	664	18%
40-44	23	16%	21	14%	15	12%	13	10%	11	11%	486	13%
45-49	14	10%	13	9%	12	9%	12	9%	9	9%	327	9%
50-54	6	4%	6	4%	6	5%	13	10%	6	6%	178	5%
55-64	6	4%	6	4%	9	7%	8	6%	4	4%	123	3%
65+	1	1%	-	-	1	1%	1	1%	2	2%	23	1%
<b>Race/Ethnicity</b>												
White, not Hispanic	69	47%	62	42%	63	48%	56	42%	40	39%	2,287	61%
Black/African American, not Hispanic	69	47%	64	44%	47	36%	62	47%	50	49%	1,260	33%
Hispanic/Latino	4	3%	10	7%	8	6%	4	3%	6	6%	118	3%
Asian/Pacific Islander	-	-	1	1%	-	-	-	-	3	3%	16	<1%
American Indian/Alaska Native	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	4	3%	9	6%	12	9%	10	8%	4	4%	95	3%
<b>Race/Sex</b>												
White, not Hispanic Males	67	46%	56	38%	56	43%	51	39%	37	36%	2,146	57%
White, not Hispanic Females	2	1%	6	4%	7	5%	5	4%	3	3%	141	4%
Black/African American, not Hispanic Males	51	35%	45	31%	33	25%	46	35%	38	37%	935	25%
Black/African American, not Hispanic Females	18	12%	19	13%	14	11%	16	12%	12	12%	325	9%
Hispanic/Latino Males	4	3%	7	5%	8	6%	3	2%	6	6%	101	3%
Hispanic/Latina Females	-	-	3	2%	-	-	1	1%	-	-	17	<1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	-	-	2	2%	13	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	1%	3	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	4	3%	9	6%	12	9%	10	8%	4	4%	95	3%
<b>Total</b>	<b>146</b>		<b>147</b>		<b>130</b>		<b>132</b>		<b>103</b>		<b>3,778</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Franklin

Transmission Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	77	62%	80	68%	69	64%	70	64%	62	71%	2,403	74%
Injection drug use (IDU)	-	-	2	2%	2	2%	-	-	-	-	149	5%
Male-to-male sexual contact and IDU	6	5%	4	3%	2	2%	2	2%	-	-	118	4%
Heterosexual contact	9	7%	5	4%	3	3%	8	7%	7	8%	170	5%
Other/unknown	33	26%	26	22%	31	29%	29	27%	18	21%	428	13%
Subtotal	125	100%	117	100%	107	100%	109	100%	87	100%	3,268	100%
<b>Female adult or adolescent</b>												
Injection drug use	2	10%	-	-	1	4%	1	4%	1	6%	64	13%
Heterosexual contact	16	76%	20	69%	18	78%	12	52%	11	69%	327	68%
Other/unknown	3	14%	9	31%	4	17%	10	43%	4	25%	93	19%
Subtotal	21	100%	29	100%	23	100%	23	100%	16	100%	484	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	1	100%	-	-	-	-	-	-	21	81%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	19%
Subtotal	-	-	1	100%	-	-	-	-	-	-	26	100%
<b>Total</b>	<b>146</b>		<b>147</b>		<b>130</b>		<b>132</b>		<b>103</b>		<b>3,778</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Franklin

Exposure Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	73	50%	78	53%	66	51%	68	52%	60	58%	2,325	62%
Injection drug use (IDU) only	-	-	2	1%	3	2%	-	-	-	-	151	4%
Heterosexual contact only	25	17%	25	17%	21	16%	20	15%	18	17%	497	13%
MSM & IDU	5	3%	4	3%	2	2%	2	2%	-	-	101	3%
IDU & Heterosexual contact	2	1%	-	-	-	-	1	1%	1	1%	62	2%
MSM & Heterosexual contact	4	3%	2	1%	3	2%	2	2%	2	2%	78	2%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-	-	-	17	<1%
Perinatal exposure	-	-	1	1%	-	-	-	-	-	-	22	1%
Other/unknown	36	25%	35	24%	35	27%	39	30%	22	21%	525	14%
<b>Total</b>	<b>146</b>		<b>147</b>		<b>130</b>		<b>132</b>		<b>103</b>		<b>3,778</b>	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Franklin

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Franklin**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	630.1	3,781	81%	2,015	81%	1,766	82%
Females	136.4	861	19%	483	19%	378	18%
<b>Age at end of year</b>							
<13	6.0	13	<1%	7	<1%	6	<1%
13-14	*	2	<1%	1	<1%	1	<1%
15-19	19.1	15	<1%	12	<1%	3	<1%
20-24	243.8	226	5%	171	7%	55	3%
25-29	376.5	446	10%	294	12%	152	7%
30-34	437.9	436	9%	284	11%	152	7%
35-39	561.0	472	10%	252	10%	220	10%
40-44	717.9	576	12%	312	12%	264	12%
45-49	1,008.6	772	17%	385	15%	387	18%
50-54	936.3	758	16%	335	13%	423	20%
55-64	530.5	739	16%	352	14%	387	18%
65+	139.8	187	4%	93	4%	94	4%
<b>Race/Ethnicity</b>							
White, not Hispanic	281.1	2,275	49%	1,255	50%	1,020	48%
Black/African American, not Hispanic	714.6	1,909	41%	1,016	41%	893	42%
Hispanic/Latino	362.4	228	5%	114	5%	114	5%
Asian/Pacific Islander	46.4	26	1%	14	1%	12	1%
American Indian/Alaska Native	202.0	5	<1%	4	<1%	1	<1%
Unknown	*	199	4%	95	4%	104	5%
<b>Race/Sex</b>							
White, not Hispanic Males	525.0	2,080	45%	1,135	45%	945	44%
White, not Hispanic Females	47.2	195	4%	120	5%	75	3%
Black/African American, not Hispanic Males	1,055.4	1,329	29%	704	28%	625	29%
Black/African American, not Hispanic Females	410.7	580	12%	312	12%	268	13%
Hispanic/Latino Males	540.1	178	4%	87	3%	91	4%
Hispanic/Latina Females	166.9	50	1%	27	1%	23	1%
Asian/Pacific Islander Males	76.0	21	<1%	11	<1%	10	<1%
Asian/Pacific Islander Females	17.6	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	199	4%	95	4%	104	5%
<b>Total</b>	<b>377.0</b>	<b>4,642</b>		<b>2,498</b>		<b>2,144</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Franklin**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	2,702	72%	1,450	72%	1,252	71%
Injection drug use (IDU)	84	2%	24	1%	60	3%
Male-to-male sexual contact and IDU	121	3%	58	3%	63	4%
Heterosexual contact	196	5%	61	3%	135	8%
Other/unknown	663	18%	414	21%	249	14%
Subtotal	3,766	100%	2,007	100%	1,759	100%
<b>Female adult or adolescent</b>						
Injection drug use	71	9%	36	8%	35	10%
Heterosexual contact	547	66%	283	60%	264	72%
Other/unknown	217	26%	151	32%	66	18%
Subtotal	835	100%	470	100%	365	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	34	83%	18	86%	16	80%
Other/unknown	7	17%	3	14%	4	20%
Subtotal	41	100%	21	100%	20	100%
<b>Total</b>	<b>4,642</b>		<b>2,498</b>		<b>2,144</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Franklin**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,564	55%	1,384	55%	1,180	55%
Injection drug use (IDU) only	80	2%	26	1%	54	3%
Heterosexual contact only	743	16%	344	14%	399	19%
MSM & IDU	97	2%	44	2%	53	2%
IDU & Heterosexual contact	75	2%	34	1%	41	2%
MSM & Heterosexual contact	138	3%	66	3%	72	3%
MSM & IDU & Heterosexual contact	24	1%	14	1%	10	<1%
Perinatal exposure	36	1%	19	1%	17	1%
Other/unknown	885	19%	567	23%	318	15%
<b>Total</b>	<b>4,642</b>		<b>2,498</b>		<b>2,144</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Franklin**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	1,649	79%	813	62%	112	64%	13	62%	3	100%	112	66%
Injection drug use (IDU)	26	1%	52	4%	4	2%	-	-	-	-	2	1%
Male-to-male sexual contact and IDU	70	3%	28	2%	9	5%	1	5%	-	-	13	8%
Heterosexual contact	45	2%	126	10%	18	10%	-	-	-	-	7	4%
Other/unknown	288	14%	299	23%	33	19%	7	33%	-	-	36	21%
<b>Subtotal</b>	<b>2,078</b>	<b>100%</b>	<b>1,318</b>	<b>100%</b>	<b>176</b>	<b>100%</b>	<b>21</b>	<b>100%</b>	<b>3</b>	<b>100%</b>	<b>170</b>	<b>100%</b>
<b>Female adult or adolescent</b>												
Injection drug use	19	10%	43	8%	7	15%	-	-	-	-	2	7%
Heterosexual contact	133	69%	358	64%	34	71%	2	50%	1	50%	19	68%
Other/unknown	41	21%	159	28%	7	15%	2	50%	1	50%	7	25%
<b>Subtotal</b>	<b>193</b>	<b>100%</b>	<b>560</b>	<b>100%</b>	<b>48</b>	<b>100%</b>	<b>4</b>	<b>100%</b>	<b>2</b>	<b>100%</b>	<b>28</b>	<b>100%</b>
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	3	75%	26	84%	4	100%	1	100%	-	-	-	-
Other/unknown	1	25%	5	16%	-	-	-	-	-	-	1	100%
<b>Subtotal</b>	<b>4</b>	<b>100%</b>	<b>31</b>	<b>100%</b>	<b>4</b>	<b>100%</b>	<b>1</b>	<b>100%</b>	<b>-</b>	<b>-</b>	<b>1</b>	<b>100%</b>
<b>Total</b>	<b>2,275</b>		<b>1,909</b>		<b>228</b>		<b>26</b>		<b>5</b>		<b>199</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Franklin**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	1,604	71%	738	39%	102	45%	12	46%	3	60%	105	53%
Injection drug use (IDU) only	20	1%	52	3%	7	3%	-	-	-	-	1	1%
Heterosexual contact only	178	8%	484	25%	52	23%	2	8%	1	20%	26	13%
MSM & IDU	54	2%	23	1%	9	4%	1	4%	-	-	10	5%
IDU & Heterosexual contact	25	1%	43	2%	4	2%	-	-	-	-	3	2%
MSM & Heterosexual contact	45	2%	75	4%	10	4%	1	4%	-	-	7	4%
MSM & IDU & Heterosexual contact	16	1%	5	0%	-	-	-	-	-	-	3	2%
Perinatal exposure	3	0%	27	1%	4	2%	1	4%	-	-	1	1%
Other/unknown	330	15%	462	24%	40	18%	9	35%	1	20%	43	22%
<b>Total</b>	<b>2,275</b>		<b>1,909</b>		<b>228</b>		<b>26</b>		<b>5</b>		<b>199</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Franklin**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	3,009	81%	3,216	81%	3,410	81%	3,599	81%	3,781	81%
Females	704	19%	750	19%	789	19%	832	19%	861	19%
<b>Age at end of year</b>										
<13	12	<1%	12	<1%	13	<1%	14	<1%	13	<1%
13-14	2	<1%	3	<1%	1	<1%	-	-	2	<1%
15-19	31	1%	24	1%	23	1%	18	<1%	15	<1%
20-24	220	6%	229	6%	238	6%	246	6%	226	5%
25-29	298	8%	342	9%	374	9%	408	9%	446	10%
30-34	359	10%	403	10%	426	10%	422	10%	436	9%
35-39	439	12%	425	11%	432	10%	446	10%	472	10%
40-44	643	17%	630	16%	598	14%	588	13%	576	12%
45-49	727	20%	760	19%	775	18%	798	18%	772	17%
50-54	500	13%	564	14%	623	15%	677	15%	758	16%
55-64	407	11%	485	12%	586	14%	668	15%	739	16%
65+	75	2%	89	2%	110	3%	146	3%	187	4%
<b>Race/Ethnicity</b>										
White, not Hispanic	1,895	51%	2,005	51%	2,099	50%	2,194	50%	2,275	49%
Black/African American, not Hispanic	1,476	40%	1,584	40%	1,700	40%	1,805	41%	1,909	41%
Hispanic/Latino	173	5%	191	5%	198	5%	213	5%	228	5%
Asian/Pacific Islander	15	<1%	17	<1%	20	<1%	22	<1%	26	1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	5	<1%
Unknown	150	4%	165	4%	178	4%	193	4%	199	4%
<b>Race/Sex</b>										
White, not Hispanic Males	1,730	47%	1,827	46%	1,915	46%	2,003	45%	2,080	45%
White, not Hispanic Females	165	4%	178	4%	184	4%	191	4%	195	4%
Black/African American, not Hispanic Males	1,001	27%	1,081	27%	1,168	28%	1,245	28%	1,329	29%
Black/African American, not Hispanic Females	475	13%	503	13%	532	13%	560	13%	580	12%
Hispanic/Latino Males	134	4%	148	4%	155	4%	165	4%	178	4%
Hispanic/Latina Females	39	1%	43	1%	43	1%	48	1%	50	1%
Asian/Pacific Islander Males	14	<1%	15	<1%	17	<1%	18	<1%	21	<1%
Asian/Pacific Islander Females	1	<1%	2	<1%	3	<1%	4	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
Unknown	150	4%	165	4%	178	4%	193	4%	199	4%
<b>Total</b>	<b>3,713</b>		<b>3,966</b>		<b>4,199</b>		<b>4,431</b>		<b>4,642</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Franklin

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	2,135	71%	2,286	71%	2,425	71%	2,563	72%	2,702	72%
Injection drug use (IDU)	78	3%	80	2%	83	2%	83	2%	84	2%
Male-to-male sexual contact and IDU	100	3%	104	3%	107	3%	114	3%	121	3%
Heterosexual contact	178	6%	181	6%	185	5%	192	5%	196	5%
Other/unknown	504	17%	551	17%	595	18%	632	18%	663	18%
Subtotal	2,995	100%	3,202	100%	3,395	100%	3,584	100%	3,766	100%
<b>Female adult or adolescent</b>										
Injection drug use	66	10%	66	9%	67	9%	70	9%	71	9%
Heterosexual contact	453	67%	482	66%	505	66%	528	65%	547	66%
Other/unknown	162	24%	178	25%	193	25%	209	26%	217	26%
Subtotal	681	100%	726	100%	765	100%	807	100%	835	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	30	81%	31	82%	32	82%	33	83%	34	83%
Other/unknown	7	19%	7	18%	7	18%	7	18%	7	17%
Subtotal	37	100%	38	100%	39	100%	40	100%	41	100%
<b>Total</b>	<b>3,713</b>		<b>3,966</b>		<b>4,199</b>		<b>4,431</b>		<b>4,642</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Franklin

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	2,023	54%	2,173	55%	2,300	55%	2,431	55%	2,564	55%
Injection drug use (IDU) only	75	2%	77	2%	80	2%	80	2%	80	2%
Heterosexual contact only	631	17%	663	17%	690	16%	720	16%	743	16%
MSM & IDU	83	2%	87	2%	89	2%	93	2%	97	2%
IDU & Heterosexual contact	69	2%	69	2%	70	2%	73	2%	75	2%
MSM & Heterosexual contact	112	3%	113	3%	125	3%	132	3%	138	3%
MSM & IDU & Heterosexual contact	17	<1%	17	<1%	18	<1%	21	<1%	24	1%
Perinatal exposure	32	1%	33	1%	34	1%	35	1%	36	1%
Other/unknown	671	18%	734	19%	793	19%	846	19%	885	19%
<b>Total</b>	<b>3,713</b>		<b>3,966</b>		<b>4,199</b>		<b>4,431</b>		<b>4,642</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.