

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Greene

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Greene

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	152.6	123	81%	57	75%	66	87%
Females	34.9	29	19%	19	25%	10	13%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	1	1%	1	1%	-	-
15-19	*	2	1%	2	3%	-	-
20-24	123.9	18	12%	16	21%	2	3%
25-29	61.1	7	5%	7	9%	-	-
30-34	89.1	9	6%	3	4%	6	8%
35-39	195.7	17	11%	8	11%	9	12%
40-44	149.6	14	9%	2	3%	12	16%
45-49	229.3	23	15%	11	14%	12	16%
50-54	196.2	23	15%	7	9%	16	21%
55-64	141.0	31	20%	17	22%	14	18%
65+	27.4	7	5%	2	3%	5	7%
Race/Ethnicity							
White, not Hispanic	65.8	91	60%	40	53%	51	67%
Black/African American, not Hispanic	478.8	56	37%	34	45%	22	29%
Hispanic/Latino	*	3	2%	-	-	3	4%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	1%	1	1%	-	-
Unknown	*	1	1%	1	1%	-	-
Race/Sex							
White, not Hispanic Males	111.6	76	50%	29	38%	47	62%
White, not Hispanic Females	21.4	15	10%	11	14%	4	5%
Black/African American, not Hispanic Males	753.5	43	28%	27	36%	16	21%
Black/African American, not Hispanic Females	217.1	13	9%	7	9%	6	8%
Hispanic/Latino Males	*	3	2%	-	-	3	4%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	1%	1	1%	-	-
Unknown	*	1	1%	1	1%	-	-
Total	92.8	152		76		76	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Greene

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	87	73%	42	75%	45	70%
Injection drug use (IDU)	4	3%	2	4%	2	3%
Male-to-male sexual contact and IDU	5	4%	1	2%	4	6%
Heterosexual contact	8	7%	2	4%	6	9%
Other/unknown	16	13%	9	16%	7	11%
Subtotal	120	100%	56	100%	64	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	22	81%	13	76%	9	90%
Other/unknown	5	19%	4	24%	1	10%
Subtotal	27	100%	17	100%	10	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	60%	3	100%	-	-
Other/unknown	2	40%	-	-	2	100%
Subtotal	5	100%	3	100%	2	100%
Total	152		76		76	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Greene

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	81	53%	39	51%	42	55%
Injection drug use (IDU) only	3	2%	1	1%	2	3%
Heterosexual contact only	30	20%	15	20%	15	20%
MSM & IDU	4	3%	1	1%	3	4%
IDU & Heterosexual contact	1	1%	1	1%	-	-
MSM & Heterosexual contact	6	4%	3	4%	3	4%
MSM & IDU & Heterosexual contact	1	1%	-	-	1	1%
Perinatal exposure	3	2%	3	4%	-	-
Other/unknown	23	15%	13	17%	10	13%
Total	152		76		76	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Greene

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	56	76%	27	64%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	1	1%	3	7%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	3	4%	2	5%	-	-	-	-	-	-	-	-
Heterosexual contact	4	5%	4	10%	-	-	-	-	-	-	-	-
Other/unknown	10	14%	6	14%	-	-	-	-	-	-	-	-
Subtotal	74	100%	42	100%	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	12	86%	9	75%	-	-	-	-	<10	*	-	-
Other/unknown	2	14%	3	25%	-	-	-	-	-	-	-	-
Subtotal	14	100%	12	100%	-	-	-	-	<10	*	-	-
Child (<13 yrs at diagnosis)												
Perinatal	1	33%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	2	67%	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	-	-	-	-	-	-	-	-
Total	91		56		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Greene

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	52	57%	26	46%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	1	1%	2	4%	-	-	-	-	-	-	-	-
Heterosexual contact only	16	18%	13	23%	-	-	-	-	<10	*	-	-
MSM & IDU	3	3%	1	2%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	2%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	4	4%	1	2%	<10	*	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	2	4%	-	-	-	-	-	-	-	-
Other/unknown	14	15%	9	16%	-	-	-	-	-	-	-	-
Total	91		56		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Greene

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	99	80%	103	79%	106	79%	114	80%	123	81%
Females	24	20%	27	21%	29	21%	29	20%	29	19%
Age at end of year										
<13	1	1%	1	1%	1	1%	1	1%	-	-
13-14	1	1%	1	1%	-	-	-	-	1	1%
15-19	4	3%	2	2%	2	1%	1	1%	2	1%
20-24	8	7%	12	9%	12	9%	16	11%	18	12%
25-29	2	2%	4	3%	5	4%	4	3%	7	5%
30-34	15	12%	14	11%	12	9%	11	8%	9	6%
35-39	11	9%	10	8%	14	10%	14	10%	17	11%
40-44	18	15%	16	12%	16	12%	20	14%	14	9%
45-49	23	19%	26	20%	24	18%	19	13%	23	15%
50-54	19	15%	19	15%	19	14%	23	16%	23	15%
55-64	17	14%	21	16%	25	19%	28	20%	31	20%
65+	4	3%	4	3%	5	4%	6	4%	7	5%
Race/Ethnicity										
White, not Hispanic	79	64%	82	63%	83	61%	88	62%	91	60%
Black/African American, not Hispanic	39	32%	43	33%	47	35%	50	35%	56	37%
Hispanic/Latino	3	2%	3	2%	3	2%	3	2%	3	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	1	1%	1	1%	1	1%	1	1%	1	1%
Race/Sex										
White, not Hispanic Males	66	54%	67	52%	68	50%	73	51%	76	50%
White, not Hispanic Females	13	11%	15	12%	15	11%	15	10%	15	10%
Black/African American, not Hispanic Males	29	24%	32	25%	34	25%	37	26%	43	28%
Black/African American, not Hispanic Females	10	8%	11	8%	13	10%	13	9%	13	9%
Hispanic/Latino Males	3	2%	3	2%	3	2%	3	2%	3	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	1	1%	1	1%	1	1%	1	1%	1	1%
Total	123		130		135		143		152	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Greene

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	69	72%	72	72%	74	72%	79	71%	87	73%
Injection drug use (IDU)	4	4%	4	4%	4	4%	4	4%	4	3%
Male-to-male sexual contact and IDU	5	5%	5	5%	5	5%	5	5%	5	4%
Heterosexual contact	8	8%	8	8%	8	8%	8	7%	8	7%
Other/unknown	10	10%	11	11%	12	12%	15	14%	16	13%
Subtotal	96	100%	100	100%	103	100%	111	100%	120	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	21	95%	22	88%	22	81%	22	81%	22	81%
Other/unknown	1	5%	3	12%	5	19%	5	19%	5	19%
Subtotal	22	100%	25	100%	27	100%	27	100%	27	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	60%	3	60%	3	60%	3	60%	3	60%
Other/unknown	2	40%	2	40%	2	40%	2	40%	2	40%
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	123		130		135		143		152	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Greene

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	63	51%	66	51%	68	50%	73	51%	81	53%
Injection drug use (IDU) only	3	2%	3	2%	3	2%	3	2%	3	2%
Heterosexual contact only	29	24%	30	23%	30	22%	30	21%	30	20%
MSM & IDU	4	3%	4	3%	4	3%	4	3%	4	3%
IDU & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
MSM & Heterosexual contact	6	5%	6	5%	6	4%	6	4%	6	4%
MSM & IDU & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
Perinatal exposure	3	2%	3	2%	3	2%	3	2%	3	2%
Other/unknown	13	11%	16	12%	19	14%	22	15%	23	15%
Total	123		130		135		143		152	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.