

Breakdown of Ryan White Part B and Part A Planning Regions

<p>1 Cleveland Ashtabula Lake Cuyahoga Lorain Geauga</p> <p>2 Columbus Delaware Madison Fairfield Pickaway Franklin Union Licking</p> <p>3 Cincinnati Adams Clinton Brown Hamilton Butler Highland Clermont Warren</p> <p>4 Dayton Clark Miami Darke Montgomery Greene Preble</p> <p>5 Toledo Defiance Ottawa Fulton Sandusky Henry Williams Lucas Wood</p> <p>6 Akron Medina Portage Summit</p> <p>7 Youngstown Columbiana Jefferson Mahoning Trumbull</p> <p>8 Canton Carroll Stark Harrison Tuscarawas Holmes Wayne</p>	<p>9A Lima Allen Mercer Auglaize Paulding Champaign Putnam Hancock Shelby Hardin Van Wert Logan</p> <p>9B Mansfield Ashland Marion Crawford Morrow Erie Richland Huron Seneca Knox Wyandot</p> <p>9C Southeast Athens Monroe Belmont Morgan Coshocton Muskingum Fayette Noble Gallia Perry Guernsey Pike Hocking Ross Jackson Scioto Lawrence Vinton Meigs Washington</p> <hr/> <p>Part A Planning Regions</p> <p>Cleveland Ashtabula Cuyahoga Geauga Lake Lorain Medina</p> <p>Columbus Delaware Madison Fairfield Morrow Franklin Pickaway Licking Union</p>
--	--

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Cleveland Consortium 1

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Cleveland Consortium 1

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	20.6	197	82%	161	85%	26	79%	10	67%	
Females	4.1	42	18%	29	15%	7	21%	5	33%	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	11.3	15	6%	14	7%	1	3%	-	-	
20-24	41.3	50	21%	44	23%	5	15%	1	7%	
25-29	30.0	36	15%	36	19%	-	-	-	-	
30-34	26.9	32	13%	25	13%	5	15%	2	13%	
35-39	19.7	22	9%	16	8%	2	6%	4	27%	
40-44	16.2	21	9%	15	8%	5	15%	1	7%	
45-49	13.5	19	8%	13	7%	3	9%	3	20%	
50-54	14.5	23	10%	14	7%	7	21%	2	13%	
55-64	6.5	18	8%	11	6%	4	12%	2	13%	
65+	*	3	1%	2	1%	1	3%	-	-	
Race/Ethnicity										
White, not Hispanic	4.8	67	28%	55	29%	9	27%	3	20%	
Black/African American, not Hispanic	34.2	140	59%	110	58%	18	55%	11	73%	
Hispanic/Latino	23.2	24	10%	17	9%	6	18%	1	7%	
Asian/Pacific Islander	*	1	<1%	1	1%	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	7	3%	7	4%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	9.0	61	26%	51	27%	7	21%	3	20%	
White, not Hispanic Females	0.8	6	3%	4	2%	2	6%	-	-	
Black/African American, not Hispanic Males	57.3	108	45%	89	47%	13	39%	6	40%	
Black/African American, not Hispanic Females	14.5	32	13%	21	11%	5	15%	5	33%	
Hispanic/Latino Males	40.5	21	9%	14	7%	6	18%	1	7%	
Hispanic/Latina Females	*	3	1%	3	2%	-	-	-	-	
Asian/Pacific Islander Males	*	1	<1%	1	1%	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	7	3%	7	4%	-	-	-	-	
Total	12.0	239		190		33		15		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Cleveland Consortium 1

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	130	66%	113	70%	13	50%	4	40%
Injection drug use (IDU)	4	2%	2	1%	2	8%	-	-
Male-to-male sexual contact and IDU	10	5%	7	4%	2	8%	1	10%
Heterosexual contact	5	3%	3	2%	1	4%	1	10%
Other/unknown	48	24%	36	22%	8	31%	4	40%
Subtotal	197	100%	161	100%	26	100%	10	100%
Female adult or adolescent								
Injection drug use	2	5%	1	4%	1	14%	-	-
Heterosexual contact	12	29%	10	36%	1	14%	1	20%
Other/unknown	27	66%	17	61%	5	71%	4	80%
Subtotal	41	100%	28	100%	7	100%	5	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	239		190		33		15	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	127	53%	110	58%	13	39%	4	27%
Injection drug use (IDU) only	5	2%	3	2%	2	6%	-	-
Heterosexual contact only	17	7%	13	7%	2	6%	2	13%
MSM & IDU	6	3%	5	3%	1	3%	-	-
IDU & Heterosexual contact	1	<1%	-	-	1	3%	-	-
MSM & Heterosexual contact	3	1%	3	2%	-	-	-	-
MSM & IDU & Heterosexual contact	4	2%	2	1%	1	3%	1	7%
Perinatal exposure	1	<1%	1	1%	-	-	-	-
Other/unknown	75	31%	53	28%	13	39%	8	53%
Total	239		190		33		15	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B
Cleveland Consortium 1**

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	38	62%	72	67%	15	71%	-	-	-	-	5	83%
Injection drug use (IDU)	1	2%	1	1%	2	10%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	7%	5	5%	1	5%	-	-	-	-	-	-
Heterosexual contact	1	2%	4	4%	-	-	-	-	-	-	-	-
Other/unknown	17	28%	26	24%	3	14%	1	100%	-	-	1	17%
Subtotal	61	100%	108	100%	21	100%	1	100%	-	-	6	100%
Female adult or adolescent												
Injection drug use	-	-	1	3%	1	33%	-	-	-	-	-	-
Heterosexual contact	2	33%	8	26%	1	33%	-	-	-	-	1	100%
Other/unknown	4	67%	22	71%	1	33%	-	-	-	-	-	-
Subtotal	6	100%	31	100%	3	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	67		140		24		1		0		7	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	37	55%	71	51%	15	63%	-	-	-	-	4	57%
Injection drug use (IDU) only	1	1%	1	1%	3	13%	-	-	-	-	-	-
Heterosexual contact only	3	4%	12	9%	1	4%	-	-	-	-	1	14%
MSM & IDU	2	3%	4	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	1%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	1%	1	1%	-	-	-	-	-	-	1	14%
MSM & IDU & Heterosexual contact	2	3%	1	1%	1	4%	-	-	-	-	-	-
Perinatal exposure	-	-	1	1%	-	-	-	-	-	-	-	-
Other/unknown	21	31%	48	34%	4	17%	1	100%	-	-	1	14%
Total	67		140		24		1		0		7	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics, Part B Cleveland Consortium 1

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	20.6	197	75%	18.2	174	75%	16.5	158	82%	18.6	178	79%
Females	6.2	64	25%	5.6	58	25%	3.4	35	18%	4.5	46	21%
Age at diagnosis (yr)												
<13	*	2	1%	*	2	1%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	1	1%	*	-	-
15-19	9.0	12	5%	8.3	11	5%	14.3	19	10%	10.5	14	6%
20-24	30.6	37	14%	24.8	30	13%	28.9	35	18%	34.7	42	19%
25-29	31.6	38	15%	32.5	39	17%	21.7	26	13%	27.5	33	15%
30-34	24.4	29	11%	13.5	16	7%	21.9	26	13%	24.4	29	13%
35-39	30.5	34	13%	34.1	38	16%	17.9	20	10%	12.6	14	6%
40-44	21.6	28	11%	20.1	26	11%	13.1	17	9%	20.9	27	12%
45-49	24.2	34	13%	20.6	29	13%	14.9	21	11%	19.2	27	12%
50-54	15.1	24	9%	12.6	20	9%	8.8	14	7%	12.0	19	8%
55-64	7.6	21	8%	6.1	17	7%	5.1	14	7%	6.1	17	8%
65+	*	2	1%	*	4	2%	*	-	-	*	2	1%
Race/Ethnicity												
White, not Hispanic	6.2	87	33%	5.3	74	32%	4.1	58	30%	4.6	64	29%
Black/African American, not Hispanic	35.7	146	56%	32.8	134	58%	27.9	114	59%	34.0	139	62%
Hispanic/Latino	23.2	24	9%	17.4	18	8%	15.5	16	8%	14.5	15	7%
Asian/Pacific Islander	*	-	-	*	-	-	*	1	1%	*	1	<1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	2%	*	6	3%	*	4	2%	*	5	2%
Race/Sex												
White, not Hispanic Males	11.3	77	30%	9.4	64	28%	7.7	52	27%	7.9	54	24%
White, not Hispanic Females	1.4	10	4%	1.4	10	4%	0.8	6	3%	1.4	10	4%
Black/African American, not Hispanic Males	54.7	103	39%	49.9	94	41%	47.8	90	47%	57.3	108	48%
Black/African American, not Hispanic Females	19.5	43	16%	18.1	40	17%	10.9	24	12%	14.1	31	14%
Hispanic/Latino Males	27.0	14	5%	25.0	13	6%	21.2	11	6%	21.2	11	5%
Hispanic/Latina Females	19.4	10	4%	9.7	5	2%	9.7	5	3%	*	4	2%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	1	1%	*	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	2%	*	6	3%	*	4	2%	*	5	2%
Total	13.1	261		11.7	232		9.7	193		11.3	224	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Cleveland Consortium 1

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	142	72%	113	66%	106	68%	129	72%
Injection drug use (IDU)	3	2%	7	4%	1	1%	3	2%
Male-to-male sexual contact and IDU	7	4%	5	3%	2	1%	2	1%
Heterosexual contact	17	9%	18	10%	3	2%	5	3%
Other/unknown	28	14%	29	17%	45	29%	39	22%
Subtotal	197	100%	172	100%	157	100%	178	100%
Female adult or adolescent								
Injection drug use	4	6%	3	5%	4	11%	4	9%
Heterosexual contact	45	73%	34	60%	10	29%	16	35%
Other/unknown	13	21%	20	35%	21	60%	26	57%
Subtotal	62	100%	57	100%	35	100%	46	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	50%	2	67%	1	100%	-	-
Other/unknown	1	50%	1	33%	-	-	-	-
Subtotal	2	100%	3	100%	1	100%	-	-
Total	261		232		193		224	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Cleveland Consortium 1

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	138	53%	110	47%	106	55%	129	58%
Injection drug use (IDU) only	4	2%	5	2%	1	1%	3	1%
Heterosexual contact only	62	24%	52	22%	13	7%	21	9%
MSM & IDU	6	2%	4	2%	1	1%	2	1%
IDU & Heterosexual contact	3	1%	5	2%	4	2%	4	2%
MSM & Heterosexual contact	4	2%	3	1%	-	-	-	-
MSM & IDU & Heterosexual contact	1	<1%	1	<1%	1	1%	-	-
Perinatal exposure	1	<1%	3	1%	1	1%	-	-
Other/unknown	42	16%	49	21%	66	34%	65	29%
Total	261		232		193		224	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Cleveland Consortium 1

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	133	74%	117	74%	106	80%	133	83%	161	85%
Females	47	26%	41	26%	27	20%	27	17%	29	15%
Age at HIV diagnosis (yr)										
<13	2	1%	2	1%	-	-	-	-	-	-
13-14	-	-	-	-	1	1%	-	-	-	-
15-19	11	6%	7	4%	16	12%	13	8%	14	7%
20-24	25	14%	24	15%	25	19%	39	24%	44	23%
25-29	28	16%	34	22%	19	14%	24	15%	36	19%
30-34	19	11%	13	8%	16	12%	22	14%	25	13%
35-39	23	13%	22	14%	14	11%	12	8%	16	8%
40-44	20	11%	17	11%	12	9%	17	11%	15	8%
45-49	22	12%	16	10%	13	10%	10	6%	13	7%
50-54	17	9%	13	8%	7	5%	12	8%	14	7%
55-64	12	7%	8	5%	10	8%	10	6%	11	6%
65+	1	1%	2	1%	-	-	1	1%	2	1%
Race/Ethnicity										
White, not Hispanic	62	34%	51	32%	37	28%	46	29%	55	29%
Black/African American, not Hispanic	105	58%	90	57%	82	62%	97	61%	110	58%
Hispanic/Latino	11	6%	13	8%	9	7%	12	8%	17	9%
Asian/Pacific Islander	-	-	-	-	1	1%	1	1%	1	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	1%	4	3%	4	3%	4	3%	7	4%
Race/Sex										
White, not Hispanic Males	55	31%	42	27%	32	24%	39	24%	51	27%
White, not Hispanic Females	7	4%	9	6%	5	4%	7	4%	4	2%
Black/African American, not Hispanic Males	71	39%	64	41%	62	47%	81	51%	89	47%
Black/African American, not Hispanic Females	34	19%	26	16%	20	15%	16	10%	21	11%
Hispanic/Latino Males	5	3%	10	6%	7	5%	9	6%	14	7%
Hispanic/Latina Females	6	3%	3	2%	2	2%	3	2%	3	2%
Asian/Pacific Islander Males	-	-	-	-	1	1%	1	1%	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	1%	4	3%	4	3%	4	3%	7	4%
Total	180		158		133		160		190	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Cleveland Consortium 1

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	96	72%	78	68%	72	69%	99	74%	113	70%
Injection drug use (IDU)	1	1%	3	3%	1	1%	3	2%	2	1%
Male-to-male sexual contact and IDU	2	2%	3	3%	1	1%	1	1%	7	4%
Heterosexual contact	11	8%	11	10%	3	3%	5	4%	3	2%
Other/unknown	23	17%	20	17%	28	27%	25	19%	36	22%
Subtotal	133	100%	115	100%	105	100%	133	100%	161	100%
Female adult or adolescent										
Injection drug use	3	7%	2	5%	2	7%	3	11%	1	4%
Heterosexual contact	33	73%	25	63%	8	30%	10	37%	10	36%
Other/unknown	9	20%	13	33%	17	63%	14	52%	17	61%
Subtotal	45	100%	40	100%	27	100%	27	100%	28	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	50%	2	67%	1	100%	-	-	-	-
Other/unknown	1	50%	1	33%	-	-	-	-	1	100%
Subtotal	2	100%	3	100%	1	100%	-	-	1	100%
Total	180		158		133		160		190	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	92	51%	75	47%	72	54%	99	62%	110	58%
Injection drug use (IDU) only	3	2%	1	1%	-	-	3	2%	3	2%
Heterosexual contact only	44	24%	36	23%	11	8%	15	9%	13	7%
MSM & IDU	2	1%	3	2%	1	1%	1	1%	5	3%
IDU & Heterosexual contact	1	1%	4	3%	3	2%	3	2%	-	-
MSM & Heterosexual contact	4	2%	3	2%	-	-	-	-	3	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	1%
Perinatal exposure	1	1%	3	2%	1	1%	-	-	1	1%
Other/unknown	33	18%	33	21%	45	34%	39	24%	53	28%
Total	180		158		133		160		190	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Cleveland Consortium 1

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	84	75%	88	73%	73	74%	75	77%	58	68%	3,829	82%
Females	28	25%	32	27%	26	26%	22	23%	27	32%	850	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	2	2%	1	1%	1	1%	59	1%
13-14	-	-	-	-	1	1%	-	-	-	-	6	<1%
15-19	2	2%	3	3%	5	5%	4	4%	5	6%	96	2%
20-24	12	11%	16	13%	16	16%	12	12%	11	13%	385	8%
25-29	19	17%	14	12%	13	13%	15	15%	4	5%	789	17%
30-34	19	17%	13	11%	20	20%	10	10%	8	9%	942	20%
35-39	15	13%	24	20%	11	11%	6	6%	13	15%	908	19%
40-44	16	14%	14	12%	8	8%	12	12%	11	13%	608	13%
45-49	14	13%	16	13%	12	12%	20	21%	9	11%	421	9%
50-54	7	6%	8	7%	6	6%	9	9%	12	14%	216	5%
55-64	7	6%	11	9%	5	5%	7	7%	9	11%	197	4%
65+	1	1%	1	1%	-	-	1	1%	2	2%	52	1%
Race/Ethnicity												
White, not Hispanic	34	30%	38	32%	29	29%	27	28%	18	21%	1,857	40%
Black/African American, not Hispanic	59	53%	68	57%	60	61%	64	66%	57	67%	2,388	51%
Hispanic/Latino	17	15%	11	9%	9	9%	5	5%	10	12%	385	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	6	<1%
Unknown	2	2%	3	3%	1	1%	1	1%	-	-	35	1%
Race/Sex												
White, not Hispanic Males	31	28%	34	28%	24	24%	24	25%	14	16%	1,700	36%
White, not Hispanic Females	3	3%	4	3%	5	5%	3	3%	4	5%	157	3%
Black/African American, not Hispanic Males	44	39%	46	38%	43	43%	46	47%	36	42%	1,827	39%
Black/African American, not Hispanic Females	15	13%	22	18%	17	17%	18	19%	21	25%	561	12%
Hispanic/Latino Males	8	7%	6	5%	5	5%	4	4%	8	9%	267	6%
Hispanic/Latina Females	9	8%	5	4%	4	4%	1	1%	2	2%	118	3%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Unknown	2	2%	3	3%	1	1%	1	1%	-	-	35	1%
Total	112		120		99		97		85		4,679	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Cleveland Consortium 1

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	57	68%	53	60%	48	66%	45	61%	33	57%	2,579	68%
Injection drug use (IDU)	2	2%	6	7%	2	3%	1	1%	2	3%	382	10%
Male-to-male sexual contact and IDU	7	8%	6	7%	3	4%	2	3%	4	7%	286	8%
Heterosexual contact	11	13%	12	14%	3	4%	2	3%	2	3%	209	5%
Other/unknown	7	8%	11	13%	17	23%	24	32%	17	29%	349	9%
Subtotal	84	100%	88	100%	73	100%	74	100%	58	100%	3,805	100%
Female adult or adolescent												
Injection drug use	2	7%	4	13%	5	21%	1	5%	2	8%	195	24%
Heterosexual contact	22	79%	23	72%	9	38%	9	41%	10	38%	474	58%
Other/unknown	4	14%	5	16%	10	42%	12	55%	14	54%	146	18%
Subtotal	28	100%	32	100%	24	100%	22	100%	26	100%	815	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	2	100%	1	100%	1	100%	51	86%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	14%
Subtotal	-	-	-	-	2	100%	1	100%	1	100%	59	100%
Total	112		120		99		97		85		4,679	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	55	49%	52	43%	45	45%	43	44%	33	39%	2,429	52%
Injection drug use (IDU) only	2	2%	8	7%	5	5%	1	1%	3	4%	353	8%
Heterosexual contact only	33	29%	35	29%	12	12%	11	11%	12	14%	683	15%
MSM & IDU	6	5%	5	4%	3	3%	1	1%	2	2%	236	5%
IDU & Heterosexual contact	2	2%	2	2%	2	2%	1	1%	1	1%	224	5%
MSM & Heterosexual contact	2	2%	1	1%	3	3%	2	2%	-	-	147	3%
MSM & IDU & Heterosexual contact	1	1%	1	1%	-	-	1	1%	2	2%	50	1%
Perinatal exposure	-	-	-	-	2	2%	1	1%	1	1%	51	1%
Other/unknown	11	10%	16	13%	27	27%	36	37%	31	36%	506	11%
Total	112		120		99		97		85		4,679	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Cleveland Consortium 1

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Cleveland Consortium 1

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	339.0	3,248	77%	1,598	77%	1,650	78%
Females	92.4	954	23%	482	23%	472	22%
Age at end of year							
<13	18.8	58	1%	32	2%	26	1%
13-14	13.3	7	<1%	3	<1%	4	<1%
15-19	121.7	162	4%	96	5%	66	3%
20-24	448.4	543	13%	329	16%	214	10%
25-29	598.0	718	17%	362	17%	356	17%
30-34	661.6	786	19%	353	17%	433	20%
35-39	619.1	690	16%	299	14%	391	18%
40-44	401.9	520	12%	246	12%	274	13%
45-49	261.7	368	9%	161	8%	207	10%
50-54	126.8	201	5%	119	6%	82	4%
55-64	43.8	121	3%	63	3%	58	3%
65+	8.8	28	1%	17	1%	11	1%
Race/Ethnicity							
White, not Hispanic	106.2	1,486	35%	729	35%	757	36%
Black/African American, not Hispanic	546.7	2,236	53%	1,109	53%	1,127	53%
Hispanic/Latino	354.9	367	9%	165	8%	202	10%
Asian/Pacific Islander	35.6	15	<1%	11	1%	4	<1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	94	2%	64	3%	30	1%
Race/Sex							
White, not Hispanic Males	190.4	1,294	31%	623	30%	671	32%
White, not Hispanic Females	26.7	192	5%	106	5%	86	4%
Black/African American, not Hispanic Males	860.4	1,621	39%	802	39%	819	39%
Black/African American, not Hispanic Females	278.8	615	15%	307	15%	308	15%
Hispanic/Latino Males	471.9	245	6%	113	5%	132	6%
Hispanic/Latina Females	236.9	122	3%	52	3%	70	3%
Asian/Pacific Islander Males	59.3	12	<1%	9	<1%	3	<1%
Asian/Pacific Islander Females	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	94	2%	64	3%	30	1%
Total	211.1	4,202		2,080		2,122	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Cleveland Consortium 1

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,148	67%	1,022	65%	1,126	69%
Injection drug use (IDU)	186	6%	67	4%	119	7%
Male-to-male sexual contact and IDU	149	5%	44	3%	105	6%
Heterosexual contact	233	7%	111	7%	122	7%
Other/unknown	502	16%	336	21%	166	10%
Subtotal	3,218	100%	1,580	100%	1,638	100%
Female adult or adolescent						
Injection drug use	120	13%	38	8%	82	18%
Heterosexual contact	571	62%	272	58%	299	65%
Other/unknown	233	25%	156	33%	77	17%
Subtotal	924	100%	466	100%	458	100%
Child (<13 yrs at diagnosis)						
Perinatal	51	85%	27	79%	24	92%
Other/unknown	9	15%	7	21%	2	8%
Subtotal	60	100%	34	100%	26	100%
Total	4,202		2,080		2,122	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,024	48%	987	47%	1,037	49%
Injection drug use (IDU) only	144	3%	53	3%	91	4%
Heterosexual contact only	804	19%	383	18%	421	20%
MSM & IDU	116	3%	40	2%	76	4%
IDU & Heterosexual contact	162	4%	52	3%	110	5%
MSM & Heterosexual contact	124	3%	35	2%	89	4%
MSM & IDU & Heterosexual contact	33	1%	4	<1%	29	1%
Perinatal exposure	52	1%	28	1%	24	1%
Other/unknown	743	18%	498	24%	245	12%
Total	4,202		2,080		2,122	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Cleveland Consortium 1

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	997	77%	1,007	63%	104	44%	5	42%	1	50%	34	46%
Injection drug use (IDU)	42	3%	84	5%	58	24%	1	8%	-	-	1	1%
Male-to-male sexual contact and IDU	68	5%	65	4%	13	5%	-	-	-	-	3	4%
Heterosexual contact	42	3%	157	10%	28	12%	-	-	-	-	6	8%
Other/unknown	140	11%	289	18%	36	15%	6	50%	1	50%	30	41%
Subtotal	1,289	100%	1,602	100%	239	100%	12	100%	2	100%	74	100%
Female adult or adolescent												
Injection drug use	27	14%	67	11%	25	21%	-	-	-	-	1	5%
Heterosexual contact	119	63%	372	63%	70	59%	1	33%	1	50%	8	40%
Other/unknown	42	22%	154	26%	23	19%	2	67%	1	50%	11	55%
Subtotal	188	100%	593	100%	118	100%	3	100%	2	100%	20	100%
Child (<13 yrs at diagnosis)												
Perinatal	4	44%	37	90%	10	100%	-	-	-	-	-	-
Other/unknown	5	56%	4	10%	-	-	-	-	-	-	-	-
Subtotal	9	100%	41	100%	10	100%	-	-	-	-	-	-
Total	1,486		2,236		367		15		4		94	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Cleveland Consortium 1

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	968	65%	923	41%	97	26%	4	27%	1	25%	31	33%
Injection drug use (IDU) only	35	2%	63	3%	44	12%	1	7%	-	-	1	1%
Heterosexual contact only	161	11%	529	24%	98	27%	1	7%	1	25%	14	15%
MSM & IDU	58	4%	46	2%	10	3%	-	-	-	-	2	2%
IDU & Heterosexual contact	34	2%	88	4%	39	11%	-	-	-	-	1	1%
MSM & Heterosexual contact	29	2%	84	4%	7	2%	1	7%	-	-	3	3%
MSM & IDU & Heterosexual contact	10	1%	19	1%	3	1%	-	-	-	-	1	1%
Perinatal exposure	5	0%	37	2%	10	3%	-	-	-	-	-	-
Other/unknown	186	13%	447	20%	59	16%	8	53%	2	50%	41	44%
Total	1,486		2,236		367		15		4		94	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Cleveland Consortium 1

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	2,575	77%	2,756	77%	2,920	77%	3,073	77%	3,248	77%
Females	760	23%	819	23%	874	23%	908	23%	954	23%
Age at end of year										
<13	54	2%	56	2%	58	2%	58	1%	58	1%
13-14	6	<1%	6	<1%	6	<1%	7	<1%	7	<1%
15-19	108	3%	120	3%	131	3%	149	4%	162	4%
20-24	403	12%	439	12%	467	12%	501	13%	543	13%
25-29	586	18%	621	17%	660	17%	685	17%	718	17%
30-34	685	21%	714	20%	731	19%	757	19%	786	19%
35-39	586	18%	620	17%	657	17%	676	17%	690	16%
40-44	425	13%	451	13%	477	13%	493	12%	520	12%
45-49	267	8%	297	8%	322	8%	343	9%	368	9%
50-54	126	4%	149	4%	168	4%	182	5%	201	5%
55-64	68	2%	79	2%	91	2%	104	3%	121	3%
65+	21	1%	23	1%	26	1%	26	1%	28	1%
Race/Ethnicity										
White, not Hispanic	1,221	37%	1,301	36%	1,369	36%	1,423	36%	1,486	35%
Black/African American, not Hispanic	1,725	52%	1,859	52%	1,987	52%	2,099	53%	2,236	53%
Hispanic/Latino	295	9%	318	9%	336	9%	352	9%	367	9%
Asian/Pacific Islander	13	<1%	13	<1%	13	<1%	14	<1%	15	<1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	77	2%	80	2%	85	2%	89	2%	94	2%
Race/Sex										
White, not Hispanic Males	1,062	32%	1,133	32%	1,192	31%	1,241	31%	1,294	31%
White, not Hispanic Females	159	5%	168	5%	177	5%	182	5%	192	5%
Black/African American, not Hispanic Males	1,244	37%	1,337	37%	1,427	38%	1,515	38%	1,621	39%
Black/African American, not Hispanic Females	481	14%	522	15%	560	15%	584	15%	615	15%
Hispanic/Latino Males	196	6%	210	6%	223	6%	234	6%	245	6%
Hispanic/Latina Females	99	3%	108	3%	113	3%	118	3%	122	3%
Asian/Pacific Islander Males	10	<1%	10	<1%	10	<1%	11	<1%	12	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	77	2%	80	2%	85	2%	89	2%	94	2%
Total	3,335		3,575		3,794		3,981		4,202	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Cleveland Consortium 1

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,674	66%	1,804	66%	1,918	66%	2,021	66%	2,148	67%
Injection drug use (IDU)	172	7%	176	6%	182	6%	183	6%	186	6%
Male-to-male sexual contact and IDU	133	5%	141	5%	145	5%	147	5%	149	5%
Heterosexual contact	191	7%	207	8%	225	8%	228	7%	233	7%
Other/unknown	378	15%	401	15%	421	15%	464	15%	502	16%
Subtotal	2,548	100%	2,729	100%	2,891	100%	3,043	100%	3,218	100%
Female adult or adolescent										
Injection drug use	106	14%	109	14%	112	13%	116	13%	120	13%
Heterosexual contact	469	64%	511	65%	545	65%	555	63%	571	62%
Other/unknown	158	22%	170	22%	187	22%	207	24%	233	25%
Subtotal	733	100%	790	100%	844	100%	878	100%	924	100%
Child (<13 yrs at diagnosis)										
Perinatal	47	87%	48	86%	50	85%	51	85%	51	85%
Other/unknown	7	13%	8	14%	9	15%	9	15%	9	15%
Subtotal	54	100%	56	100%	59	100%	60	100%	60	100%
Total	3,335		3,575		3,794		3,981		4,202	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Cleveland Consortium 1

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	1,557	47%	1,683	47%	1,794	47%	1,897	48%	2,024	48%
Injection drug use (IDU) only	132	4%	136	4%	140	4%	141	4%	144	3%
Heterosexual contact only	660	20%	718	20%	770	20%	783	20%	804	19%
MSM & IDU	103	3%	110	3%	113	3%	114	3%	116	3%
IDU & Heterosexual contact	146	4%	149	4%	154	4%	158	4%	162	4%
MSM & Heterosexual contact	117	4%	121	3%	124	3%	124	3%	124	3%
MSM & IDU & Heterosexual contact	30	1%	31	1%	32	1%	33	1%	33	1%
Perinatal exposure	47	1%	48	1%	51	1%	52	1%	52	1%
Other/unknown	543	16%	579	16%	616	16%	679	17%	743	18%
Total	3,335		3,575		3,794		3,981		4,202	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Columbus Consortium 2

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Columbus Consortium 2

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	28.1	259	84%	182	82%	48	86%	29	94%	
Females	5.2	50	16%	40	18%	8	14%	2	6%	
Age at diagnosis (yr)										
<13	*	2	1%	2	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	11.9	15	5%	13	6%	1	2%	1	3%	
20-24	44.1	61	20%	52	23%	5	9%	4	13%	
25-29	37.1	53	17%	40	18%	10	18%	3	10%	
30-34	34.1	47	15%	32	14%	6	11%	9	29%	
35-39	22.3	28	9%	16	7%	9	16%	3	10%	
40-44	25.4	34	11%	22	10%	9	16%	3	10%	
45-49	22.4	29	9%	18	8%	6	11%	5	16%	
50-54	12.7	17	6%	12	5%	4	7%	1	3%	
55-64	9.2	20	6%	13	6%	5	9%	2	6%	
65+	*	3	1%	2	1%	1	2%	-	-	
Race/Ethnicity										
White, not Hispanic	10.0	142	46%	100	45%	27	48%	15	48%	
Black/African American, not Hispanic	47.5	134	43%	104	47%	20	36%	10	32%	
Hispanic/Latino	14.0	10	3%	5	2%	3	5%	2	6%	
Asian/Pacific Islander	*	4	1%	4	2%	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	19	6%	9	4%	6	11%	4	13%	
Race/Sex										
White, not Hispanic Males	19.1	133	43%	92	41%	27	48%	14	45%	
White, not Hispanic Females	1.3	9	3%	8	4%	-	-	1	3%	
Black/African American, not Hispanic Males	72.3	98	32%	75	34%	14	25%	9	29%	
Black/African American, not Hispanic Females	24.5	36	12%	29	13%	6	11%	1	3%	
Hispanic/Latino Males	26.6	10	3%	5	2%	3	5%	2	6%	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	3	1%	3	1%	-	-	-	-	
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	19	6%	9	4%	6	11%	4	13%	
Total	16.4	309		222		56		31		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Columbus Consortium 2

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	168	65%	118	65%	35	73%	15	52%
Injection drug use (IDU)	2	1%	1	1%	1	2%	-	-
Male-to-male sexual contact and IDU	3	1%	2	1%	-	-	1	3%
Heterosexual contact	5	2%	4	2%	1	2%	-	-
Other/unknown	80	31%	56	31%	11	23%	13	45%
Subtotal	258	100%	181	100%	48	100%	29	100%
Female adult or adolescent								
Injection drug use	1	2%	1	3%	-	-	-	-
Heterosexual contact	12	24%	10	26%	1	13%	1	50%
Other/unknown	36	73%	28	72%	7	88%	1	50%
Subtotal	49	100%	39	100%	8	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	2	100%	2	100%	-	-	-	-
Subtotal	2	100%	2	100%	-	-	-	-
Total	309		222		56		31	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Columbus Consortium 2

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	155	50%	108	49%	33	59%	14	45%
Injection drug use (IDU) only	3	1%	2	1%	1	2%	-	-
Heterosexual contact only	17	6%	14	6%	2	4%	1	3%
MSM & IDU	3	1%	2	1%	-	-	1	3%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	13	4%	10	5%	2	4%	1	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	118	38%	86	39%	18	32%	14	45%
Total	309		222		56		31	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Columbus Consortium 2

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	96	72%	55	57%	7	70%	1	33%	-	-	9	60%
Injection drug use (IDU)	1	1%	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	2%	-	-	-	-	-	-	-	-	1	7%
Heterosexual contact	1	1%	4	4%	-	-	-	-	-	-	-	-
Other/unknown	33	25%	37	38%	3	30%	2	67%	-	-	5	33%
Subtotal	133	100%	97	100%	10	100%	3	100%	-	-	15	100%
Female adult or adolescent												
Injection drug use	-	-	1	3%	-	-	-	-	-	-	-	-
Heterosexual contact	6	67%	5	14%	-	-	-	-	-	-	1	25%
Other/unknown	3	33%	29	83%	-	-	1	100%	-	-	3	75%
Subtotal	9	100%	35	100%	-	-	1	100%	-	-	4	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	2	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	-	-	-	-	-	-	-	-
Total	142		134		10		4		0		19	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Columbus Consortium 2

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	89	63%	50	37%	6	60%	1	25%	-	-	9	47%
Injection drug use (IDU) only	1	1%	2	1%	-	-	-	-	-	-	-	-
Heterosexual contact only	7	5%	9	7%	-	-	-	-	-	-	1	5%
MSM & IDU	2	1%	-	-	-	-	-	-	-	-	1	5%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	7	5%	5	4%	1	10%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	36	25%	68	51%	3	30%	3	75%	-	-	8	42%
Total	142		134		10		4		0		19	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Columbus Consortium 2**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	26.5	245	84%	26.1	241	81%	26.5	245	86%	25.5	235	83%
Females	4.8	46	16%	6.0	57	19%	4.1	39	14%	5.0	48	17%
Age at diagnosis (yr)												
<13	*	2	1%	*	2	1%	*	-	-	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	10.3	13	4%	21.4	27	9%	10.3	13	5%	13.5	17	6%
20-24	35.4	49	17%	37.6	52	17%	43.4	60	21%	39.8	55	19%
25-29	25.9	37	13%	39.9	57	19%	30.1	43	15%	30.8	44	16%
30-34	31.9	44	15%	25.4	35	12%	29.0	40	14%	33.4	46	16%
35-39	31.1	39	13%	32.7	41	14%	30.3	38	13%	31.1	39	14%
40-44	31.3	42	14%	27.6	37	12%	27.6	37	13%	22.4	30	11%
45-49	22.4	29	10%	13.9	18	6%	19.3	25	9%	16.2	21	7%
50-54	18.7	25	9%	10.5	14	5%	9.0	12	4%	11.2	15	5%
55-64	2.8	6	2%	5.5	12	4%	6.9	15	5%	5.5	12	4%
65+	2.4	5	2%	*	3	1%	*	1	<1%	*	3	1%
Race/Ethnicity												
White, not Hispanic	11.0	156	54%	10.9	154	52%	10.2	144	51%	9.5	134	47%
Black/African American, not Hispanic	40.7	115	40%	43.9	124	42%	41.5	117	41%	41.8	118	42%
Hispanic/Latino	14.0	10	3%	14.0	10	3%	19.6	14	5%	23.8	17	6%
Asian/Pacific Islander	*	2	1%	*	2	1%	*	2	1%	*	2	1%
American Indian/Alaska Native	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	7	2%	*	8	3%	*	7	2%	*	12	4%
Race/Sex												
White, not Hispanic Males	20.4	142	49%	20.0	139	47%	19.1	133	47%	17.5	122	43%
White, not Hispanic Females	1.9	14	5%	2.1	15	5%	1.5	11	4%	1.7	12	4%
Black/African American, not Hispanic Males	63.4	86	30%	63.4	86	29%	67.9	92	32%	64.2	87	31%
Black/African American, not Hispanic Females	19.8	29	10%	25.9	38	13%	17.0	25	9%	21.1	31	11%
Hispanic/Latino Males	24.0	9	3%	24.0	9	3%	34.6	13	5%	34.6	13	5%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	1	<1%	*	4	1%
Asian/Pacific Islander Males	*	2	1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Females	*	-	-	*	1	<1%	*	1	<1%	*	1	<1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	7	2%	*	8	3%	*	7	2%	*	12	4%
Total	15.5	291		15.9	298		15.1	284		15.1	283	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Columbus Consortium 2

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	191	78%	178	74%	171	70%	163	69%
Injection drug use (IDU)	2	1%	1	<1%	-	-	3	1%
Male-to-male sexual contact and IDU	9	4%	7	3%	8	3%	5	2%
Heterosexual contact	27	11%	18	7%	10	4%	4	2%
Other/unknown	16	7%	37	15%	56	23%	60	26%
Subtotal	245	100%	241	100%	245	100%	235	100%
Female adult or adolescent								
Injection drug use	3	7%	1	2%	3	8%	-	-
Heterosexual contact	25	58%	36	65%	14	36%	16	34%
Other/unknown	15	35%	18	33%	22	56%	31	66%
Subtotal	43	100%	55	100%	39	100%	47	100%
Child (<13 yrs at diagnosis)								
Perinatal	2	67%	2	100%	-	-	1	100%
Other/unknown	1	33%	-	-	-	-	-	-
Subtotal	3	100%	2	100%	-	-	1	100%
Total	291		298		284		283	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Columbus Consortium 2

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	185	64%	171	57%	166	58%	161	57%
Injection drug use (IDU) only	1	<1%	-	-	-	-	2	1%
Heterosexual contact only	52	18%	54	18%	24	8%	20	7%
MSM & IDU	8	3%	4	1%	8	3%	5	2%
IDU & Heterosexual contact	4	1%	2	1%	3	1%	1	<1%
MSM & Heterosexual contact	6	2%	7	2%	5	2%	2	1%
MSM & IDU & Heterosexual contact	1	<1%	3	1%	-	-	-	-
Perinatal exposure	3	1%	2	1%	-	-	1	<1%
Other/unknown	31	11%	55	18%	78	27%	91	32%
Total	291		298		284		283	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Columbus Consortium 2

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	153	86%	154	81%	158	86%	156	83%	182	82%
Females	25	14%	37	19%	25	14%	32	17%	40	18%
Age at HIV diagnosis (yr)										
<13	1	1%	1	1%	-	-	-	-	2	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	10	6%	21	11%	10	5%	14	7%	13	6%
20-24	30	17%	42	22%	49	27%	41	22%	52	23%
25-29	24	13%	36	19%	35	19%	34	18%	40	18%
30-34	25	14%	18	9%	21	11%	32	17%	32	14%
35-39	23	13%	26	14%	23	13%	15	8%	16	7%
40-44	27	15%	27	14%	18	10%	19	10%	22	10%
45-49	17	10%	8	4%	13	7%	13	7%	18	8%
50-54	14	8%	5	3%	6	3%	11	6%	12	5%
55-64	4	2%	6	3%	7	4%	6	3%	13	6%
65+	3	2%	1	1%	1	1%	3	2%	2	1%
Race/Ethnicity										
White, not Hispanic	101	57%	104	54%	90	49%	95	51%	100	45%
Black/African American, not Hispanic	63	35%	74	39%	76	42%	76	40%	104	47%
Hispanic/Latino	9	5%	6	3%	10	5%	8	4%	5	2%
Asian/Pacific Islander	1	1%	1	1%	2	1%	1	1%	4	2%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	4	2%	6	3%	5	3%	8	4%	9	4%
Race/Sex										
White, not Hispanic Males	93	52%	92	48%	81	44%	87	46%	92	41%
White, not Hispanic Females	8	4%	12	6%	9	5%	8	4%	8	4%
Black/African American, not Hispanic Males	47	26%	52	27%	62	34%	54	29%	75	34%
Black/African American, not Hispanic Females	16	9%	22	12%	14	8%	22	12%	29	13%
Hispanic/Latino Males	9	5%	5	3%	9	5%	7	4%	5	2%
Hispanic/Latina Females	-	-	1	1%	1	1%	1	1%	-	-
Asian/Pacific Islander Males	1	1%	1	1%	1	1%	-	-	3	1%
Asian/Pacific Islander Females	-	-	-	-	1	1%	1	1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	4	2%	6	3%	5	3%	8	4%	9	4%
Total	178		191		183		188		222	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Columbus Consortium 2

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	122	80%	118	77%	117	74%	109	70%	118	65%
Injection drug use (IDU)	1	1%	-	-	-	-	1	1%	1	1%
Male-to-male sexual contact and IDU	4	3%	4	3%	5	3%	3	2%	2	1%
Heterosexual contact	14	9%	9	6%	6	4%	2	1%	4	2%
Other/unknown	12	8%	23	15%	30	19%	41	26%	56	31%
Subtotal	153	100%	154	100%	158	100%	156	100%	181	100%
Female adult or adolescent										
Injection drug use	-	-	1	3%	2	8%	-	-	1	3%
Heterosexual contact	15	65%	24	67%	9	36%	12	38%	10	26%
Other/unknown	8	35%	11	31%	14	56%	20	63%	28	72%
Subtotal	23	100%	36	100%	25	100%	32	100%	39	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	50%	1	100%	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	-	-	2	100%
Subtotal	2	100%	1	100%	-	-	-	-	2	100%
Total	178		191		183		188		222	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Columbus Consortium 2

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	120	67%	114	60%	114	62%	107	57%	108	49%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	2	1%
Heterosexual contact only	29	16%	33	17%	15	8%	14	7%	14	6%
MSM & IDU	4	2%	2	1%	5	3%	3	2%	2	1%
IDU & Heterosexual contact	1	1%	1	1%	2	1%	1	1%	-	-
MSM & Heterosexual contact	2	1%	4	2%	3	2%	2	1%	10	5%
MSM & IDU & Heterosexual contact	-	-	2	1%	-	-	-	-	-	-
Perinatal exposure	2	1%	1	1%	-	-	-	-	-	-
Other/unknown	20	11%	34	18%	44	24%	61	32%	86	39%
Total	178		191		183		188		222	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Columbus Consortium 2

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	110	83%	125	78%	137	86%	129	81%	126	82%	3,441	87%
Females	22	17%	35	22%	22	14%	31	19%	28	18%	510	13%
Age at AIDS diagnosis (yr)												
<13	3	2%	1	1%	-	-	1	1%	-	-	24	1%
13-14	-	-	-	-	-	-	-	-	-	-	6	<1%
15-19	4	3%	7	4%	10	6%	8	5%	4	3%	74	2%
20-24	17	13%	19	12%	20	13%	22	14%	22	14%	366	9%
25-29	14	11%	29	18%	17	11%	16	10%	26	17%	679	17%
30-34	35	27%	24	15%	31	19%	26	16%	25	16%	899	23%
35-39	21	16%	24	15%	24	15%	38	24%	24	16%	718	18%
40-44	13	10%	20	13%	27	17%	23	14%	17	11%	518	13%
45-49	13	10%	16	10%	14	9%	13	8%	18	12%	334	8%
50-54	10	8%	13	8%	8	5%	6	4%	7	5%	184	5%
55-64	2	2%	4	3%	7	4%	7	4%	10	6%	127	3%
65+	-	-	3	2%	1	1%	-	-	1	1%	22	1%
Race/Ethnicity												
White, not Hispanic	63	48%	69	43%	82	52%	74	46%	81	53%	2,531	64%
Black/African American, not Hispanic	55	42%	79	49%	70	44%	66	41%	51	33%	1,227	31%
Hispanic/Latino	5	4%	6	4%	4	3%	9	6%	9	6%	112	3%
Asian/Pacific Islander	1	1%	1	1%	-	-	1	1%	-	-	13	<1%
American Indian/Alaska Native	-	-	-	-	-	-	1	1%	-	-	2	<1%
Unknown	8	6%	5	3%	3	2%	9	6%	13	8%	66	2%
Race/Sex												
White, not Hispanic Males	59	45%	60	38%	80	50%	67	42%	71	46%	2,363	60%
White, not Hispanic Females	4	3%	9	6%	2	1%	7	4%	10	6%	168	4%
Black/African American, not Hispanic Males	40	30%	55	34%	51	32%	47	29%	36	23%	917	23%
Black/African American, not Hispanic Females	15	11%	24	15%	19	12%	19	12%	15	10%	310	8%
Hispanic/Latino Males	4	3%	6	4%	4	3%	6	4%	8	5%	93	2%
Hispanic/Latina Females	1	1%	-	-	-	-	3	2%	1	1%	19	<1%
Asian/Pacific Islander Males	1	1%	-	-	-	-	1	1%	-	-	11	<1%
Asian/Pacific Islander Females	-	-	1	1%	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	1	1%	-	-	2	<1%
Unknown	8	6%	5	3%	3	2%	9	6%	13	8%	66	2%
Total	132		160		159		160		154		3,951	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Columbus Consortium 2

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	74	68%	91	73%	82	60%	87	67%	81	64%	2,479	72%
Injection drug use (IDU)	3	3%	3	2%	-	-	2	2%	2	2%	183	5%
Male-to-male sexual contact and IDU	7	6%	3	2%	6	4%	4	3%	3	2%	137	4%
Heterosexual contact	15	14%	16	13%	9	7%	5	4%	3	2%	178	5%
Other/unknown	10	9%	12	10%	40	29%	31	24%	37	29%	452	13%
Subtotal	109	100%	125	100%	137	100%	129	100%	126	100%	3,429	100%
Female adult or adolescent												
Injection drug use	1	5%	2	6%	2	9%	-	-	1	4%	72	15%
Heterosexual contact	12	60%	20	59%	11	50%	14	47%	12	43%	267	54%
Other/unknown	7	35%	12	35%	9	41%	16	53%	15	54%	155	31%
Subtotal	20	100%	34	100%	22	100%	30	100%	28	100%	494	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	67%	1	100%	-	-	1	100%	-	-	21	75%
Other/unknown	1	33%	-	-	-	-	-	-	-	-	7	25%
Subtotal	3	100%	1	100%	-	-	1	100%	-	-	28	100%
Total	132		160		159		160		154		3,951	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Columbus Consortium 2

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	69	52%	82	51%	78	49%	85	53%	77	50%	2,401	61%
Injection drug use (IDU) only	2	2%	1	1%	-	-	2	1%	3	2%	179	5%
Heterosexual contact only	27	20%	36	23%	20	13%	19	12%	15	10%	445	11%
MSM & IDU	4	3%	3	2%	5	3%	4	3%	3	2%	117	3%
IDU & Heterosexual contact	2	2%	4	3%	2	1%	-	-	-	-	76	2%
MSM & Heterosexual contact	5	4%	9	6%	4	3%	2	1%	4	3%	78	2%
MSM & IDU & Heterosexual contact	3	2%	-	-	1	1%	-	-	-	-	20	1%
Perinatal exposure	2	2%	1	1%	-	-	1	1%	-	-	22	1%
Other/unknown	18	14%	24	15%	49	31%	47	29%	52	34%	613	16%
Total	132		160		159		160		154		3,951	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Columbus Consortium 2

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Columbus Consortium 2

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	367.4	3,391	82%	1,844	81%	1,547	83%
Females	78.5	750	18%	423	19%	327	17%
Age at end of year							
<13	9.6	32	1%	18	1%	14	1%
13-14	16.1	8	<1%	4	<1%	4	<1%
15-19	138.1	174	4%	118	5%	56	3%
20-24	399.0	552	13%	327	14%	225	12%
25-29	490.8	701	17%	405	18%	296	16%
30-34	564.1	777	19%	367	16%	410	22%
35-39	602.3	755	18%	415	18%	340	18%
40-44	393.2	527	13%	285	13%	242	13%
45-49	240.4	311	8%	156	7%	155	8%
50-54	137.7	184	4%	99	4%	85	5%
55-64	47.1	102	2%	60	3%	42	2%
65+	9.0	19	<1%	14	1%	5	<1%
Race/Ethnicity							
White, not Hispanic	165.7	2,345	57%	1,300	57%	1,045	56%
Black/African American, not Hispanic	532.5	1,503	36%	816	36%	687	37%
Hispanic/Latino	211.7	151	4%	71	3%	80	4%
Asian/Pacific Islander	32.8	21	1%	13	1%	8	<1%
American Indian/Alaska Native	*	4	<1%	3	<1%	1	<1%
Unknown	*	118	3%	65	3%	53	3%
Race/Sex							
White, not Hispanic Males	304.4	2,118	51%	1,167	51%	951	51%
White, not Hispanic Females	31.6	227	5%	133	6%	94	5%
Black/African American, not Hispanic Males	763.5	1,035	25%	554	24%	481	26%
Black/African American, not Hispanic Females	319.1	468	11%	262	12%	206	11%
Hispanic/Latino Males	324.7	122	3%	59	3%	63	3%
Hispanic/Latina Females	85.9	29	1%	12	1%	17	1%
Asian/Pacific Islander Males	47.7	15	<1%	9	<1%	6	<1%
Asian/Pacific Islander Females	18.4	6	<1%	4	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	118	3%	65	3%	53	3%
Total	220.5	4,142		2,268		1,874	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Columbus Consortium 2

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,331	69%	1,264	69%	1,067	69%
Injection drug use (IDU)	74	2%	19	1%	55	4%
Male-to-male sexual contact and IDU	92	3%	38	2%	54	4%
Heterosexual contact	219	6%	82	4%	137	9%
Other/unknown	660	20%	432	24%	228	15%
Subtotal	3,376	100%	1,835	100%	1,541	100%
Female adult or adolescent						
Injection drug use	61	8%	27	7%	34	11%
Heterosexual contact	391	54%	201	49%	190	60%
Other/unknown	277	38%	185	45%	92	29%
Subtotal	729	100%	413	100%	316	100%
Child (<13 yrs at diagnosis)						
Perinatal	28	78%	15	79%	13	76%
Other/unknown	8	22%	4	21%	4	24%
Subtotal	36	100%	19	100%	17	100%
Total	4,142		2,268		1,874	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Columbus Consortium 2

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,241	54%	1,227	54%	1,014	54%
Injection drug use (IDU) only	67	2%	17	1%	50	3%
Heterosexual contact only	610	15%	283	12%	327	17%
MSM & IDU	79	2%	36	2%	43	2%
IDU & Heterosexual contact	68	2%	29	1%	39	2%
MSM & Heterosexual contact	90	2%	37	2%	53	3%
MSM & IDU & Heterosexual contact	13	<1%	2	<1%	11	1%
Perinatal exposure	30	1%	16	1%	14	1%
Other/unknown	944	23%	621	27%	323	17%
Total	4,142		2,268		1,874	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Columbus Consortium 2

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	1,618	76%	569	56%	76	63%	8	57%	3	100%	57	58%
Injection drug use (IDU)	30	1%	37	4%	5	4%	-	-	-	-	2	2%
Male-to-male sexual contact and IDU	66	3%	22	2%	-	-	1	7%	-	-	3	3%
Heterosexual contact	60	3%	138	13%	16	13%	1	7%	-	-	4	4%
Other/unknown	342	16%	259	25%	24	20%	4	29%	-	-	32	33%
Subtotal	2,116	100%	1,025	100%	121	100%	14	100%	3	100%	98	100%
Female adult or adolescent												
Injection drug use	23	10%	31	7%	6	22%	-	-	-	-	1	6%
Heterosexual contact	128	58%	235	52%	17	63%	3	50%	-	-	8	47%
Other/unknown	71	32%	190	42%	4	15%	3	50%	1	100%	8	47%
Subtotal	222	100%	456	100%	27	100%	6	100%	1	100%	17	100%
Child (<13 yrs at diagnosis)												
Perinatal	5	71%	17	77%	3	100%	1	100%	-	-	2	67%
Other/unknown	2	29%	5	23%	-	-	-	-	-	-	1	33%
Subtotal	7	100%	22	100%	3	100%	1	100%	-	-	3	100%
Total	2,345		1,503		151		21		4		118	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Columbus Consortium 2

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	1,586	68%	520	35%	71	47%	8	38%	3	75%	53	45%
Injection drug use (IDU) only	24	1%	32	2%	10	7%	-	-	-	-	1	1%
Heterosexual contact only	188	8%	373	25%	33	22%	4	19%	-	-	12	10%
MSM & IDU	58	2%	17	1%	-	-	1	5%	-	-	3	3%
IDU & Heterosexual contact	29	1%	36	2%	1	1%	-	-	-	-	2	2%
MSM & Heterosexual contact	32	1%	49	3%	5	3%	-	-	-	-	4	3%
MSM & IDU & Heterosexual contact	8	0%	5	0%	-	-	-	-	-	-	-	-
Perinatal exposure	5	0%	18	1%	3	2%	1	5%	-	-	3	3%
Other/unknown	415	18%	453	30%	28	19%	7	33%	1	25%	40	34%
Total	2,345		1,503		151		21		4		118	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Columbus Consortium 2

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	2,473	81%	2,698	81%	2,925	81%	3,159	82%	3,391	82%
Females	567	19%	612	18%	666	19%	704	18%	750	18%
Age at end of year										
<13	27	1%	29	1%	31	1%	31	1%	32	1%
13-14	8	<1%	8	<1%	8	<1%	8	<1%	8	<1%
15-19	104	3%	116	4%	143	4%	157	4%	174	4%
20-24	343	11%	391	12%	439	12%	497	13%	552	13%
25-29	525	17%	561	17%	617	17%	659	17%	701	17%
30-34	620	20%	660	20%	692	19%	731	19%	777	19%
35-39	600	20%	638	19%	679	19%	716	19%	755	18%
40-44	393	13%	429	13%	462	13%	498	13%	527	13%
45-49	228	7%	253	8%	269	7%	292	8%	311	8%
50-54	123	4%	146	4%	159	4%	169	4%	184	4%
55-64	62	2%	68	2%	78	2%	90	2%	102	2%
65+	8	<1%	12	<1%	15	<1%	16	<1%	19	<1%
Race/Ethnicity										
White, not Hispanic	1,787	59%	1,933	58%	2,077	58%	2,215	57%	2,345	57%
Black/African American, not Hispanic	1,053	35%	1,158	35%	1,275	35%	1,387	36%	1,503	36%
Hispanic/Latino	100	3%	109	3%	119	3%	133	3%	151	4%
Asian/Pacific Islander	14	<1%	16	<1%	18	1%	19	<1%	21	1%
American Indian/Alaska Native	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	84	3%	91	3%	99	3%	106	3%	118	3%
Race/Sex										
White, not Hispanic Males	1,611	53%	1,743	53%	1,872	52%	1,999	52%	2,118	51%
White, not Hispanic Females	176	6%	190	6%	205	6%	216	6%	227	5%
Black/African American, not Hispanic Males	703	23%	779	24%	861	24%	949	25%	1,035	25%
Black/African American, not Hispanic Females	350	12%	379	11%	414	12%	438	11%	468	11%
Hispanic/Latino Males	77	3%	86	3%	95	3%	108	3%	122	3%
Hispanic/Latina Females	23	1%	23	1%	24	1%	25	1%	29	1%
Asian/Pacific Islander Males	11	<1%	13	<1%	14	<1%	14	<1%	15	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	5	<1%	6	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	84	3%	91	3%	99	3%	106	3%	118	3%
Total	3,041		3,311		3,592		3,864		4,142	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Columbus Consortium 2

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,660	68%	1,836	68%	2,004	69%	2,169	69%	2,331	69%
Injection drug use (IDU)	68	3%	69	3%	70	2%	70	2%	74	2%
Male-to-male sexual contact and IDU	65	3%	72	3%	79	3%	87	3%	92	3%
Heterosexual contact	161	7%	187	7%	205	7%	215	7%	219	6%
Other/unknown	504	21%	519	19%	552	19%	603	19%	660	20%
Subtotal	2,458	100%	2,683	100%	2,910	100%	3,144	100%	3,376	100%
Female adult or adolescent										
Injection drug use	54	10%	57	10%	58	9%	61	9%	61	8%
Heterosexual contact	299	54%	325	55%	361	56%	375	55%	391	54%
Other/unknown	199	36%	212	36%	227	35%	248	36%	277	38%
Subtotal	552	100%	594	100%	646	100%	684	100%	729	100%
Child (<13 yrs at diagnosis)										
Perinatal	23	77%	25	76%	27	77%	27	77%	28	78%
Other/unknown	7	23%	8	24%	8	23%	8	23%	8	22%
Subtotal	30	100%	33	100%	35	100%	35	100%	36	100%
Total	3,041		3,311		3,592		3,864		4,142	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Columbus Consortium 2

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	1,587	52%	1,759	53%	1,921	53%	2,081	54%	2,241	54%
Injection drug use (IDU) only	64	2%	64	2%	64	2%	64	2%	67	2%
Heterosexual contact only	460	15%	512	15%	566	16%	590	15%	610	15%
MSM & IDU	55	2%	62	2%	66	2%	74	2%	79	2%
IDU & Heterosexual contact	58	2%	62	2%	64	2%	67	2%	68	2%
MSM & Heterosexual contact	73	2%	77	2%	83	2%	88	2%	90	2%
MSM & IDU & Heterosexual contact	10	<1%	10	<1%	13	<1%	13	<1%	13	<1%
Perinatal exposure	24	1%	27	1%	29	1%	29	1%	30	1%
Other/unknown	710	23%	738	22%	786	22%	858	22%	944	23%
Total	3,041		3,311		3,592		3,864		4,142	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Cincinnati Consortium 3

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Cincinnati Consortium 3

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	16.9	144	77%	96	74%	42	84%	6	75%	
Females	4.9	44	23%	34	26%	8	16%	2	25%	
Age at diagnosis (yr)										
<13	*	1	1%	1	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	11.7	14	7%	13	10%	1	2%	-	-	
20-24	34.5	41	22%	36	28%	5	10%	-	-	
25-29	22.0	25	13%	19	15%	5	10%	1	13%	
30-34	25.9	29	15%	20	15%	8	16%	1	13%	
35-39	11.6	12	6%	6	5%	4	8%	2	25%	
40-44	17.1	20	11%	11	8%	7	14%	2	25%	
45-49	17.7	22	12%	16	12%	4	8%	2	25%	
50-54	8.3	11	6%	3	2%	8	16%	-	-	
55-64	5.0	11	6%	4	3%	7	14%	-	-	
65+	*	2	1%	1	1%	1	2%	-	-	
Race/Ethnicity										
White, not Hispanic	4.6	63	34%	41	32%	18	36%	4	50%	
Black/African American, not Hispanic	45.3	111	59%	80	62%	27	54%	4	50%	
Hispanic/Latino	14.8	7	4%	5	4%	2	4%	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	7	4%	4	3%	3	6%	-	-	
Race/Sex										
White, not Hispanic Males	7.8	53	28%	32	25%	18	36%	3	38%	
White, not Hispanic Females	1.4	10	5%	9	7%	-	-	1	13%	
Black/African American, not Hispanic Males	68.7	79	42%	56	43%	20	40%	3	38%	
Black/African American, not Hispanic Females	24.6	32	17%	24	18%	7	14%	1	13%	
Hispanic/Latino Males	24.2	6	3%	4	3%	2	4%	-	-	
Hispanic/Latina Females	*	1	1%	1	1%	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	7	4%	4	3%	3	6%	-	-	
Total	10.8	188		130		50		8		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	100	70%	69	73%	27	64%	4	67%
Injection drug use (IDU)	1	1%	-	-	1	2%	-	-
Male-to-male sexual contact and IDU	2	1%	2	2%	-	-	-	-
Heterosexual contact	3	2%	3	3%	-	-	-	-
Other/unknown	37	26%	21	22%	14	33%	2	33%
Subtotal	143	100%	95	100%	42	100%	6	100%
Female adult or adolescent								
Injection drug use	2	5%	2	6%	-	-	-	-
Heterosexual contact	7	16%	6	18%	-	-	1	50%
Other/unknown	34	79%	25	76%	8	100%	1	50%
Subtotal	43	100%	33	100%	8	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	2	100%	2	100%	-	-	-	-
Subtotal	2	100%	2	100%	-	-	-	-
Total	188		130		50		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	94	50%	65	50%	25	50%	4	50%
Injection drug use (IDU) only	1	1%	1	1%	-	-	-	-
Heterosexual contact only	10	5%	9	7%	-	-	1	13%
MSM & IDU	1	1%	1	1%	-	-	-	-
IDU & Heterosexual contact	2	1%	1	1%	1	2%	-	-
MSM & Heterosexual contact	6	3%	4	3%	2	4%	-	-
MSM & IDU & Heterosexual contact	1	1%	1	1%	-	-	-	-
Perinatal exposure	1	1%	1	1%	-	-	-	-
Other/unknown	72	38%	47	36%	22	44%	3	38%
Total	188		130		50		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	39	74%	51	65%	5	83%	-	-	-	-	5	83%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	1	17%
Male-to-male sexual contact and IDU	1	2%	1	1%	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	3	4%	-	-	-	-	-	-	-	-
Other/unknown	13	25%	23	29%	1	17%	-	-	-	-	-	-
Subtotal	53	100%	78	100%	6	100%	-	-	-	-	6	100%
Female adult or adolescent												
Injection drug use	2	20%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	7	23%	-	-	-	-	-	-	-	-
Other/unknown	8	80%	24	77%	1	100%	-	-	-	-	1	100%
Subtotal	10	100%	31	100%	1	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	2	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	-	-	-	-	-	-	-	-
Total	63		111		7		0		0		7	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	36	57%	48	43%	5	71%	-	-	-	-	5	71%
Injection drug use (IDU) only	1	2%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	10	9%	-	-	-	-	-	-	-	-
MSM & IDU	1	2%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	2%	-	-	-	-	-	-	-	-	1	14%
MSM & Heterosexual contact	3	5%	3	3%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	1	1%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	1%	-	-	-	-	-	-	-	-
Other/unknown	21	33%	48	43%	2	29%	-	-	-	-	1	14%
Total	63		111		7		0		0		7	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Cincinnati Consortium 3**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	14.8	126	76%	17.0	145	74%	16.0	136	75%	15.4	131	72%
Females	4.4	39	24%	5.6	50	26%	5.0	45	25%	5.7	51	28%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	7.5	9	5%	14.2	17	9%	15.0	18	10%	9.2	11	6%
20-24	33.6	40	24%	37.0	44	23%	32.8	39	22%	35.3	42	23%
25-29	15.9	18	11%	25.6	29	15%	28.2	32	18%	24.7	28	15%
30-34	12.5	14	8%	22.3	25	13%	15.2	17	9%	25.0	28	15%
35-39	19.3	20	12%	13.5	14	7%	19.3	20	11%	13.5	14	8%
40-44	25.6	30	18%	12.0	14	7%	17.1	20	11%	14.5	17	9%
45-49	11.3	14	8%	15.3	19	10%	18.5	23	13%	12.9	16	9%
50-54	9.8	13	8%	13.6	18	9%	6.8	9	5%	11.3	15	8%
55-64	3.2	7	4%	6.3	14	7%	*	1	1%	3.6	8	4%
65+	*	-	-	*	1	1%	*	2	1%	*	3	2%
Race/Ethnicity												
White, not Hispanic	4.2	58	35%	4.2	58	30%	4.4	61	34%	4.9	68	37%
Black/African American, not Hispanic	38.7	95	58%	51.4	126	65%	45.7	112	62%	43.2	106	58%
Hispanic/Latino	23.2	11	7%	12.7	6	3%	*	2	1%	10.5	5	3%
Asian/Pacific Islander	*	-	-	*	1	1%	*	1	1%	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	4	2%	*	5	3%	*	1	1%
Race/Sex												
White, not Hispanic Males	7.2	49	30%	6.8	46	24%	6.8	46	25%	8.3	56	31%
White, not Hispanic Females	1.3	9	5%	1.7	12	6%	2.1	15	8%	1.7	12	7%
Black/African American, not Hispanic Males	57.4	66	40%	77.4	89	46%	73.9	85	47%	60.9	70	38%
Black/African American, not Hispanic Females	22.3	29	18%	28.4	37	19%	20.7	27	15%	27.6	36	20%
Hispanic/Latino Males	40.3	10	6%	20.1	5	3%	*	2	1%	*	4	2%
Hispanic/Latina Females	*	1	1%	*	1	1%	*	-	-	*	1	1%
Asian/Pacific Islander Males	*	-	-	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	1	1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	1%	*	4	2%	*	5	3%	*	1	1%
Total	9.4	165		11.2	195		10.4	181		10.4	182	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Cincinnati Consortium 3

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	81	64%	104	72%	98	72%	78	60%
Injection drug use (IDU)	2	2%	2	1%	-	-	1	1%
Male-to-male sexual contact and IDU	5	4%	3	2%	3	2%	2	2%
Heterosexual contact	11	9%	9	6%	10	7%	3	2%
Other/unknown	27	21%	27	19%	25	18%	47	36%
Subtotal	126	100%	145	100%	136	100%	131	100%
Female adult or adolescent								
Injection drug use	4	10%	7	14%	4	9%	3	6%
Heterosexual contact	20	51%	17	34%	14	31%	13	25%
Other/unknown	15	38%	26	52%	27	60%	35	69%
Subtotal	39	100%	50	100%	45	100%	51	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	165		195		181		182	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Cincinnati Consortium 3

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	81	49%	96	49%	95	52%	74	41%
Injection drug use (IDU) only	6	4%	6	3%	2	1%	2	1%
Heterosexual contact only	31	19%	26	13%	24	13%	16	9%
MSM & IDU	3	2%	3	2%	3	2%	2	1%
IDU & Heterosexual contact	-	-	3	2%	2	1%	2	1%
MSM & Heterosexual contact	-	-	8	4%	3	2%	4	2%
MSM & IDU & Heterosexual contact	2	1%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	41	25%	53	27%	52	29%	82	45%
Total	165		195		181		182	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Cincinnati Consortium 3

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	62	77%	73	73%	72	71%	87	70%	96	74%
Females	19	23%	27	27%	29	29%	37	30%	34	26%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	5	6%	12	12%	16	16%	10	8%	13	10%
20-24	28	35%	30	30%	29	29%	29	23%	36	28%
25-29	7	9%	13	13%	15	15%	24	19%	19	15%
30-34	7	9%	16	16%	10	10%	14	11%	20	15%
35-39	6	7%	3	3%	10	10%	8	6%	6	5%
40-44	17	21%	5	5%	8	8%	11	9%	11	8%
45-49	6	7%	8	8%	10	10%	11	9%	16	12%
50-54	3	4%	7	7%	3	3%	12	10%	3	2%
55-64	2	2%	6	6%	-	-	4	3%	4	3%
65+	-	-	-	-	-	-	1	1%	1	1%
Race/Ethnicity										
White, not Hispanic	23	28%	30	30%	38	38%	44	35%	41	32%
Black/African American, not Hispanic	51	63%	66	66%	58	57%	73	59%	80	62%
Hispanic/Latino	7	9%	2	2%	1	1%	4	3%	5	4%
Asian/Pacific Islander	-	-	-	-	1	1%	2	2%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	2%	3	3%	1	1%	4	3%
Race/Sex										
White, not Hispanic Males	18	22%	22	22%	28	28%	35	28%	32	25%
White, not Hispanic Females	5	6%	8	8%	10	10%	9	7%	9	7%
Black/African American, not Hispanic Males	38	47%	47	47%	41	41%	48	39%	56	43%
Black/African American, not Hispanic Females	13	16%	19	19%	17	17%	25	20%	24	18%
Hispanic/Latino Males	6	7%	2	2%	1	1%	3	2%	4	3%
Hispanic/Latina Females	1	1%	-	-	-	-	1	1%	1	1%
Asian/Pacific Islander Males	-	-	-	-	1	1%	1	1%	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	1	1%	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	2%	3	3%	1	1%	4	3%
Total	81		100		101		124		130	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	37	60%	57	78%	54	75%	50	57%	69	73%
Injection drug use (IDU)	2	3%	-	-	-	-	1	1%	-	-
Male-to-male sexual contact and IDU	2	3%	3	4%	2	3%	2	2%	2	2%
Heterosexual contact	7	11%	1	1%	7	10%	3	3%	3	3%
Other/unknown	14	23%	12	16%	9	13%	31	36%	21	22%
Subtotal	62	100%	73	100%	72	100%	87	100%	95	100%
Female adult or adolescent										
Injection drug use	-	-	3	11%	4	14%	2	5%	2	6%
Heterosexual contact	9	47%	8	30%	9	31%	11	30%	6	18%
Other/unknown	10	53%	16	59%	16	55%	24	65%	25	76%
Subtotal	19	100%	27	100%	29	100%	37	100%	33	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	100%
Subtotal	-	-	-	-	-	-	-	-	2	100%
Total	81		100		101		124		130	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	37	46%	52	52%	53	52%	46	37%	65	50%
Injection drug use (IDU) only	2	2%	3	3%	2	2%	2	2%	1	1%
Heterosexual contact only	16	20%	9	9%	16	16%	14	11%	9	7%
MSM & IDU	1	1%	3	3%	2	2%	2	2%	1	1%
IDU & Heterosexual contact	-	-	-	-	2	2%	1	1%	1	1%
MSM & Heterosexual contact	-	-	5	5%	1	1%	4	3%	4	3%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-	1	1%
Perinatal exposure	-	-	-	-	-	-	-	-	1	1%
Other/unknown	24	30%	28	28%	25	25%	55	44%	47	36%
Total	81		100		101		124		130	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Cincinnati Consortium 3

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	86	77%	93	73%	85	80%	76	74%	63	83%	2,500	86%
Females	26	23%	35	27%	21	20%	27	26%	13	17%	397	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	1%	-	-	23	1%
13-14	-	-	-	-	-	-	-	-	-	-	3	<1%
15-19	5	4%	9	7%	4	4%	6	6%	1	1%	61	2%
20-24	16	14%	20	16%	14	13%	24	23%	10	13%	264	9%
25-29	18	16%	20	16%	23	22%	10	10%	7	9%	496	17%
30-34	13	12%	13	10%	15	14%	17	17%	12	16%	616	21%
35-39	19	17%	20	16%	8	8%	11	11%	10	13%	531	18%
40-44	17	15%	15	12%	12	11%	9	9%	13	17%	393	14%
45-49	10	9%	15	12%	15	14%	9	9%	6	8%	227	8%
50-54	8	7%	10	8%	8	8%	7	7%	8	11%	149	5%
55-64	6	5%	5	4%	4	4%	7	7%	8	11%	101	3%
65+	-	-	1	1%	3	3%	2	2%	1	1%	33	1%
Race/Ethnicity												
White, not Hispanic	44	39%	41	32%	38	36%	38	37%	27	36%	1,522	53%
Black/African American, not Hispanic	62	55%	77	60%	66	62%	60	58%	42	55%	1,297	45%
Hispanic/Latino	4	4%	5	4%	1	1%	2	2%	3	4%	44	2%
Asian/Pacific Islander	-	-	1	1%	-	-	-	-	-	-	7	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	2	2%	4	3%	1	1%	3	3%	4	5%	25	1%
Race/Sex												
White, not Hispanic Males	39	35%	36	28%	33	31%	29	28%	26	34%	1,411	49%
White, not Hispanic Females	5	4%	5	4%	5	5%	9	9%	1	1%	111	4%
Black/African American, not Hispanic Males	41	37%	51	40%	50	47%	43	42%	31	41%	1,023	35%
Black/African American, not Hispanic Females	21	19%	26	20%	16	15%	17	17%	11	14%	274	9%
Hispanic/Latino Males	4	4%	3	2%	1	1%	2	2%	3	4%	38	1%
Hispanic/Latina Females	-	-	2	2%	-	-	-	-	-	-	6	<1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	-	-	-	-	6	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	4	3%	1	1%	3	3%	4	5%	25	1%
Total	112		128		106		103		76		2,897	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	57	66%	59	63%	63	74%	50	66%	39	62%	1,791	72%
Injection drug use (IDU)	1	1%	2	2%	-	-	-	-	2	3%	107	4%
Male-to-male sexual contact and IDU	1	1%	3	3%	4	5%	1	1%	1	2%	140	6%
Heterosexual contact	12	14%	12	13%	3	4%	2	3%	1	2%	111	4%
Other/unknown	15	17%	17	18%	15	18%	23	30%	20	32%	336	14%
Subtotal	86	100%	93	100%	85	100%	76	100%	63	100%	2,485	100%
Female adult or adolescent												
Injection drug use	1	4%	5	14%	1	5%	3	12%	1	8%	57	15%
Heterosexual contact	20	77%	18	51%	5	24%	8	31%	2	15%	205	53%
Other/unknown	5	19%	12	34%	15	71%	15	58%	10	77%	127	33%
Subtotal	26	100%	35	100%	21	100%	26	100%	13	100%	389	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	1	100%	-	-	17	74%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	26%
Subtotal	-	-	-	-	-	-	1	100%	-	-	23	100%
Total	112		128		106		103		76		2,897	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	55	49%	54	42%	62	58%	50	49%	36	47%	1,714	59%
Injection drug use (IDU) only	2	2%	4	3%	-	-	2	2%	1	1%	98	3%
Heterosexual contact only	32	29%	30	23%	8	8%	10	10%	3	4%	316	11%
MSM & IDU	1	1%	3	2%	2	2%	1	1%	1	1%	122	4%
IDU & Heterosexual contact	-	-	3	2%	1	1%	1	1%	2	3%	66	2%
MSM & Heterosexual contact	2	2%	5	4%	1	1%	-	-	3	4%	75	3%
MSM & IDU & Heterosexual contact	-	-	-	-	2	2%	-	-	-	-	18	1%
Perinatal exposure	-	-	-	-	-	-	1	1%	-	-	17	1%
Other/unknown	20	18%	29	23%	30	28%	38	37%	30	39%	471	16%
Total	112		128		106		103		76		2,897	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Cincinnati Consortium 3

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Cincinnati Consortium 3

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	260.7	2,221	79%	1,039	76%	1,182	81%
Females	66.2	592	21%	320	24%	272	19%
Age at end of year							
<13	5.9	18	1%	9	1%	9	1%
13-14	*	3	<1%	-	-	3	<1%
15-19	120.2	144	5%	97	7%	47	3%
20-24	347.2	413	15%	244	18%	169	12%
25-29	435.6	494	18%	236	17%	258	18%
30-34	468.9	525	19%	244	18%	281	19%
35-39	435.3	450	16%	193	14%	257	18%
40-44	289.6	339	12%	143	11%	196	13%
45-49	162.7	202	7%	86	6%	116	8%
50-54	103.4	137	5%	68	5%	69	5%
55-64	32.5	72	3%	31	2%	41	3%
65+	6.9	16	1%	8	1%	8	1%
Race/Ethnicity							
White, not Hispanic	91.3	1,261	45%	617	45%	644	44%
Black/African American, not Hispanic	585.5	1,436	51%	684	50%	752	52%
Hispanic/Latino	126.5	60	2%	28	2%	32	2%
Asian/Pacific Islander	32.8	13	<1%	9	1%	4	<1%
American Indian/Alaska Native	*	2	<1%	-	-	2	<1%
Unknown	*	41	1%	21	2%	20	1%
Race/Sex							
White, not Hispanic Males	158.2	1,071	38%	499	37%	572	39%
White, not Hispanic Females	27.0	190	7%	118	9%	72	5%
Black/African American, not Hispanic Males	917.7	1,055	38%	495	36%	560	39%
Black/African American, not Hispanic Females	292.4	381	14%	189	14%	192	13%
Hispanic/Latino Males	197.3	49	2%	21	2%	28	2%
Hispanic/Latina Females	48.7	11	<1%	7	1%	4	<1%
Asian/Pacific Islander Males	57.5	11	<1%	8	1%	3	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	-	-	2	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	41	1%	21	2%	20	1%
Total	161.1	2,813		1,359		1,454	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,494	68%	663	64%	831	71%
Injection drug use (IDU)	71	3%	30	3%	41	3%
Male-to-male sexual contact and IDU	88	4%	31	3%	57	5%
Heterosexual contact	139	6%	57	6%	82	7%
Other/unknown	420	19%	253	24%	167	14%
Subtotal	2,212	100%	1,034	100%	1,178	100%
Female adult or adolescent						
Injection drug use	52	9%	22	7%	30	11%
Heterosexual contact	315	54%	159	50%	156	58%
Other/unknown	216	37%	135	43%	81	30%
Subtotal	583	100%	316	100%	267	100%
Child (<13 yrs at diagnosis)						
Perinatal	14	78%	8	89%	6	67%
Other/unknown	4	22%	1	11%	3	33%
Subtotal	18	100%	9	100%	9	100%
Total	2,813		1,359		1,454	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	1,424	51%	643	47%	781	54%
Injection drug use (IDU) only	72	3%	33	2%	39	3%
Heterosexual contact only	454	16%	216	16%	238	16%
MSM & IDU	73	3%	26	2%	47	3%
IDU & Heterosexual contact	51	2%	19	1%	32	2%
MSM & Heterosexual contact	70	2%	20	1%	50	3%
MSM & IDU & Heterosexual contact	15	1%	5	<1%	10	1%
Perinatal exposure	14	<1%	8	1%	6	<1%
Other/unknown	640	23%	389	29%	251	17%
Total	2,813		1,359		1,454	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Cincinnati Consortium 3

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	819	77%	626	60%	21	43%	8	73%	1	50%	19	58%
Injection drug use (IDU)	15	1%	46	4%	10	20%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	42	4%	43	4%	1	2%	-	-	1	50%	1	3%
Heterosexual contact	27	3%	101	10%	8	16%	2	18%	-	-	1	3%
Other/unknown	164	15%	234	22%	9	18%	1	9%	-	-	12	36%
Subtotal	1,067	100%	1,050	100%	49	100%	11	100%	2	100%	33	100%
Female adult or adolescent												
Injection drug use	25	13%	26	7%	-	-	-	-	-	-	1	13%
Heterosexual contact	104	55%	198	53%	8	73%	2	100%	-	-	3	38%
Other/unknown	59	31%	150	40%	3	27%	-	-	-	-	4	50%
Subtotal	188	100%	374	100%	11	100%	2	100%	-	-	8	100%
Child (<13 yrs at diagnosis)												
Perinatal	4	67%	10	83%	-	-	-	-	-	-	-	-
Other/unknown	2	33%	2	17%	-	-	-	-	-	-	-	-
Subtotal	6	100%	12	100%	-	-	-	-	-	-	-	-
Total	1,261		1,436		60		13		2		41	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	795	63%	583	41%	18	30%	8	62%	1	50%	19	46%
Injection drug use (IDU) only	27	2%	39	3%	6	10%	-	-	-	-	-	-
Heterosexual contact only	131	10%	299	21%	16	27%	4	31%	-	-	4	10%
MSM & IDU	36	3%	36	3%	-	-	-	-	1	50%	-	-
IDU & Heterosexual contact	13	1%	33	2%	4	7%	-	-	-	-	1	2%
MSM & Heterosexual contact	24	2%	43	3%	3	5%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	6	0%	7	0%	1	2%	-	-	-	-	1	2%
Perinatal exposure	4	0%	10	1%	-	-	-	-	-	-	-	-
Other/unknown	225	18%	386	27%	12	20%	1	8%	-	-	16	39%
Total	1,261		1,436		60		13		2		41	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Cincinnati Consortium 3

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	1,706	80%	1,821	80%	1,958	80%	2,092	79%	2,221	79%
Females	419	20%	455	20%	501	20%	543	21%	592	21%
Age at end of year										
<13	18	1%	18	1%	18	1%	18	1%	18	1%
13-14	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
15-19	88	4%	98	4%	115	5%	133	5%	144	5%
20-24	252	12%	291	13%	334	14%	372	14%	413	15%
25-29	393	18%	409	18%	435	18%	466	18%	494	18%
30-34	444	21%	458	20%	481	20%	497	19%	525	19%
35-39	384	18%	402	18%	416	17%	436	17%	450	16%
40-44	263	12%	289	13%	302	12%	322	12%	339	12%
45-49	136	6%	149	7%	167	7%	189	7%	202	7%
50-54	86	4%	97	4%	113	5%	122	5%	137	5%
55-64	47	2%	51	2%	63	3%	64	2%	72	3%
65+	11	1%	11	<1%	12	<1%	13	<1%	16	1%
Race/Ethnicity										
White, not Hispanic	1,028	48%	1,082	48%	1,136	46%	1,195	45%	1,261	45%
Black/African American, not Hispanic	1,017	48%	1,103	48%	1,222	50%	1,332	51%	1,436	51%
Hispanic/Latino	38	2%	48	2%	53	2%	55	2%	60	2%
Asian/Pacific Islander	9	<1%	9	<1%	10	<1%	11	<1%	13	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	31	1%	32	1%	36	1%	40	2%	41	1%
Race/Sex										
White, not Hispanic Males	883	42%	929	41%	972	40%	1,017	39%	1,071	38%
White, not Hispanic Females	145	7%	153	7%	164	7%	178	7%	190	7%
Black/African American, not Hispanic Males	757	36%	816	36%	901	37%	985	37%	1,055	38%
Black/African American, not Hispanic Females	260	12%	287	13%	321	13%	347	13%	381	14%
Hispanic/Latino Males	30	1%	39	2%	43	2%	45	2%	49	2%
Hispanic/Latina Females	8	<1%	9	<1%	10	<1%	10	<1%	11	<1%
Asian/Pacific Islander Males	8	<1%	8	<1%	9	<1%	10	<1%	11	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	31	1%	32	1%	36	1%	40	2%	41	1%
Total	2,125		2,276		2,459		2,635		2,813	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Cincinnati Consortium 3

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,144	67%	1,221	67%	1,321	68%	1,417	68%	1,494	68%
Injection drug use (IDU)	66	4%	68	4%	70	4%	70	3%	71	3%
Male-to-male sexual contact and IDU	75	4%	80	4%	83	4%	86	4%	88	4%
Heterosexual contact	106	6%	117	6%	126	6%	136	7%	139	6%
Other/unknown	306	18%	326	18%	349	18%	374	18%	420	19%
Subtotal	1,697	100%	1,812	100%	1,949	100%	2,083	100%	2,212	100%
Female adult or adolescent										
Injection drug use	35	9%	39	9%	45	9%	49	9%	52	9%
Heterosexual contact	254	62%	273	61%	289	59%	303	57%	315	54%
Other/unknown	121	30%	134	30%	158	32%	182	34%	216	37%
Subtotal	410	100%	446	100%	492	100%	534	100%	583	100%
Child (<13 yrs at diagnosis)										
Perinatal	14	78%	14	78%	14	78%	14	78%	14	78%
Other/unknown	4	22%	4	22%	4	22%	4	22%	4	22%
Subtotal	18	100%	18	100%	18	100%	18	100%	18	100%
Total	2,125		2,276		2,459		2,635		2,813	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Cincinnati Consortium 3

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	1,089	51%	1,166	51%	1,258	51%	1,351	51%	1,424	51%
Injection drug use (IDU) only	57	3%	63	3%	68	3%	70	3%	72	3%
Heterosexual contact only	360	17%	390	17%	415	17%	439	17%	454	16%
MSM & IDU	62	3%	65	3%	68	3%	71	3%	73	3%
IDU & Heterosexual contact	44	2%	44	2%	47	2%	49	2%	51	2%
MSM & Heterosexual contact	55	3%	55	2%	63	3%	66	3%	70	2%
MSM & IDU & Heterosexual contact	13	1%	15	1%	15	1%	15	1%	15	1%
Perinatal exposure	14	1%	14	1%	14	1%	14	1%	14	<1%
Other/unknown	431	20%	464	20%	511	21%	560	21%	640	23%
Total	2,125		2,276		2,459		2,635		2,813	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B Dayton
Consortium 4

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Dayton Consortium 4

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	13.4	67	80%	42	75%	20	91%	5	83%
Females	3.2	17	20%	14	25%	2	9%	1	17%
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	3	4%	2	4%	1	5%	-	-
20-24	25.8	19	23%	16	29%	3	14%	-	-
25-29	25.1	16	19%	13	23%	2	9%	1	17%
30-34	9.7	6	7%	3	5%	1	5%	2	33%
35-39	21.4	12	14%	9	16%	3	14%	-	-
40-44	9.3	6	7%	3	5%	2	9%	1	17%
45-49	11.5	8	10%	1	2%	6	27%	1	17%
50-54	11.7	9	11%	6	11%	2	9%	1	17%
55-64	3.6	5	6%	3	5%	2	9%	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	4.3	36	43%	23	41%	11	50%	2	33%
Black/African American, not Hispanic	30.4	42	50%	29	52%	10	45%	3	50%
Hispanic/Latino	*	3	4%	2	4%	1	5%	-	-
Asian/Pacific Islander	*	2	2%	1	2%	-	-	1	17%
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	1%	1	2%	-	-	-	-
Race/Sex									
White, not Hispanic Males	7.4	30	36%	18	32%	10	45%	2	33%
White, not Hispanic Females	1.4	6	7%	5	9%	1	5%	-	-
Black/African American, not Hispanic Males	49.3	32	38%	21	38%	9	41%	2	33%
Black/African American, not Hispanic Females	13.7	10	12%	8	14%	1	5%	1	17%
Hispanic/Latino Males	*	3	4%	2	4%	1	5%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	-	-	-	-	1	17%
Asian/Pacific Islander Females	*	1	1%	1	2%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	1%	1	2%	-	-	-	-
Total	8.1	84		56		22		6	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Dayton Consortium 4

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	50	75%	33	79%	14	70%	3	60%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	3%	-	-	2	10%	-	-
Heterosexual contact	5	7%	3	7%	2	10%	-	-
Other/unknown	10	15%	6	14%	2	10%	2	40%
Subtotal	67	100%	42	100%	20	100%	5	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	6	35%	5	36%	1	50%	-	-
Other/unknown	11	65%	9	64%	1	50%	1	100%
Subtotal	17	100%	14	100%	2	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	84		56		22		6	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Dayton Consortium 4

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	48	57%	32	57%	13	59%	3	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	11	13%	8	14%	3	14%	-	-
MSM & IDU	1	1%	-	-	1	5%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	2	2%	1	2%	1	5%	-	-
MSM & IDU & Heterosexual contact	1	1%	-	-	1	5%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	21	25%	15	27%	3	14%	3	50%
Total	84		56		22		6	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B
Dayton Consortium 4**

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	22	73%	25	78%	2	67%	-	-	-	-	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	7%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	3%	3	9%	1	33%	-	-	-	-	-	-
Other/unknown	5	17%	4	13%	-	-	1	100%	-	-	-	-
Subtotal	30	100%	32	100%	3	100%	1	100%	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	4	67%	2	20%	-	-	-	-	-	-	-	-
Other/unknown	2	33%	8	80%	-	-	1	100%	-	-	-	-
Subtotal	6	100%	10	100%	-	-	1	100%	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	36		42		3		2		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Dayton Consortium 4

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	22	61%	23	55%	2	67%	-	-	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	5	14%	5	12%	1	33%	-	-	-	-	-	-
MSM & IDU	1	3%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	2	5%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	7	19%	12	29%	-	-	2	100%	-	-	-	-
Total	36		42		3		2		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Dayton Consortium 4**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex												
Males	12.2	61	80%	11.8	59	78%	16.4	82	85%	15.6	78	77%
Females	2.8	15	20%	3.2	17	22%	2.8	15	15%	4.3	23	23%
Age at diagnosis (yr)												
<13	*	1	1%	*	1	1%	*	4	4%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	8.5	6	8%	9.9	7	9%	19.9	14	14%	9.9	7	7%
20-24	10.8	8	11%	21.7	16	21%	32.5	24	25%	38.0	28	28%
25-29	17.2	11	14%	15.7	10	13%	25.1	16	16%	15.7	10	10%
30-34	16.1	10	13%	14.5	9	12%	11.3	7	7%	19.3	12	12%
35-39	16.0	9	12%	*	3	4%	10.7	6	6%	23.2	13	13%
40-44	18.7	12	16%	9.3	6	8%	14.0	9	9%	12.5	8	8%
45-49	10.1	7	9%	14.4	10	13%	8.6	6	6%	15.8	11	11%
50-54	9.1	7	9%	7.8	6	8%	7.8	6	6%	10.4	8	8%
55-64	*	4	5%	5.8	8	11%	*	2	2%	*	2	2%
65+	*	1	1%	*	-	-	*	3	3%	*	2	2%
Race/Ethnicity												
White, not Hispanic	4.1	34	45%	3.1	26	34%	4.2	35	36%	4.7	39	39%
Black/African American, not Hispanic	26.8	37	49%	35.5	49	64%	42.7	59	61%	41.3	57	56%
Hispanic/Latino	*	2	3%	*	-	-	*	-	-	*	3	3%
Asian/Pacific Islander	*	1	1%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	3%	*	1	1%	*	3	3%	*	2	2%
Race/Sex												
White, not Hispanic Males	7.2	29	38%	5.7	23	30%	8.4	34	35%	7.7	31	31%
White, not Hispanic Females	1.2	5	7%	*	3	4%	*	1	1%	1.9	8	8%
Black/African American, not Hispanic Males	43.1	28	37%	53.9	35	46%	70.9	46	47%	64.7	42	42%
Black/African American, not Hispanic Females	12.3	9	12%	19.1	14	18%	17.7	13	13%	20.5	15	15%
Hispanic/Latino Males	*	2	3%	*	-	-	*	-	-	*	3	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	3%	*	1	1%	*	3	3%	*	2	2%
Total	7.4	76		7.4	76		9.4	97		9.8	101	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Dayton Consortium 4

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	47	78%	36	62%	66	84%	47	60%
Injection drug use (IDU)	1	2%	2	3%	1	1%	1	1%
Male-to-male sexual contact and IDU	-	-	1	2%	1	1%	-	-
Heterosexual contact	2	3%	3	5%	3	4%	8	10%
Other/unknown	10	17%	16	28%	8	10%	22	28%
Subtotal	60	100%	58	100%	79	100%	78	100%
Female adult or adolescent								
Injection drug use	1	7%	1	6%	-	-	3	13%
Heterosexual contact	9	64%	7	41%	5	36%	9	39%
Other/unknown	4	29%	9	53%	9	64%	11	48%
Subtotal	14	100%	17	100%	14	100%	23	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	1	100%	3	75%	-	-
Other/unknown	2	100%	-	-	1	25%	-	-
Subtotal	2	100%	1	100%	4	100%	-	-
Total	76		76		97		101	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Dayton Consortium 4

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	45	59%	36	47%	66	68%	44	44%
Injection drug use (IDU) only	1	1%	2	3%	1	1%	1	1%
Heterosexual contact only	11	14%	10	13%	8	8%	17	17%
MSM & IDU	-	-	1	1%	-	-	-	-
IDU & Heterosexual contact	1	1%	1	1%	-	-	3	3%
MSM & Heterosexual contact	2	3%	-	-	-	-	3	3%
MSM & IDU & Heterosexual contact	-	-	-	-	1	1%	-	-
Perinatal exposure	1	1%	1	1%	3	3%	-	-
Other/unknown	15	20%	25	33%	18	19%	33	33%
Total	76		76		97		101	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Dayton Consortium 4

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	33	79%	39	75%	55	86%	56	76%	42	75%
Females	9	21%	13	25%	9	14%	18	24%	14	25%
Age at HIV diagnosis (yr)										
<13	-	-	1	2%	4	6%	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	4	10%	4	8%	11	17%	6	8%	2	4%
20-24	4	10%	13	25%	20	31%	23	31%	16	29%
25-29	6	14%	7	13%	11	17%	7	9%	13	23%
30-34	4	10%	7	13%	2	3%	10	14%	3	5%
35-39	5	12%	2	4%	4	6%	8	11%	9	16%
40-44	6	14%	4	8%	5	8%	5	7%	3	5%
45-49	6	14%	6	12%	1	2%	6	8%	1	2%
50-54	3	7%	2	4%	3	5%	5	7%	6	11%
55-64	3	7%	6	12%	1	2%	2	3%	3	5%
65+	1	2%	-	-	2	3%	2	3%	-	-
Race/Ethnicity										
White, not Hispanic	17	40%	16	31%	20	31%	28	38%	23	41%
Black/African American, not Hispanic	22	52%	36	69%	42	66%	43	58%	29	52%
Hispanic/Latino	1	2%	-	-	-	-	1	1%	2	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	2%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	5%	-	-	2	3%	2	3%	1	2%
Race/Sex										
White, not Hispanic Males	15	36%	14	27%	20	31%	21	28%	18	32%
White, not Hispanic Females	2	5%	2	4%	-	-	7	9%	5	9%
Black/African American, not Hispanic Males	16	38%	25	48%	34	53%	32	43%	21	38%
Black/African American, not Hispanic Females	6	14%	11	21%	8	13%	11	15%	8	14%
Hispanic/Latino Males	1	2%	-	-	-	-	1	1%	2	4%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	2%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	5%	-	-	2	3%	2	3%	1	2%
Total	42		52		64		74		56	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Dayton Consortium 4

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	23	70%	23	61%	42	81%	33	59%	33	79%
Injection drug use (IDU)	1	3%	1	3%	-	-	1	2%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	-	-	-	-
Heterosexual contact	1	3%	3	8%	2	4%	3	5%	3	7%
Other/unknown	8	24%	11	29%	7	13%	19	34%	6	14%
Subtotal	33	100%	38	100%	52	100%	56	100%	42	100%
Female adult or adolescent										
Injection drug use	1	13%	1	8%	-	-	3	17%	-	-
Heterosexual contact	4	50%	5	38%	3	38%	6	33%	5	36%
Other/unknown	3	38%	7	54%	5	63%	9	50%	9	64%
Subtotal	8	100%	13	100%	8	100%	18	100%	14	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	1	100%	3	75%	-	-	-	-
Other/unknown	1	100%	-	-	1	25%	-	-	-	-
Subtotal	1	100%	1	100%	4	100%	-	-	-	-
Total	42		52		64		74		56	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Dayton Consortium 4

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	22	52%	23	44%	42	66%	30	41%	32	57%
Injection drug use (IDU) only	1	2%	2	4%	-	-	1	1%	-	-
Heterosexual contact only	5	12%	8	15%	5	8%	9	12%	8	14%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	2%	-	-	-	-	3	4%	-	-
MSM & Heterosexual contact	1	2%	-	-	-	-	3	4%	1	2%
MSM & IDU & Heterosexual contact	-	-	-	-	1	2%	-	-	-	-
Perinatal exposure	1	2%	1	2%	3	5%	-	-	-	-
Other/unknown	11	26%	18	35%	13	20%	28	38%	15	27%
Total	42		52		64		74		56	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Dayton Consortium 4

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	44	80%	34	79%	32	74%	33	85%	46	90%	1,415	85%
Females	11	20%	9	21%	11	26%	6	15%	5	10%	251	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	1	2%	-	-	-	-	23	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	2	4%	3	7%	4	9%	3	8%	4	8%	43	3%
20-24	8	15%	8	19%	4	9%	5	13%	5	10%	140	8%
25-29	8	15%	6	14%	5	12%	5	13%	9	18%	283	17%
30-34	7	13%	8	19%	10	23%	8	21%	6	12%	317	19%
35-39	10	18%	2	5%	4	9%	4	10%	7	14%	293	18%
40-44	10	18%	4	9%	5	12%	4	10%	5	10%	248	15%
45-49	3	5%	6	14%	5	12%	7	18%	8	16%	162	10%
50-54	6	11%	4	9%	3	7%	3	8%	5	10%	77	5%
55-64	1	2%	2	5%	1	2%	-	-	2	4%	54	3%
65+	-	-	-	-	1	2%	-	-	-	-	24	1%
Race/Ethnicity												
White, not Hispanic	25	45%	17	40%	18	42%	20	51%	23	45%	1,017	61%
Black/African American, not Hispanic	28	51%	25	58%	24	56%	16	41%	24	47%	613	37%
Hispanic/Latino	-	-	-	-	-	-	1	3%	3	6%	19	1%
Asian/Pacific Islander	2	4%	-	-	-	-	-	-	1	2%	6	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	1	2%	1	2%	2	5%	-	-	10	1%
Race/Sex												
White, not Hispanic Males	23	42%	14	33%	16	37%	18	46%	21	41%	916	55%
White, not Hispanic Females	2	4%	3	7%	2	5%	2	5%	2	4%	101	6%
Black/African American, not Hispanic Males	19	35%	19	44%	15	35%	13	33%	21	41%	467	28%
Black/African American, not Hispanic Females	9	16%	6	14%	9	21%	3	8%	3	6%	146	9%
Hispanic/Latino Males	-	-	-	-	-	-	1	3%	3	6%	18	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	2	4%	-	-	-	-	-	-	1	2%	5	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	2%	1	2%	2	5%	-	-	10	1%
Total	55		43		43		39		51		1,666	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Dayton Consortium 4

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	37	84%	25	74%	27	87%	21	64%	33	72%	1,016	73%
Injection drug use (IDU)	1	2%	1	3%	1	3%	1	3%	-	-	60	4%
Male-to-male sexual contact and IDU	-	-	-	-	1	3%	1	3%	2	4%	89	6%
Heterosexual contact	4	9%	2	6%	1	3%	5	15%	4	9%	85	6%
Other/unknown	2	5%	6	18%	1	3%	5	15%	7	15%	150	11%
Subtotal	44	100%	34	100%	31	100%	33	100%	46	100%	1,400	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	41	17%
Heterosexual contact	10	91%	6	67%	6	55%	4	67%	2	40%	141	58%
Other/unknown	1	9%	3	33%	5	45%	2	33%	3	60%	61	25%
Subtotal	11	100%	9	100%	11	100%	6	100%	5	100%	243	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	15	65%
Other/unknown	-	-	-	-	1	100%	-	-	-	-	8	35%
Subtotal	-	-	-	-	1	100%	-	-	-	-	23	100%
Total	55		43		43		39		51		1,666	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Dayton Consortium 4

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	36	65%	23	53%	27	63%	21	54%	32	63%	975	59%
Injection drug use (IDU) only	1	2%	-	-	1	2%	1	3%	-	-	64	4%
Heterosexual contact only	14	25%	8	19%	7	16%	9	23%	6	12%	226	14%
MSM & IDU	-	-	-	-	1	2%	1	3%	1	2%	75	5%
IDU & Heterosexual contact	-	-	1	2%	-	-	-	-	-	-	36	2%
MSM & Heterosexual contact	1	2%	2	5%	-	-	-	-	1	2%	41	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	2%	14	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	15	1%
Other/unknown	3	5%	9	21%	7	16%	7	18%	10	20%	220	13%
Total	55		43		43		39		51		1,666	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Dayton Consortium 4

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Dayton Consortium 4

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	239.3	1,199	79%	549	76%	650	81%
Females	61.7	328	21%	174	24%	154	19%
Age at end of year							
<13	10.2	17	1%	10	1%	7	1%
13-14	18.9	5	<1%	3	<1%	2	<1%
15-19	129.2	91	6%	53	7%	38	5%
20-24	298.2	220	14%	127	18%	93	12%
25-29	410.5	262	17%	127	18%	135	17%
30-34	432.9	269	18%	109	15%	160	20%
35-39	392.3	220	14%	94	13%	126	16%
40-44	310.0	199	13%	84	12%	115	14%
45-49	175.3	122	8%	55	8%	67	8%
50-54	102.9	79	5%	39	5%	40	5%
55-64	23.9	33	2%	19	3%	14	2%
65+	6.1	10	1%	3	<1%	7	1%
Race/Ethnicity							
White, not Hispanic	91.9	762	50%	333	46%	429	53%
Black/African American, not Hispanic	510.3	705	46%	360	50%	345	43%
Hispanic/Latino	93.9	22	1%	8	1%	14	2%
Asian/Pacific Islander	39.8	7	<1%	3	<1%	4	<1%
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	29	2%	18	2%	11	1%
Race/Sex							
White, not Hispanic Males	161.0	651	43%	277	38%	374	47%
White, not Hispanic Females	26.1	111	7%	56	8%	55	7%
Black/African American, not Hispanic Males	773.6	502	33%	250	35%	252	31%
Black/African American, not Hispanic Females	277.1	203	13%	110	15%	93	12%
Hispanic/Latino Males	164.1	20	1%	8	1%	12	1%
Hispanic/Latina Females	*	2	<1%	-	-	2	<1%
Asian/Pacific Islander Males	73.6	6	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	29	2%	18	2%	11	1%
Total	147.9	1,527		723		804	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Dayton Consortium 4

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	869	73%	375	69%	494	77%
Injection drug use (IDU)	29	2%	15	3%	14	2%
Male-to-male sexual contact and IDU	40	3%	14	3%	26	4%
Heterosexual contact	74	6%	26	5%	48	7%
Other/unknown	177	15%	114	21%	63	10%
Subtotal	1,189	100%	544	100%	645	100%
Female adult or adolescent						
Injection drug use	35	11%	21	13%	14	9%
Heterosexual contact	200	63%	93	55%	107	70%
Other/unknown	85	27%	54	32%	31	20%
Subtotal	320	100%	168	100%	152	100%
Child (<13 yrs at diagnosis)						
Perinatal	13	72%	9	82%	4	57%
Other/unknown	5	28%	2	18%	3	43%
Subtotal	18	100%	11	100%	7	100%
Total	1,527		723		804	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Dayton Consortium 4

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	823	54%	355	49%	468	58%
Injection drug use (IDU) only	37	2%	19	3%	18	2%
Heterosexual contact only	274	18%	119	16%	155	19%
MSM & IDU	36	2%	11	2%	25	3%
IDU & Heterosexual contact	27	2%	17	2%	10	1%
MSM & Heterosexual contact	46	3%	20	3%	26	3%
MSM & IDU & Heterosexual contact	4	<1%	3	<1%	1	<1%
Perinatal exposure	14	1%	10	1%	4	<1%
Other/unknown	266	17%	169	23%	97	12%
Total	1,527		723		804	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Dayton Consortium 4

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	518	80%	319	64%	18	90%	2	33%	-	-	12	63%
Injection drug use (IDU)	12	2%	16	3%	-	-	-	-	1	100%	-	-
Male-to-male sexual contact and IDU	20	3%	20	4%	-	-	-	-	-	-	-	-
Heterosexual contact	24	4%	49	10%	-	-	-	-	-	-	1	5%
Other/unknown	74	11%	91	18%	2	10%	4	67%	-	-	6	32%
Subtotal	648	100%	495	100%	20	100%	6	100%	1	100%	19	100%
Female adult or adolescent												
Injection drug use	16	14%	18	9%	-	-	1	100%	-	-	-	-
Heterosexual contact	75	68%	116	59%	2	100%	-	-	1	100%	6	60%
Other/unknown	20	18%	61	31%	-	-	-	-	-	-	4	40%
Subtotal	111	100%	195	100%	2	100%	1	100%	1	100%	10	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	33%	12	80%	-	-	-	-	-	-	-	-
Other/unknown	2	67%	3	20%	-	-	-	-	-	-	-	-
Subtotal	3	100%	15	100%	-	-	-	-	-	-	-	-
Total	762		705		22		7		2		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Dayton Consortium 4

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	497	65%	298	42%	16	73%	1	14%	-	-	11	38%
Injection drug use (IDU) only	16	2%	19	3%	-	-	1	14%	1	50%	-	-
Heterosexual contact only	99	13%	165	23%	2	9%	-	-	1	50%	7	24%
MSM & IDU	18	2%	18	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	12	2%	15	2%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	21	3%	21	3%	2	9%	1	14%	-	-	1	3%
MSM & IDU & Heterosexual contact	2	<1%	2	<1%	-	-	-	-	-	-	-	-
Perinatal exposure	1	<1%	13	2%	-	-	-	-	-	-	-	-
Other/unknown	96	13%	154	22%	2	9%	4	57%	-	-	10	34%
Total	762		705		22		7		2		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Dayton Consortium 4

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	933	78%	990	78%	1,044	78%	1,123	79%	1,199	79%
Females	260	22%	275	22%	290	22%	305	21%	328	21%
Age at end of year										
<13	11	1%	12	1%	13	1%	17	1%	17	1%
13-14	5	<1%	5	<1%	5	<1%	5	<1%	5	<1%
15-19	57	5%	63	5%	70	5%	83	6%	91	6%
20-24	146	12%	153	12%	168	13%	192	13%	220	14%
25-29	216	18%	227	18%	237	18%	253	18%	262	17%
30-34	231	19%	241	19%	250	19%	257	18%	269	18%
35-39	190	16%	198	16%	201	15%	207	14%	220	14%
40-44	165	14%	177	14%	182	14%	191	13%	199	13%
45-49	90	8%	97	8%	106	8%	111	8%	122	8%
50-54	54	5%	60	5%	65	5%	71	5%	79	5%
55-64	21	2%	25	2%	30	2%	31	2%	33	2%
65+	7	1%	7	1%	7	1%	10	1%	10	1%
Race/Ethnicity										
White, not Hispanic	636	53%	667	53%	690	52%	723	51%	762	50%
Black/African American, not Hispanic	511	43%	547	43%	592	44%	650	46%	705	46%
Hispanic/Latino	17	1%	19	2%	19	1%	19	1%	22	1%
Asian/Pacific Islander	6	1%	7	1%	7	1%	7	<1%	7	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	21	2%	23	2%	24	2%	27	2%	29	2%
Race/Sex										
White, not Hispanic Males	542	45%	568	45%	588	44%	620	43%	651	43%
White, not Hispanic Females	94	8%	99	8%	102	8%	103	7%	111	7%
Black/African American, not Hispanic Males	357	30%	384	30%	417	31%	462	32%	502	33%
Black/African American, not Hispanic Females	154	13%	163	13%	175	13%	188	13%	203	13%
Hispanic/Latino Males	15	1%	17	1%	17	1%	17	1%	20	1%
Hispanic/Latina Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Asian/Pacific Islander Males	5	<1%	6	<1%	6	<1%	6	<1%	6	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	21	2%	23	2%	24	2%	27	2%	29	2%
Total	1,193		1,265		1,334		1,428		1,527	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Dayton Consortium 4

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	678	73%	722	73%	756	73%	821	74%	869	73%
Injection drug use (IDU)	25	3%	26	3%	28	3%	28	3%	29	2%
Male-to-male sexual contact and IDU	38	4%	38	4%	39	4%	40	4%	40	3%
Heterosexual contact	58	6%	60	6%	63	6%	66	6%	74	6%
Other/unknown	129	14%	138	14%	151	15%	158	14%	177	15%
Subtotal	928	100%	984	100%	1,037	100%	1,113	100%	1,189	100%
Female adult or adolescent										
Injection drug use	30	12%	31	12%	32	11%	32	11%	35	11%
Heterosexual contact	170	67%	179	67%	186	66%	191	64%	200	63%
Other/unknown	54	21%	58	22%	65	23%	74	25%	85	27%
Subtotal	254	100%	268	100%	283	100%	297	100%	320	100%
Child (<13 yrs at diagnosis)										
Perinatal	9	82%	9	69%	10	71%	13	72%	13	72%
Other/unknown	2	18%	4	31%	4	29%	5	28%	5	28%
Subtotal	11	100%	13	100%	14	100%	18	100%	18	100%
Total	1,193		1,265		1,334		1,428		1,527	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Dayton Consortium 4

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	637	53%	679	54%	713	53%	778	54%	823	54%
Injection drug use (IDU) only	33	3%	34	3%	36	3%	36	3%	37	2%
Heterosexual contact only	228	19%	239	19%	249	19%	257	18%	274	18%
MSM & IDU	35	3%	35	3%	36	3%	36	3%	36	2%
IDU & Heterosexual contact	22	2%	23	2%	24	2%	24	2%	27	2%
MSM & Heterosexual contact	41	3%	43	3%	43	3%	43	3%	46	3%
MSM & IDU & Heterosexual contact	3	<1%	3	<1%	3	<1%	4	<1%	4	<1%
Perinatal exposure	9	1%	10	1%	11	1%	14	1%	14	1%
Other/unknown	185	16%	199	16%	219	16%	236	17%	266	17%
Total	1,193		1,265		1,334		1,428		1,527	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B Toledo
Consortium 5

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Toledo Consortium 5

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	13.1	52	84%	42	88%	7	78%	3	60%
Females	2.4	10	16%	6	13%	2	22%	2	40%
Age at diagnosis (yr)									
<13	*	1	2%	1	2%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	31.6	20	32%	18	38%	2	22%	-	-
25-29	15.8	8	13%	8	17%	-	-	-	-
30-34	18.5	9	15%	7	15%	2	22%	-	-
35-39	13.0	6	10%	5	10%	1	11%	-	-
40-44	17.8	9	15%	5	10%	3	33%	1	20%
45-49	*	4	6%	2	4%	1	11%	1	20%
50-54	*	2	3%	1	2%	-	-	1	20%
55-64	*	2	3%	-	-	-	-	2	40%
65+	*	1	2%	1	2%	-	-	-	-
Race/Ethnicity									
White, not Hispanic	5.6	36	58%	29	60%	5	56%	2	40%
Black/African American, not Hispanic	22.4	20	32%	16	33%	2	22%	2	40%
Hispanic/Latino	*	3	5%	2	4%	1	11%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	3	5%	1	2%	1	11%	1	20%
Race/Sex									
White, not Hispanic Males	9.5	30	48%	26	54%	4	44%	-	-
White, not Hispanic Females	1.8	6	10%	3	6%	1	11%	2	40%
Black/African American, not Hispanic Males	39.8	17	27%	13	27%	2	22%	2	40%
Black/African American, not Hispanic Females	*	3	5%	3	6%	-	-	-	-
Hispanic/Latino Males	*	2	3%	2	4%	-	-	-	-
Hispanic/Latina Females	*	1	2%	-	-	1	11%	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	3	5%	1	2%	1	11%	1	20%
Total	7.6	62		48		9		5	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Toledo Consortium 5

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	37	71%	31	74%	5	71%	1	33%
Injection drug use (IDU)	1	2%	1	2%	-	-	-	-
Male-to-male sexual contact and IDU	4	8%	4	10%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	10	19%	6	14%	2	29%	2	67%
Subtotal	52	100%	42	100%	7	100%	3	100%
Female adult or adolescent								
Injection drug use	1	11%	1	20%	-	-	-	-
Heterosexual contact	4	44%	3	60%	-	-	1	50%
Other/unknown	4	44%	1	20%	2	100%	1	50%
Subtotal	9	100%	5	100%	2	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	62		48		9		5	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	36	58%	30	63%	5	56%	1	20%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	6%	3	6%	-	-	1	20%
MSM & IDU	4	6%	4	8%	-	-	-	-
IDU & Heterosexual contact	2	3%	2	4%	-	-	-	-
MSM & Heterosexual contact	1	2%	1	2%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	1	2%	-	-	-	-
Other/unknown	14	23%	7	15%	4	44%	3	60%
Total	62		48		9		5	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Toledo Consortium 5

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	22	73%	11	65%	2	100%	-	-	-	-	2	67%
Injection drug use (IDU)	1	3%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	3	10%	-	-	-	-	-	-	-	-	1	33%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	13%	6	35%	-	-	-	-	-	-	-	-
Subtotal	30	100%	17	100%	2	100%	-	-	-	-	3	100%
Female adult or adolescent												
Injection drug use	1	17%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	50%	1	50%	-	-	-	-	-	-	-	-
Other/unknown	2	33%	1	50%	1	100%	-	-	-	-	-	-
Subtotal	6	100%	2	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	36		20		3		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Toledo Consortium 5

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	21	58%	11	55%	2	67%	-	-	-	-	2	67%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	8%	1	5%	-	-	-	-	-	-	-	-
MSM & IDU	3	8%	-	-	-	-	-	-	-	-	1	33%
IDU & Heterosexual contact	2	6%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	5%	-	-	-	-	-	-	-	-
Other/unknown	6	17%	7	35%	1	33%	-	-	-	-	-	-
Total	36		20		3		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Toledo Consortium 5**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	11.1	44	72%	8.5	34	79%	8.5	34	83%	11.3	45	80%
Females	4.1	17	28%	2.2	9	21%	1.7	7	17%	2.6	11	20%
Age at diagnosis (yr)												
<13	*	1	2%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	3%	*	1	2%	*	2	5%	*	4	7%
20-24	20.6	13	21%	11.1	7	16%	12.6	8	20%	17.4	11	20%
25-29	19.7	10	16%	*	4	9%	*	2	5%	11.8	6	11%
30-34	16.4	8	13%	*	3	7%	12.3	6	15%	14.4	7	13%
35-39	32.6	15	25%	15.2	7	16%	21.7	10	24%	21.7	10	18%
40-44	*	3	5%	*	4	9%	*	2	5%	11.9	6	11%
45-49	*	4	7%	11.2	6	14%	*	4	10%	*	4	7%
50-54	*	3	5%	8.2	5	12%	*	3	7%	8.2	5	9%
55-64	*	2	3%	5.5	6	14%	*	4	10%	*	3	5%
65+	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity												
White, not Hispanic	3.2	21	34%	2.8	18	42%	3.7	24	59%	3.6	23	41%
Black/African American, not Hispanic	35.8	32	52%	24.6	22	51%	15.7	14	34%	31.3	28	50%
Hispanic/Latino	9.6	5	8%	*	2	5%	*	1	2%	*	2	4%
Asian/Pacific Islander	*	2	3%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	1	2%	*	2	5%	*	3	5%
Race/Sex												
White, not Hispanic Males	5.7	18	30%	4.7	15	35%	6.6	21	51%	6.6	21	38%
White, not Hispanic Females	*	3	5%	*	3	7%	*	3	7%	*	2	4%
Black/African American, not Hispanic Males	46.9	20	33%	39.8	17	40%	23.4	10	24%	44.5	19	34%
Black/African American, not Hispanic Females	25.7	12	20%	10.7	5	12%	*	4	10%	19.3	9	16%
Hispanic/Latino Males	*	4	7%	*	1	2%	*	1	2%	*	2	4%
Hispanic/Latina Females	*	1	2%	*	1	2%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	2%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	1	2%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	1	2%	*	2	5%	*	3	5%
Total	7.5	61		5.3	43		5.0	41		6.9	56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Toledo Consortium 5

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	31	70%	28	82%	29	85%	32	71%
Injection drug use (IDU)	1	2%	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	3	7%	1	3%	1	3%	2	4%
Heterosexual contact	8	18%	4	12%	3	9%	4	9%
Other/unknown	1	2%	1	3%	1	3%	6	13%
Subtotal	44	100%	34	100%	34	100%	45	100%
Female adult or adolescent								
Injection drug use	1	6%	-	-	-	-	-	-
Heterosexual contact	14	88%	6	67%	6	86%	8	73%
Other/unknown	1	6%	3	33%	1	14%	3	27%
Subtotal	16	100%	9	100%	7	100%	11	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-
Total	61		43		41		56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Toledo Consortium 5

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	30	49%	26	60%	28	68%	31	55%
Injection drug use (IDU) only	1	2%	-	-	-	-	1	2%
Heterosexual contact only	22	36%	10	23%	9	22%	12	21%
MSM & IDU	1	2%	1	2%	1	2%	1	2%
IDU & Heterosexual contact	1	2%	-	-	-	-	-	-
MSM & Heterosexual contact	1	2%	2	5%	1	2%	1	2%
MSM & IDU & Heterosexual contact	2	3%	-	-	-	-	1	2%
Perinatal exposure	1	2%	-	-	-	-	-	-
Other/unknown	2	3%	4	9%	2	5%	9	16%
Total	61		43		41		56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Toledo Consortium 5

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	27	69%	21	81%	22	85%	27	79%	42	88%
Females	12	31%	5	19%	4	15%	7	21%	6	13%
Age at HIV diagnosis (yr)										
<13	1	3%	-	-	-	-	-	-	1	2%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	2	5%	1	4%	2	8%	3	9%	-	-
20-24	11	28%	6	23%	7	27%	10	29%	18	38%
25-29	6	15%	4	15%	2	8%	5	15%	8	17%
30-34	2	5%	2	8%	2	8%	5	15%	7	15%
35-39	10	26%	4	15%	7	27%	4	12%	5	10%
40-44	2	5%	1	4%	1	4%	3	9%	5	10%
45-49	2	5%	5	19%	1	4%	2	6%	2	4%
50-54	2	5%	2	8%	1	4%	1	3%	1	2%
55-64	1	3%	1	4%	3	12%	1	3%	-	-
65+	-	-	-	-	-	-	-	-	1	2%
Race/Ethnicity										
White, not Hispanic	13	33%	11	42%	13	50%	13	38%	29	60%
Black/African American, not Hispanic	21	54%	12	46%	11	42%	17	50%	16	33%
Hispanic/Latino	3	8%	2	8%	-	-	1	3%	2	4%
Asian/Pacific Islander	1	3%	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	1	4%	2	8%	3	9%	1	2%
Race/Sex										
White, not Hispanic Males	11	28%	10	38%	11	42%	11	32%	26	54%
White, not Hispanic Females	2	5%	1	4%	2	8%	2	6%	3	6%
Black/African American, not Hispanic Males	11	28%	9	35%	9	35%	12	35%	13	27%
Black/African American, not Hispanic Females	10	26%	3	12%	2	8%	5	15%	3	6%
Hispanic/Latino Males	3	8%	1	4%	-	-	1	3%	2	4%
Hispanic/Latina Females	-	-	1	4%	-	-	-	-	-	-
Asian/Pacific Islander Males	1	3%	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	1	4%	2	8%	3	9%	1	2%
Total	39		26		26		34		48	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Toledo Consortium 5

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	20	74%	18	86%	17	77%	21	78%	31	74%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	1	4%	1	5%	1	5%	2	7%	4	10%
Heterosexual contact	5	19%	2	10%	3	14%	1	4%	-	-
Other/unknown	1	4%	-	-	1	5%	3	11%	6	14%
Subtotal	27	100%	21	100%	22	100%	27	100%	42	100%
Female adult or adolescent										
Injection drug use	1	9%	-	-	-	-	-	-	1	20%
Heterosexual contact	10	91%	3	60%	4	100%	5	71%	3	60%
Other/unknown	-	-	2	40%	-	-	2	29%	1	20%
Subtotal	11	100%	5	100%	4	100%	7	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	1	100%
Total	39		26		26		34		48	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	20	51%	16	62%	16	62%	21	62%	30	63%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	15	38%	5	19%	7	27%	6	18%	3	6%
MSM & IDU	-	-	1	4%	1	4%	1	3%	4	8%
IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	2	4%
MSM & Heterosexual contact	-	-	2	8%	1	4%	-	-	1	2%
MSM & IDU & Heterosexual contact	1	3%	-	-	-	-	1	3%	-	-
Perinatal exposure	1	3%	-	-	-	-	-	-	1	2%
Other/unknown	1	3%	2	8%	1	4%	5	15%	7	15%
Total	39		26		26		34		48	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Toledo Consortium 5

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	23	64%	27	75%	24	77%	29	83%	14	78%	873	80%
Females	13	36%	9	25%	7	23%	6	17%	4	22%	220	20%
Age at AIDS diagnosis (yr)												
<13	1	3%	-	-	-	-	-	-	-	-	18	2%
13-14	1	3%	-	-	-	-	-	-	-	-	1	<1%
15-19	1	3%	1	3%	3	10%	1	3%	-	-	33	3%
20-24	7	19%	5	14%	1	3%	3	9%	2	11%	102	9%
25-29	5	14%	7	19%	5	16%	2	6%	-	-	173	16%
30-34	7	19%	7	19%	6	19%	5	14%	3	17%	216	20%
35-39	9	25%	7	19%	4	13%	8	23%	3	17%	210	19%
40-44	1	3%	2	6%	5	16%	6	17%	4	22%	134	12%
45-49	2	6%	-	-	4	13%	3	9%	3	17%	91	8%
50-54	1	3%	3	8%	2	6%	4	11%	1	6%	63	6%
55-64	1	3%	4	11%	1	3%	3	9%	2	11%	44	4%
65+	-	-	-	-	-	-	-	-	-	-	8	1%
Race/Ethnicity												
White, not Hispanic	16	44%	13	36%	18	58%	16	46%	11	61%	649	59%
Black/African American, not Hispanic	18	50%	22	61%	10	32%	16	46%	4	22%	377	34%
Hispanic/Latino	1	3%	1	3%	3	10%	3	9%	1	6%	51	5%
Asian/Pacific Islander	1	3%	-	-	-	-	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	2	11%	11	1%
Race/Sex												
White, not Hispanic Males	14	39%	10	28%	14	45%	15	43%	8	44%	565	52%
White, not Hispanic Females	2	6%	3	8%	4	13%	1	3%	3	17%	84	8%
Black/African American, not Hispanic Males	9	25%	16	44%	7	23%	11	31%	4	22%	257	24%
Black/African American, not Hispanic Females	9	25%	6	17%	3	10%	5	14%	-	-	120	11%
Hispanic/Latino Males	-	-	1	3%	3	10%	3	9%	-	-	37	3%
Hispanic/Latina Females	1	3%	-	-	-	-	-	-	1	6%	14	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	1	3%	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	2	11%	11	1%
Total	36		36		31		35		18		1,093	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Toledo Consortium 5

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	18	78%	21	78%	20	83%	19	66%	9	64%	626	73%
Injection drug use (IDU)	1	4%	-	-	-	-	1	3%	-	-	47	5%
Male-to-male sexual contact and IDU	1	4%	1	4%	1	4%	1	3%	1	7%	66	8%
Heterosexual contact	2	9%	4	15%	3	13%	3	10%	-	-	58	7%
Other/unknown	1	4%	1	4%	-	-	5	17%	4	29%	66	8%
Subtotal	23	100%	27	100%	24	100%	29	100%	14	100%	863	100%
Female adult or adolescent												
Injection drug use	1	8%	-	-	-	-	-	-	-	-	31	15%
Heterosexual contact	10	83%	6	67%	6	86%	5	83%	1	25%	150	71%
Other/unknown	1	8%	3	33%	1	14%	1	17%	3	75%	31	15%
Subtotal	12	100%	9	100%	7	100%	6	100%	4	100%	212	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	15	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	17%
Subtotal	1	100%	-	-	-	-	-	-	-	-	18	100%
Total	36		36		31		35		18		1,093	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	17	47%	20	56%	20	65%	18	51%	8	44%	588	54%
Injection drug use (IDU) only	1	3%	-	-	-	-	1	3%	-	-	39	4%
Heterosexual contact only	12	33%	10	28%	9	29%	8	23%	1	6%	208	19%
MSM & IDU	-	-	1	3%	1	3%	1	3%	1	6%	50	5%
IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	39	4%
MSM & Heterosexual contact	1	3%	1	3%	-	-	1	3%	1	6%	38	3%
MSM & IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	16	1%
Perinatal exposure	1	3%	-	-	-	-	-	-	-	-	15	1%
Other/unknown	2	6%	4	11%	1	3%	6	17%	7	39%	100	9%
Total	36		36		31		35		18		1,093	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Toledo Consortium 5

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Toledo Consortium 5

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	184.5	734	72%	347	71%	387	74%
Females	67.4	281	28%	144	29%	137	26%
Age at end of year							
<13	12.0	16	2%	9	2%	7	1%
13-14	*	2	<1%	1	<1%	1	<1%
15-19	77.3	45	4%	22	4%	23	4%
20-24	254.5	161	16%	101	21%	60	11%
25-29	303.4	154	15%	81	16%	73	14%
30-34	365.5	178	18%	76	15%	102	19%
35-39	430.6	198	20%	90	18%	108	21%
40-44	225.2	114	11%	48	10%	66	13%
45-49	145.7	78	8%	35	7%	43	8%
50-54	65.4	40	4%	18	4%	22	4%
55-64	25.7	28	3%	10	2%	18	3%
65+	*	1	<1%	-	-	1	<1%
Race/Ethnicity							
White, not Hispanic	80.3	519	51%	251	51%	268	51%
Black/African American, not Hispanic	458.7	410	40%	200	41%	210	40%
Hispanic/Latino	107.4	56	6%	22	4%	34	6%
Asian/Pacific Islander	67.9	7	1%	4	1%	3	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	23	2%	14	3%	9	2%
Race/Sex							
White, not Hispanic Males	132.9	420	41%	196	40%	224	43%
White, not Hispanic Females	29.9	99	10%	55	11%	44	8%
Black/African American, not Hispanic Males	585.7	250	25%	121	25%	129	25%
Black/African American, not Hispanic Females	342.6	160	16%	79	16%	81	15%
Hispanic/Latino Males	145.7	38	4%	14	3%	24	5%
Hispanic/Latina Females	69.1	18	2%	8	2%	10	2%
Asian/Pacific Islander Males	121.5	6	1%	4	1%	2	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	23	2%	14	3%	9	2%
Total	124.6	1,015		491		524	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Toledo Consortium 5

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	532	73%	249	73%	283	74%
Injection drug use (IDU)	31	4%	13	4%	18	5%
Male-to-male sexual contact and IDU	48	7%	21	6%	27	7%
Heterosexual contact	70	10%	33	10%	37	10%
Other/unknown	46	6%	27	8%	19	5%
Subtotal	727	100%	343	100%	384	100%
Female adult or adolescent						
Injection drug use	24	9%	10	7%	14	11%
Heterosexual contact	210	77%	106	76%	104	78%
Other/unknown	38	14%	23	17%	15	11%
Subtotal	272	100%	139	100%	133	100%
Child (<13 yrs at diagnosis)						
Perinatal	16	100%	9	100%	7	100%
Other/unknown	-	-	-	-	-	-
Subtotal	16	100%	9	100%	7	100%
Total	1,015		491		524	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	486	48%	229	47%	257	49%
Injection drug use (IDU) only	23	2%	10	2%	13	2%
Heterosexual contact only	280	28%	139	28%	141	27%
MSM & IDU	35	3%	17	3%	18	3%
IDU & Heterosexual contact	32	3%	13	3%	19	4%
MSM & Heterosexual contact	46	5%	20	4%	26	5%
MSM & IDU & Heterosexual contact	13	1%	4	1%	9	2%
Perinatal exposure	16	2%	9	2%	7	1%
Other/unknown	84	8%	50	10%	34	6%
Total	1,015		491		524	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Toledo Consortium 5

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	336	81%	154	62%	28	74%	3	50%	-	-	11	58%
Injection drug use (IDU)	12	3%	13	5%	4	11%	-	-	-	-	2	11%
Male-to-male sexual contact and IDU	31	7%	14	6%	3	8%	-	-	-	-	-	-
Heterosexual contact	23	6%	44	18%	1	3%	-	-	-	-	2	11%
Other/unknown	15	4%	22	9%	2	5%	3	50%	-	-	4	21%
Subtotal	417	100%	247	100%	38	100%	6	100%	-	-	19	100%
Female adult or adolescent												
Injection drug use	7	7%	17	11%	-	-	-	-	-	-	-	-
Heterosexual contact	79	81%	119	77%	12	71%	-	-	-	-	-	-
Other/unknown	12	12%	18	12%	5	29%	1	100%	-	-	2	100%
Subtotal	98	100%	154	100%	17	100%	1	100%	-	-	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	4	100%	9	100%	1	100%	-	-	-	-	2	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	9	100%	1	100%	-	-	-	-	2	100%
Total	519		410		56		7		0		23	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	310	60%	138	34%	25	45%	2	29%	-	-	11	48%
Injection drug use (IDU) only	12	2%	10	2%	1	2%	-	-	-	-	-	-
Heterosexual contact only	102	20%	163	40%	13	23%	-	-	-	-	2	9%
MSM & IDU	21	4%	12	3%	2	4%	-	-	-	-	-	-
IDU & Heterosexual contact	7	1%	20	5%	3	5%	-	-	-	-	2	9%
MSM & Heterosexual contact	26	5%	16	4%	3	5%	1	14%	-	-	-	-
MSM & IDU & Heterosexual contact	10	2%	2	0%	1	2%	-	-	-	-	-	-
Perinatal exposure	4	1%	9	2%	1	2%	-	-	-	-	2	9%
Other/unknown	27	5%	40	10%	7	13%	4	57%	-	-	6	26%
Total	519		410		56		7		0		23	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Toledo Consortium 5

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	581	71%	625	71%	659	71%	690	72%	734	72%
Females	240	29%	257	29%	265	29%	271	28%	281	28%
Age at end of year										
<13	15	2%	16	2%	16	2%	16	2%	16	2%
13-14	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
15-19	36	4%	38	4%	39	4%	41	4%	45	4%
20-24	122	15%	136	15%	143	15%	151	16%	161	16%
25-29	132	16%	142	16%	146	16%	148	15%	154	15%
30-34	156	19%	163	18%	166	18%	171	18%	178	18%
35-39	157	19%	172	20%	179	19%	188	20%	198	20%
40-44	99	12%	102	12%	106	11%	108	11%	114	11%
45-49	61	7%	65	7%	70	8%	74	8%	78	8%
50-54	24	3%	27	3%	32	3%	35	4%	40	4%
55-64	16	2%	18	2%	24	3%	26	3%	28	3%
65+	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Race/Ethnicity										
White, not Hispanic	436	53%	457	52%	475	51%	496	52%	519	51%
Black/African American, not Hispanic	318	39%	350	40%	371	40%	384	40%	410	40%
Hispanic/Latino	46	6%	51	6%	53	6%	54	6%	56	6%
Asian/Pacific Islander	5	1%	7	1%	7	1%	7	1%	7	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	16	2%	17	2%	18	2%	20	2%	23	2%
Race/Sex										
White, not Hispanic Males	348	42%	366	41%	381	41%	399	42%	420	41%
White, not Hispanic Females	88	11%	91	10%	94	10%	97	10%	99	10%
Black/African American, not Hispanic Males	185	23%	205	23%	222	24%	232	24%	250	25%
Black/African American, not Hispanic Females	133	16%	145	16%	149	16%	152	16%	160	16%
Hispanic/Latino Males	30	4%	34	4%	35	4%	36	4%	38	4%
Hispanic/Latina Females	16	2%	17	2%	18	2%	18	2%	18	2%
Asian/Pacific Islander Males	5	1%	6	1%	6	1%	6	1%	6	1%
Asian/Pacific Islander Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	16	2%	17	2%	18	2%	20	2%	23	2%
Total	821		882		924		961		1,015	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Toledo Consortium 5

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	414	72%	446	72%	474	73%	501	73%	532	73%
Injection drug use (IDU)	29	5%	30	5%	30	5%	30	4%	31	4%
Male-to-male sexual contact and IDU	41	7%	44	7%	45	7%	46	7%	48	7%
Heterosexual contact	52	9%	59	10%	63	10%	66	10%	70	10%
Other/unknown	38	7%	39	6%	40	6%	40	6%	46	6%
Subtotal	574	100%	618	100%	652	100%	683	100%	727	100%
Female adult or adolescent										
Injection drug use	23	10%	24	10%	24	9%	24	9%	24	9%
Heterosexual contact	178	77%	192	77%	198	77%	203	77%	210	77%
Other/unknown	31	13%	32	13%	34	13%	35	13%	38	14%
Subtotal	232	100%	248	100%	256	100%	262	100%	272	100%
Child (<13 yrs at diagnosis)										
Perinatal	15	100%	16	100%	16	100%	16	100%	16	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	15	100%	16	100%	16	100%	16	100%	16	100%
Total	821		882		924		961		1,015	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Toledo Consortium 5

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	373	45%	404	46%	430	47%	456	47%	486	48%
Injection drug use (IDU) only	21	3%	22	2%	22	2%	22	2%	23	2%
Heterosexual contact only	230	28%	251	28%	261	28%	269	28%	280	28%
MSM & IDU	31	4%	32	4%	33	4%	34	4%	35	3%
IDU & Heterosexual contact	31	4%	32	4%	32	3%	32	3%	32	3%
MSM & Heterosexual contact	41	5%	42	5%	44	5%	45	5%	46	5%
MSM & IDU & Heterosexual contact	10	1%	12	1%	12	1%	12	1%	13	1%
Perinatal exposure	15	2%	16	2%	16	2%	16	2%	16	2%
Other/unknown	69	8%	71	8%	74	8%	75	8%	84	8%
Total	821		882		924		961		1,015	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B Akron
Consortium 6

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Akron Consortium 6

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	8.7	37	79%	30	81%	6	75%	1	50%
Females	2.2	10	21%	7	19%	2	25%	1	50%
Age at diagnosis (yr)									
<13	*	1	2%	1	3%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	9.6	6	13%	6	16%	-	-	-	-
20-24	11.1	7	15%	6	16%	1	13%	-	-
25-29	15.7	8	17%	6	16%	2	25%	-	-
30-34	17.7	9	19%	8	22%	1	13%	-	-
35-39	*	2	4%	2	5%	-	-	-	-
40-44	*	3	6%	2	5%	1	13%	-	-
45-49	*	2	4%	2	5%	-	-	-	-
50-54	*	4	9%	1	3%	2	25%	1	50%
55-64	*	3	6%	1	3%	1	13%	1	50%
65+	*	2	4%	2	5%	-	-	-	-
Race/Ethnicity									
White, not Hispanic	3.4	25	53%	20	54%	4	50%	1	50%
Black/African American, not Hispanic	22.9	20	43%	15	41%	4	50%	1	50%
Hispanic/Latino	*	1	2%	1	3%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	2%	1	3%	-	-	-	-
Race/Sex									
White, not Hispanic Males	5.3	19	40%	15	41%	3	38%	1	50%
White, not Hispanic Females	1.6	6	13%	5	14%	1	13%	-	-
Black/African American, not Hispanic Males	39.4	16	34%	13	35%	3	38%	-	-
Black/African American, not Hispanic Females	*	4	9%	2	5%	1	13%	1	50%
Hispanic/Latino Males	*	1	2%	1	3%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	2%	1	3%	-	-	-	-
Total	5.4	47		37		8		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Akron Consortium 6

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	23	62%	19	63%	4	67%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	3%	1	3%	-	-	-	-
Other/unknown	13	35%	10	33%	2	33%	1	100%
Subtotal	37	100%	30	100%	6	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	4	44%	3	50%	1	50%	-	-
Other/unknown	5	56%	3	50%	1	50%	1	100%
Subtotal	9	100%	6	100%	2	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	47		37		8		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Akron Consortium 6

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	22	47%	18	49%	4	50%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	5	11%	4	11%	1	13%	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	2%	1	3%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	19	40%	14	38%	3	38%	2	100%
Total	47		37		8		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Akron Consortium 6

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	12	63%	10	63%	1	100%	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	6%	-	-	-	-	-	-	-	-
Other/unknown	7	37%	5	31%	-	-	-	-	-	-	1	100%
Subtotal	19	100%	16	100%	1	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	33%	2	67%	-	-	-	-	-	-	-	-
Other/unknown	4	67%	1	33%	-	-	-	-	-	-	-	-
Subtotal	6	100%	3	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	25		20		1		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Akron Consortium 6

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	48%	10	50%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	8%	3	15%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	1	100%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	11	44%	7	35%	-	-	-	-	-	-	1	100%
Total	25		20		1		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Akron Consortium 6**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	8.4	36	84%	9.6	41	82%	5.6	24	75%	9.9	42	82%
Females	1.6	7	16%	2.0	9	18%	1.8	8	25%	2.0	9	18%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	2	4%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	3	7%	*	2	4%	*	-	-	*	3	6%
20-24	15.9	10	23%	23.9	15	30%	8.0	5	16%	19.1	12	24%
25-29	15.7	8	19%	13.7	7	14%	*	2	6%	15.7	8	16%
30-34	*	4	9%	*	4	8%	17.7	9	28%	*	2	4%
35-39	*	3	7%	*	2	4%	*	2	6%	14.2	7	14%
40-44	*	4	9%	13.6	8	16%	*	-	-	10.2	6	12%
45-49	7.9	5	12%	12.6	8	16%	7.9	5	16%	*	2	4%
50-54	*	2	5%	*	1	2%	7.3	5	16%	10.2	7	14%
55-64	*	4	9%	*	3	6%	*	1	3%	*	2	4%
65+	*	-	-	*	-	-	*	3	9%	*	-	-
Race/Ethnicity												
White, not Hispanic	3.0	22	51%	3.9	29	58%	2.4	18	56%	3.8	28	55%
Black/African American, not Hispanic	21.8	19	44%	22.9	20	40%	12.6	11	34%	22.9	20	39%
Hispanic/Latino	*	2	5%	*	1	2%	*	-	-	*	1	2%
Asian/Pacific Islander	*	-	-	*	-	-	*	2	6%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	1	3%	*	2	4%
Race/Sex												
White, not Hispanic Males	5.5	20	47%	6.4	23	46%	3.6	13	41%	6.4	23	45%
White, not Hispanic Females	*	2	5%	1.6	6	12%	1.3	5	16%	1.3	5	10%
Black/African American, not Hispanic Males	34.4	14	33%	41.8	17	34%	22.1	9	28%	39.4	16	31%
Black/African American, not Hispanic Females	10.8	5	12%	*	3	6%	*	2	6%	*	4	8%
Hispanic/Latino Males	*	2	5%	*	1	2%	*	-	-	*	1	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	1	3%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	3%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	1	3%	*	2	4%
Total	4.9	43		5.7	50		3.7	32		5.8	51	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Akron Consortium 6

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	32	89%	22	54%	15	63%	26	65%
Injection drug use (IDU)	-	-	2	5%	1	4%	1	3%
Male-to-male sexual contact and IDU	-	-	-	-	2	8%	-	-
Heterosexual contact	3	8%	4	10%	1	4%	2	5%
Other/unknown	1	3%	13	32%	5	21%	11	28%
Subtotal	36	100%	41	100%	24	100%	40	100%
Female adult or adolescent								
Injection drug use	2	29%	1	11%	-	-	-	-
Heterosexual contact	1	14%	7	78%	4	50%	2	22%
Other/unknown	4	57%	1	11%	4	50%	7	78%
Subtotal	7	100%	9	100%	8	100%	9	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	2	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	2	100%
Total	43		50		32		51	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Akron Consortium 6

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	31	72%	22	44%	14	44%	25	49%
Injection drug use (IDU) only	1	2%	-	-	1	3%	-	-
Heterosexual contact only	4	9%	11	22%	5	16%	4	8%
MSM & IDU	-	-	-	-	1	3%	-	-
IDU & Heterosexual contact	1	2%	3	6%	-	-	1	2%
MSM & Heterosexual contact	1	2%	-	-	1	3%	1	2%
MSM & IDU & Heterosexual contact	-	-	-	-	1	3%	-	-
Perinatal exposure	-	-	-	-	-	-	2	4%
Other/unknown	5	12%	14	28%	9	28%	18	35%
Total	43		50		32		51	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Akron Consortium 6

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	25	83%	31	86%	16	73%	38	84%	30	81%
Females	5	17%	5	14%	6	27%	7	16%	7	19%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	2	4%	1	3%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	10%	1	3%	-	-	3	7%	6	16%
20-24	6	20%	14	39%	5	23%	11	24%	6	16%
25-29	7	23%	4	11%	1	5%	6	13%	6	16%
30-34	2	7%	3	8%	5	23%	2	4%	8	22%
35-39	1	3%	2	6%	2	9%	7	16%	2	5%
40-44	2	7%	3	8%	-	-	5	11%	2	5%
45-49	4	13%	7	19%	4	18%	1	2%	2	5%
50-54	1	3%	1	3%	2	9%	6	13%	1	3%
55-64	4	13%	1	3%	1	5%	2	4%	1	3%
65+	-	-	-	-	2	9%	-	-	2	5%
Race/Ethnicity										
White, not Hispanic	19	63%	20	56%	12	55%	25	56%	20	54%
Black/African American, not Hispanic	10	33%	15	42%	9	41%	18	40%	15	41%
Hispanic/Latino	1	3%	1	3%	-	-	1	2%	1	3%
Asian/Pacific Islander	-	-	-	-	1	5%	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	1	2%	1	3%
Race/Sex										
White, not Hispanic Males	17	57%	17	47%	9	41%	20	44%	15	41%
White, not Hispanic Females	2	7%	3	8%	3	14%	5	11%	5	14%
Black/African American, not Hispanic Males	7	23%	13	36%	7	32%	16	36%	13	35%
Black/African American, not Hispanic Females	3	10%	2	6%	2	9%	2	4%	2	5%
Hispanic/Latino Males	1	3%	1	3%	-	-	1	2%	1	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	1	5%	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	1	2%	1	3%
Total	30		36		22		45		37	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Akron Consortium 6

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	23	92%	16	52%	11	69%	24	67%	19	63%
Injection drug use (IDU)	-	-	-	-	-	-	1	3%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	2	13%	-	-	-	-
Heterosexual contact	2	8%	3	10%	-	-	1	3%	1	3%
Other/unknown	-	-	12	39%	3	19%	10	28%	10	33%
Subtotal	25	100%	31	100%	16	100%	36	100%	30	100%
Female adult or adolescent										
Injection drug use	1	20%	1	20%	-	-	-	-	-	-
Heterosexual contact	1	20%	4	80%	4	67%	2	29%	3	50%
Other/unknown	3	60%	-	-	2	33%	5	71%	3	50%
Subtotal	5	100%	5	100%	6	100%	7	100%	6	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	2	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	100%
Subtotal	-	-	-	-	-	-	2	100%	1	100%
Total	30		36		22		45		37	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Akron Consortium 6

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	23	77%	16	44%	10	45%	23	51%	18	49%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	10%	7	19%	4	18%	3	7%	4	11%
MSM & IDU	-	-	-	-	1	5%	-	-	-	-
IDU & Heterosexual contact	1	3%	1	3%	-	-	1	2%	-	-
MSM & Heterosexual contact	-	-	-	-	1	5%	1	2%	1	3%
MSM & IDU & Heterosexual contact	-	-	-	-	1	5%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	2	4%	-	-
Other/unknown	3	10%	12	33%	5	23%	15	33%	14	38%
Total	30		36		22		45		37	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Akron Consortium 6

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	15	79%	20	87%	12	75%	14	82%	9	75%	755	86%
Females	4	21%	3	13%	4	25%	3	18%	3	25%	118	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	1	<1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	-	-	-	-	1	6%	1	6%	-	-	18	2%
20-24	3	16%	4	17%	1	6%	4	24%	1	8%	79	9%
25-29	1	5%	4	17%	2	13%	3	18%	2	17%	125	14%
30-34	2	11%	4	17%	2	13%	3	18%	2	17%	192	22%
35-39	6	32%	2	9%	1	6%	-	-	-	-	168	19%
40-44	4	21%	5	22%	2	13%	4	24%	1	8%	125	14%
45-49	2	11%	2	9%	1	6%	1	6%	-	-	77	9%
50-54	1	5%	1	4%	3	19%	1	6%	4	33%	50	6%
55-64	-	-	1	4%	2	13%	-	-	2	17%	31	4%
65+	-	-	-	-	1	6%	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	7	37%	10	43%	11	69%	10	59%	5	42%	552	63%
Black/African American, not Hispanic	12	63%	10	43%	3	19%	6	35%	7	58%	298	34%
Hispanic/Latino	-	-	1	4%	-	-	-	-	-	-	14	2%
Asian/Pacific Islander	-	-	1	4%	1	6%	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	4%	1	6%	1	6%	-	-	6	1%
Race/Sex												
White, not Hispanic Males	6	32%	9	39%	8	50%	9	53%	4	33%	499	57%
White, not Hispanic Females	1	5%	1	4%	3	19%	1	6%	1	8%	53	6%
Black/African American, not Hispanic Males	9	47%	9	39%	2	13%	4	24%	5	42%	238	27%
Black/African American, not Hispanic Females	3	16%	1	4%	1	6%	2	12%	2	17%	60	7%
Hispanic/Latino Males	-	-	1	4%	-	-	-	-	-	-	11	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Males	-	-	-	-	1	6%	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	1	4%	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	4%	1	6%	1	6%	-	-	6	1%
Total	19		23		16		17		12		873	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Akron Consortium 6

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	7	47%	14	70%	7	58%	10	71%	4	44%	540	72%
Injection drug use (IDU)	-	-	3	15%	1	8%	1	7%	-	-	45	6%
Male-to-male sexual contact and IDU	1	7%	-	-	-	-	-	-	-	-	43	6%
Heterosexual contact	3	20%	2	10%	1	8%	2	14%	1	11%	40	5%
Other/unknown	4	27%	1	5%	3	25%	1	7%	4	44%	86	11%
Subtotal	15	100%	20	100%	12	100%	14	100%	9	100%	754	100%
Female adult or adolescent												
Injection drug use	2	50%	-	-	-	-	-	-	-	-	17	14%
Heterosexual contact	1	25%	2	67%	1	25%	1	33%	1	33%	67	57%
Other/unknown	1	25%	1	33%	3	75%	2	67%	2	67%	34	29%
Subtotal	4	100%	3	100%	4	100%	3	100%	3	100%	118	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	1	100%
Total	19		23		16		17		12		873	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Akron Consortium 6

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	6	32%	14	61%	7	44%	9	53%	4	33%	526	60%
Injection drug use (IDU) only	1	5%	1	4%	1	6%	-	-	-	-	36	4%
Heterosexual contact only	4	21%	4	17%	2	13%	3	18%	2	17%	107	12%
MSM & IDU	1	5%	-	-	-	-	-	-	-	-	40	5%
IDU & Heterosexual contact	1	5%	2	9%	-	-	1	6%	-	-	26	3%
MSM & Heterosexual contact	1	5%	-	-	-	-	1	6%	-	-	14	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	<1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	1	<1%
Other/unknown	5	26%	2	9%	6	38%	3	18%	6	50%	120	14%
Total	19		23		16		17		12		873	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Akron Consortium 6

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Akron Consortium 6

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	158.5	676	81%	340	79%	336	83%
Females	35.6	160	19%	90	21%	70	17%
Age at end of year							
<13	7.4	10	1%	9	2%	1	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	43.3	27	3%	19	4%	8	2%
20-24	182.9	115	14%	73	17%	42	10%
25-29	256.7	131	16%	77	18%	54	13%
30-34	286.8	146	17%	63	15%	83	20%
35-39	286.6	141	17%	65	15%	76	19%
40-44	213.7	126	15%	55	13%	71	17%
45-49	115.1	73	9%	35	8%	38	9%
50-54	58.1	40	5%	17	4%	23	6%
55-64	16.6	20	2%	13	3%	7	2%
65+	4.6	6	1%	4	1%	2	<1%
Race/Ethnicity							
White, not Hispanic	64.8	479	57%	237	55%	242	60%
Black/African American, not Hispanic	364.8	318	38%	172	40%	146	36%
Hispanic/Latino	147.4	22	3%	11	3%	11	3%
Asian/Pacific Islander	34.1	6	1%	4	1%	2	<1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	10	1%	5	1%	5	1%
Race/Sex							
White, not Hispanic Males	115.4	417	50%	206	48%	211	52%
White, not Hispanic Females	16.4	62	7%	31	7%	31	8%
Black/African American, not Hispanic Males	560.8	228	27%	117	27%	111	27%
Black/African American, not Hispanic Females	193.5	90	11%	55	13%	35	9%
Hispanic/Latino Males	239.2	18	2%	9	2%	9	2%
Hispanic/Latina Females	*	4	<1%	2	<1%	2	<1%
Asian/Pacific Islander Males	*	3	<1%	3	1%	-	-
Asian/Pacific Islander Females	*	3	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	10	1%	5	1%	5	1%
Total	95.4	836		430		406	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Akron Consortium 6

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	452	68%	213	64%	239	71%
Injection drug use (IDU)	24	4%	8	2%	16	5%
Male-to-male sexual contact and IDU	24	4%	7	2%	17	5%
Heterosexual contact	47	7%	19	6%	28	8%
Other/unknown	121	18%	86	26%	35	10%
Subtotal	668	100%	333	100%	335	100%
Female adult or adolescent						
Injection drug use	15	9%	8	9%	7	10%
Heterosexual contact	93	59%	48	55%	45	64%
Other/unknown	50	32%	32	36%	18	26%
Subtotal	158	100%	88	100%	70	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	100%	9	100%	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	10	100%	9	100%	1	100%
Total	836		430		406	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Akron Consortium 6

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	435	52%	207	48%	228	56%
Injection drug use (IDU) only	21	3%	9	2%	12	3%
Heterosexual contact only	140	17%	67	16%	73	18%
MSM & IDU	22	3%	6	1%	16	4%
IDU & Heterosexual contact	18	2%	7	2%	11	3%
MSM & Heterosexual contact	17	2%	6	1%	11	3%
MSM & IDU & Heterosexual contact	2	<1%	1	<1%	1	<1%
Perinatal exposure	10	1%	9	2%	1	<1%
Other/unknown	171	20%	118	27%	53	13%
Total	836		430		406	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Akron Consortium 6

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	299	72%	131	59%	12	75%	2	67%	1	100%	7	78%
Injection drug use (IDU)	15	4%	8	4%	1	6%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	13	3%	11	5%	-	-	-	-	-	-	-	-
Heterosexual contact	22	5%	23	10%	1	6%	1	33%	-	-	-	-
Other/unknown	67	16%	50	22%	2	13%	-	-	-	-	2	22%
Subtotal	416	100%	223	100%	16	100%	3	100%	1	100%	9	100%
Female adult or adolescent												
Injection drug use	6	10%	8	9%	1	25%	-	-	-	-	-	-
Heterosexual contact	38	61%	50	57%	3	75%	2	67%	-	-	-	-
Other/unknown	18	29%	30	34%	-	-	1	33%	-	-	1	100%
Subtotal	62	100%	88	100%	4	100%	3	100%	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	7	100%	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	7	100%	2	100%	-	-	-	-	-	-
Total	479		318		22		6		1		10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Akron Consortium 6

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	289	60%	125	39%	12	55%	2	33%	1	100%	6	60%
Injection drug use (IDU) only	12	3%	8	3%	1	5%	-	-	-	-	-	-
Heterosexual contact only	60	13%	73	23%	4	18%	3	50%	-	-	-	-
MSM & IDU	13	3%	9	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	9	2%	8	3%	1	5%	-	-	-	-	-	-
MSM & Heterosexual contact	10	2%	6	2%	-	-	-	-	-	-	1	10%
MSM & IDU & Heterosexual contact	-	-	2	1%	-	-	-	-	-	-	-	-
Perinatal exposure	1	0%	7	2%	2	9%	-	-	-	-	-	-
Other/unknown	85	18%	80	25%	2	9%	1	17%	-	-	3	30%
Total	479		318		22		6		1		10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Akron Consortium 6

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	541	81%	575	81%	614	81%	637	81%	676	81%
Females	128	19%	135	19%	143	19%	151	19%	160	19%
Age at end of year										
<13	8	1%	8	1%	8	1%	8	1%	10	1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	20	3%	23	3%	25	3%	25	3%	27	3%
20-24	73	11%	83	12%	98	13%	103	13%	115	14%
25-29	106	16%	114	16%	121	16%	123	16%	131	16%
30-34	127	19%	131	18%	135	18%	144	18%	146	17%
35-39	127	19%	130	18%	132	17%	134	17%	141	17%
40-44	108	16%	112	16%	120	16%	120	15%	126	15%
45-49	55	8%	59	8%	66	9%	71	9%	73	9%
50-54	28	4%	30	4%	31	4%	35	4%	40	5%
55-64	13	2%	16	2%	17	2%	18	2%	20	2%
65+	3	<1%	3	<1%	3	<1%	6	1%	6	1%
Race/Ethnicity										
White, not Hispanic	386	58%	406	57%	434	57%	452	57%	479	57%
Black/African American, not Hispanic	252	38%	271	38%	289	38%	300	38%	318	38%
Hispanic/Latino	18	3%	20	3%	21	3%	21	3%	22	3%
Asian/Pacific Islander	5	1%	5	1%	5	1%	6	1%	6	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	7	1%	7	1%	7	1%	8	1%	10	1%
Race/Sex										
White, not Hispanic Males	342	51%	360	51%	382	50%	395	50%	417	50%
White, not Hispanic Females	44	7%	46	6%	52	7%	57	7%	62	7%
Black/African American, not Hispanic Males	175	26%	189	27%	205	27%	214	27%	228	27%
Black/African American, not Hispanic Females	77	12%	82	12%	84	11%	86	11%	90	11%
Hispanic/Latino Males	14	2%	16	2%	17	2%	17	2%	18	2%
Hispanic/Latina Females	4	1%	4	1%	4	1%	4	1%	4	<1%
Asian/Pacific Islander Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	2	<1%	2	<1%	2	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	7	1%	7	1%	7	1%	8	1%	10	1%
Total	669		710		757		788		836	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Akron Consortium 6

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	360	67%	390	69%	412	68%	427	68%	452	68%
Injection drug use (IDU)	20	4%	20	4%	22	4%	23	4%	24	4%
Male-to-male sexual contact and IDU	22	4%	22	4%	22	4%	24	4%	24	4%
Heterosexual contact	39	7%	42	7%	45	7%	45	7%	47	7%
Other/unknown	94	18%	95	17%	107	18%	112	18%	121	18%
Subtotal	535	100%	569	100%	608	100%	631	100%	668	100%
Female adult or adolescent										
Injection drug use	12	10%	14	11%	15	11%	15	10%	15	9%
Heterosexual contact	80	63%	81	61%	87	62%	91	61%	93	59%
Other/unknown	34	27%	38	29%	39	28%	43	29%	50	32%
Subtotal	126	100%	133	100%	141	100%	149	100%	158	100%
Child (<13 yrs at diagnosis)										
Perinatal	8	100%	8	100%	8	100%	8	100%	10	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	8	100%	8	100%	8	100%	8	100%	10	100%
Total	669		710		757		788		836	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Akron Consortium 6

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	346	52%	375	53%	397	52%	411	52%	435	52%
Injection drug use (IDU) only	19	3%	20	3%	20	3%	21	3%	21	3%
Heterosexual contact only	119	18%	123	17%	132	17%	136	17%	140	17%
MSM & IDU	21	3%	21	3%	21	3%	22	3%	22	3%
IDU & Heterosexual contact	13	2%	14	2%	17	2%	17	2%	18	2%
MSM & Heterosexual contact	14	2%	15	2%	15	2%	16	2%	17	2%
MSM & IDU & Heterosexual contact	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
Perinatal exposure	8	1%	8	1%	8	1%	8	1%	10	1%
Other/unknown	128	19%	133	19%	146	19%	155	20%	171	20%
Total	669		710		757		788		836	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Youngstown Consortium 7

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Youngstown Consortium 7

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	9.9	30	83%	19	83%	8	80%	3	100%
Females	1.9	6	17%	4	17%	2	20%	-	-
Age at diagnosis (yr)									
<13	*	1	3%	1	4%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	2	6%	2	9%	-	-	-	-
20-24	18.8	7	19%	4	17%	2	20%	1	33%
25-29	*	4	11%	3	13%	1	10%	-	-
30-34	*	3	8%	2	9%	1	10%	-	-
35-39	17.6	6	17%	3	13%	2	20%	1	33%
40-44	*	1	3%	1	4%	-	-	-	-
45-49	*	4	11%	3	13%	1	10%	-	-
50-54	12.2	6	17%	3	13%	3	30%	-	-
55-64	*	2	6%	1	4%	-	-	1	33%
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	3.4	18	50%	11	48%	5	50%	2	67%
Black/African American, not Hispanic	22.0	13	36%	8	35%	4	40%	1	33%
Hispanic/Latino	*	4	11%	3	13%	1	10%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	3%	1	4%	-	-	-	-
Race/Sex									
White, not Hispanic Males	5.9	15	42%	8	35%	5	50%	2	67%
White, not Hispanic Females	*	3	8%	3	13%	-	-	-	-
Black/African American, not Hispanic Males	38.2	11	31%	8	35%	2	20%	1	33%
Black/African American, not Hispanic Females	*	2	6%	-	-	2	20%	-	-
Hispanic/Latino Males	*	4	11%	3	13%	1	10%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	3%	1	4%	-	-	-	-
Total	5.8	36		23		10		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Youngstown Consortium 7

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	15	52%	10	56%	4	50%	1	33%
Injection drug use (IDU)	2	7%	2	11%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	3%	1	6%	-	-	-	-
Other/unknown	11	38%	5	28%	4	50%	2	67%
Subtotal	29	100%	18	100%	8	100%	3	100%
Female adult or adolescent								
Injection drug use	1	17%	1	25%	-	-	-	-
Heterosexual contact	1	17%	-	-	1	50%	-	-
Other/unknown	4	67%	3	75%	1	50%	-	-
Subtotal	6	100%	4	100%	2	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	36		23		10		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	39%	9	39%	4	40%	1	33%
Injection drug use (IDU) only	3	8%	3	13%	-	-	-	-
Heterosexual contact only	2	6%	1	4%	1	10%	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	3%	1	4%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	16	44%	9	39%	5	50%	2	67%
Total	36		23		10		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B
Youngstown Consortium 7**

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	7	47%	6	60%	2	50%	-	-	-	-	-	-
Injection drug use (IDU)	1	7%	-	-	1	25%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	10%	-	-	-	-	-	-	-	-
Other/unknown	7	47%	3	30%	1	25%	-	-	-	-	-	-
Subtotal	15	100%	10	100%	4	100%	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	1	33%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	50%	-	-	-	-	-	-	-	-
Other/unknown	2	67%	1	50%	-	-	-	-	-	-	1	100%
Subtotal	3	100%	2	100%	-	-	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	18		13		4		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B
Youngstown Consortium 7**

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	39%	5	38%	2	50%	-	-	-	-	-	-
Injection drug use (IDU) only	2	11%	-	-	1	25%	-	-	-	-	-	-
Heterosexual contact only	-	-	2	15%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	1	8%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	9	50%	5	38%	1	25%	-	-	-	-	1	100%
Total	18		13		4		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Youngstown Consortium 7**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	5.6	17	59%	8.0	24	71%	8.6	26	90%	9.0	27	75%
Females	3.8	12	41%	3.2	10	29%	*	3	10%	2.8	9	25%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	3%	*	4	14%	12.9	5	14%
20-24	*	3	10%	18.8	7	21%	*	4	14%	18.8	7	19%
25-29	15.3	5	17%	*	4	12%	*	3	10%	24.5	8	22%
30-34	*	2	7%	17.7	6	18%	*	-	-	*	2	6%
35-39	14.7	5	17%	14.7	5	15%	*	3	10%	*	3	8%
40-44	*	2	7%	*	3	9%	12.9	5	17%	*	3	8%
45-49	19.2	8	28%	*	2	6%	*	4	14%	*	1	3%
50-54	*	3	10%	*	3	9%	*	3	10%	*	3	8%
55-64	*	1	3%	*	3	9%	*	3	10%	*	3	8%
65+	*	-	-	*	-	-	*	-	-	*	1	3%
Race/Ethnicity												
White, not Hispanic	2.5	13	45%	3.4	18	53%	2.1	11	38%	2.3	12	33%
Black/African American, not Hispanic	22.0	13	45%	20.3	12	35%	25.4	15	52%	32.1	19	53%
Hispanic/Latino	*	2	7%	*	2	6%	*	1	3%	*	2	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	3%
Unknown	*	1	3%	*	2	6%	*	2	7%	*	2	6%
Race/Sex												
White, not Hispanic Males	3.5	9	31%	5.5	14	41%	3.9	10	34%	3.1	8	22%
White, not Hispanic Females	*	4	14%	*	4	12%	*	1	3%	*	4	11%
Black/African American, not Hispanic Males	20.8	6	21%	24.3	7	21%	45.1	13	45%	55.5	16	44%
Black/African American, not Hispanic Females	23.1	7	24%	16.5	5	15%	*	2	7%	*	3	8%
Hispanic/Latino Males	*	1	3%	*	1	3%	*	1	3%	*	1	3%
Hispanic/Latina Females	*	1	3%	*	1	3%	*	-	-	*	1	3%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	3%	*	2	6%	*	2	7%	*	2	6%
Total	4.7	29		5.5	34		4.7	29		5.8	36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Youngstown Consortium 7

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	10	59%	15	63%	14	54%	14	52%
Injection drug use (IDU)	2	12%	-	-	3	12%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	2	12%	5	21%	3	12%	2	7%
Other/unknown	3	18%	4	17%	6	23%	11	41%
Subtotal	17	100%	24	100%	26	100%	27	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	1	33%	1	11%
Heterosexual contact	7	58%	8	80%	1	33%	3	33%
Other/unknown	5	42%	2	20%	1	33%	5	56%
Subtotal	12	100%	10	100%	3	100%	9	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	29		34		29		36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Youngstown Consortium 7

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	34%	12	35%	13	45%	14	39%
Injection drug use (IDU) only	2	7%	-	-	3	10%	1	3%
Heterosexual contact only	9	31%	13	38%	4	14%	5	14%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	3%	-	-
MSM & Heterosexual contact	-	-	3	9%	1	3%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	8	28%	6	18%	7	24%	16	44%
Total	29		34		29		36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Youngstown Consortium 7

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	6	40%	12	67%	16	89%	18	72%	19	83%
Females	9	60%	6	33%	2	11%	7	28%	4	17%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	1	4%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	1	6%	2	11%	5	20%	2	9%
20-24	3	20%	5	28%	3	17%	6	24%	4	17%
25-29	2	13%	3	17%	3	17%	6	24%	3	13%
30-34	1	7%	3	17%	-	-	1	4%	2	9%
35-39	3	20%	3	17%	3	17%	3	12%	3	13%
40-44	-	-	1	6%	3	17%	-	-	1	4%
45-49	4	27%	-	-	1	6%	-	-	3	13%
50-54	2	13%	1	6%	2	11%	2	8%	3	13%
55-64	-	-	1	6%	1	6%	1	4%	1	4%
65+	-	-	-	-	-	-	1	4%	-	-
Race/Ethnicity										
White, not Hispanic	4	27%	8	44%	6	33%	7	28%	11	48%
Black/African American, not Hispanic	10	67%	7	39%	9	50%	14	56%	8	35%
Hispanic/Latino	1	7%	2	11%	1	6%	2	8%	3	13%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	1	4%	-	-
Unknown	-	-	1	6%	2	11%	1	4%	1	4%
Race/Sex										
White, not Hispanic Males	2	13%	6	33%	5	28%	4	16%	8	35%
White, not Hispanic Females	2	13%	2	11%	1	6%	3	12%	3	13%
Black/African American, not Hispanic Males	4	27%	4	22%	8	44%	11	44%	8	35%
Black/African American, not Hispanic Females	6	40%	3	17%	1	6%	3	12%	-	-
Hispanic/Latino Males	-	-	1	6%	1	6%	1	4%	3	13%
Hispanic/Latina Females	1	7%	1	6%	-	-	1	4%	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	1	4%	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	6%	2	11%	1	4%	1	4%
Total	15		18		18		25		23	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Youngstown Consortium 7

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	4	67%	7	58%	9	56%	12	67%	10	56%
Injection drug use (IDU)	1	17%	-	-	1	6%	-	-	2	11%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	4	33%	2	13%	1	6%	1	6%
Other/unknown	1	17%	1	8%	4	25%	5	28%	5	28%
Subtotal	6	100%	12	100%	16	100%	18	100%	18	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	1	50%	-	-	1	25%
Heterosexual contact	5	56%	6	100%	-	-	3	43%	-	-
Other/unknown	4	44%	-	-	1	50%	4	57%	3	75%
Subtotal	9	100%	6	100%	2	100%	7	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	100%
Subtotal	-	-	-	-	-	-	-	-	1	100%
Total	15		18		18		25		23	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	4	27%	5	28%	8	44%	12	48%	9	39%
Injection drug use (IDU) only	1	7%	-	-	2	11%	-	-	3	13%
Heterosexual contact only	5	33%	10	56%	2	11%	4	16%	1	4%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	2	11%	1	6%	-	-	1	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	33%	1	6%	5	28%	9	36%	9	39%
Total	15		18		18		25		23	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Youngstown Consortium 7

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	15	75%	11	65%	12	71%	19	79%	13	87%	497	77%
Females	5	25%	6	35%	5	29%	5	21%	2	13%	149	23%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	-	-	-	-	4	24%	1	4%	-	-	16	2%
20-24	1	5%	-	-	-	-	4	17%	3	20%	41	6%
25-29	5	25%	1	6%	1	6%	3	13%	1	7%	101	16%
30-34	1	5%	6	35%	1	6%	3	13%	1	7%	126	20%
35-39	2	10%	3	18%	2	12%	-	-	3	20%	128	20%
40-44	5	25%	2	12%	3	18%	5	21%	-	-	94	15%
45-49	3	15%	1	6%	4	24%	3	13%	2	13%	60	9%
50-54	2	10%	2	12%	1	6%	2	8%	3	20%	32	5%
55-64	1	5%	2	12%	1	6%	3	13%	2	13%	35	5%
65+	-	-	-	-	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	11	55%	12	71%	6	35%	11	46%	8	53%	363	56%
Black/African American, not Hispanic	6	30%	4	24%	9	53%	10	42%	6	40%	244	38%
Hispanic/Latino	1	5%	-	-	1	6%	1	4%	1	7%	30	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	5%	-	-	-	-	-	-	-	-	1	<1%
Unknown	1	5%	1	6%	1	6%	2	8%	-	-	8	1%
Race/Sex												
White, not Hispanic Males	8	40%	9	53%	5	29%	8	33%	8	53%	312	48%
White, not Hispanic Females	3	15%	3	18%	1	6%	3	13%	-	-	51	8%
Black/African American, not Hispanic Males	4	20%	1	6%	5	29%	10	42%	4	27%	159	25%
Black/African American, not Hispanic Females	2	10%	3	18%	4	24%	-	-	2	13%	85	13%
Hispanic/Latino Males	1	5%	-	-	1	6%	-	-	1	7%	20	3%
Hispanic/Latina Females	-	-	-	-	-	-	1	4%	-	-	10	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	5%	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	1	6%	1	6%	2	8%	-	-	8	1%
Total	20		17		17		24		15		646	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Youngstown Consortium 7

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	10	67%	7	64%	6	50%	8	42%	5	38%	281	57%
Injection drug use (IDU)	2	13%	-	-	2	17%	-	-	1	8%	46	9%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	33	7%
Heterosexual contact	2	13%	2	18%	1	8%	4	21%	-	-	44	9%
Other/unknown	1	7%	2	18%	3	25%	7	37%	7	54%	89	18%
Subtotal	15	100%	11	100%	12	100%	19	100%	13	100%	493	100%
Female adult or adolescent												
Injection drug use	2	40%	1	17%	1	20%	2	40%	-	-	33	22%
Heterosexual contact	3	60%	2	33%	4	80%	1	20%	1	50%	72	49%
Other/unknown	-	-	3	50%	-	-	2	40%	1	50%	42	29%
Subtotal	5	100%	6	100%	5	100%	5	100%	2	100%	147	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	33%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	20		17		17		24		15		646	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	9	45%	6	35%	6	35%	6	25%	5	33%	264	41%
Injection drug use (IDU) only	3	15%	-	-	2	12%	2	8%	-	-	43	7%
Heterosexual contact only	5	25%	4	24%	5	29%	5	21%	1	7%	116	18%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	27	4%
IDU & Heterosexual contact	1	5%	1	6%	1	6%	-	-	1	7%	35	5%
MSM & Heterosexual contact	1	5%	1	6%	-	-	2	8%	-	-	17	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	6	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	4	1%
Other/unknown	1	5%	5	29%	3	18%	9	38%	8	53%	134	21%
Total	20		17		17		24		15		646	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Youngstown Consortium 7

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Youngstown Consortium 7

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	131.0	395	69%	202	68%	193	69%
Females	57.3	181	31%	96	32%	85	31%
Age at end of year							
<13	14.4	13	2%	9	3%	4	1%
13-14	*	1	<1%	-	-	1	<1%
15-19	62.0	24	4%	15	5%	9	3%
20-24	182.6	68	12%	52	17%	16	6%
25-29	314.8	103	18%	53	18%	50	18%
30-34	248.1	84	15%	38	13%	46	17%
35-39	284.5	97	17%	45	15%	52	19%
40-44	209.1	81	14%	33	11%	48	17%
45-49	115.3	48	8%	24	8%	24	9%
50-54	67.0	33	6%	19	6%	14	5%
55-64	20.5	19	3%	6	2%	13	5%
65+	4.4	5	1%	4	1%	1	<1%
Race/Ethnicity							
White, not Hispanic	48.0	253	44%	124	42%	129	46%
Black/African American, not Hispanic	429.5	254	44%	128	43%	126	45%
Hispanic/Latino	222.4	38	7%	23	8%	15	5%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	29	5%	22	7%	7	3%
Race/Sex							
White, not Hispanic Males	75.7	194	34%	92	31%	102	37%
White, not Hispanic Females	21.8	59	10%	32	11%	27	10%
Black/African American, not Hispanic Males	551.6	159	28%	83	28%	76	27%
Black/African American, not Hispanic Females	313.4	95	16%	45	15%	50	18%
Hispanic/Latino Males	224.9	21	4%	11	4%	10	4%
Hispanic/Latina Females	219.4	17	3%	12	4%	5	2%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	29	5%	22	7%	7	3%
Total	93.3	576		298		278	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Youngstown Consortium 7

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	208	53%	97	49%	111	58%
Injection drug use (IDU)	25	6%	7	4%	18	9%
Male-to-male sexual contact and IDU	11	3%	4	2%	7	4%
Heterosexual contact	49	13%	22	11%	27	14%
Other/unknown	96	25%	68	34%	28	15%
Subtotal	389	100%	198	100%	191	100%
Female adult or adolescent						
Injection drug use	18	10%	5	5%	13	16%
Heterosexual contact	97	56%	45	49%	52	63%
Other/unknown	59	34%	41	45%	18	22%
Subtotal	174	100%	91	100%	83	100%
Child (<13 yrs at diagnosis)						
Perinatal	11	85%	8	89%	3	75%
Other/unknown	2	15%	1	11%	1	25%
Subtotal	13	100%	9	100%	4	100%
Total	576		298		278	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	189	33%	90	30%	99	36%
Injection drug use (IDU) only	31	5%	11	4%	20	7%
Heterosexual contact only	146	25%	67	22%	79	28%
MSM & IDU	8	1%	2	1%	6	2%
IDU & Heterosexual contact	12	2%	1	<1%	11	4%
MSM & Heterosexual contact	19	3%	7	2%	12	4%
MSM & IDU & Heterosexual contact	3	1%	2	1%	1	<1%
Perinatal exposure	11	2%	8	3%	3	1%
Other/unknown	157	27%	110	37%	47	17%
Total	576		298		278	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Youngstown Consortium 7

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	113	59%	85	54%	7	35%	-	-	-	-	3	16%
Injection drug use (IDU)	11	6%	9	6%	4	20%	-	-	1	50%	-	-
Male-to-male sexual contact and IDU	7	4%	4	3%	-	-	-	-	-	-	-	-
Heterosexual contact	20	10%	21	13%	5	25%	-	-	-	-	3	16%
Other/unknown	41	21%	37	24%	4	20%	-	-	1	50%	13	68%
Subtotal	192	100%	156	100%	20	100%	-	-	2	100%	19	100%
Female adult or adolescent												
Injection drug use	7	12%	11	12%	-	-	-	-	-	-	-	-
Heterosexual contact	37	64%	51	55%	9	60%	-	-	-	-	-	-
Other/unknown	14	24%	31	33%	6	40%	-	-	-	-	8	100%
Subtotal	58	100%	93	100%	15	100%	-	-	-	-	8	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	100%	4	80%	3	100%	-	-	-	-	1	50%
Other/unknown	-	-	1	20%	-	-	-	-	-	-	1	50%
Subtotal	3	100%	5	100%	3	100%	-	-	-	-	2	100%
Total	253		254		38		0		2		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	104	41%	77	30%	6	16%	-	-	-	-	2	7%
Injection drug use (IDU) only	13	5%	15	6%	2	5%	-	-	1	50%	-	-
Heterosexual contact only	57	23%	72	28%	14	37%	-	-	-	-	3	10%
MSM & IDU	5	2%	3	1%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	5	2%	5	2%	2	5%	-	-	-	-	-	-
MSM & Heterosexual contact	9	4%	8	3%	1	3%	-	-	-	-	1	3%
MSM & IDU & Heterosexual contact	2	1%	1	0%	-	-	-	-	-	-	-	-
Perinatal exposure	3	1%	4	2%	3	8%	-	-	-	-	1	3%
Other/unknown	55	22%	69	27%	10	26%	-	-	1	50%	22	76%
Total	253		254		38		0		2		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Youngstown Consortium 7

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	307	67%	325	67%	345	67%	370	68%	395	69%
Females	149	33%	160	33%	169	33%	172	32%	181	31%
Age at end of year										
<13	13	3%	13	3%	13	3%	13	2%	13	2%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	14	3%	14	3%	15	3%	19	4%	24	4%
20-24	46	10%	50	10%	57	11%	61	11%	68	12%
25-29	83	18%	88	18%	92	18%	95	18%	103	18%
30-34	73	16%	76	16%	82	16%	82	15%	84	15%
35-39	82	18%	87	18%	92	18%	95	18%	97	17%
40-44	69	15%	71	15%	74	14%	78	14%	81	14%
45-49	37	8%	43	9%	43	8%	47	9%	48	8%
50-54	22	5%	25	5%	27	5%	30	6%	33	6%
55-64	12	3%	13	3%	14	3%	17	3%	19	3%
65+	4	1%	4	1%	4	1%	4	1%	5	1%
Race/Ethnicity										
White, not Hispanic	205	45%	218	45%	232	45%	242	45%	253	44%
Black/African American, not Hispanic	196	43%	210	43%	221	43%	236	44%	254	44%
Hispanic/Latino	32	7%	33	7%	35	7%	36	7%	38	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
Unknown	22	5%	23	5%	25	5%	27	5%	29	5%
Race/Sex										
White, not Hispanic Males	159	35%	168	35%	178	35%	187	35%	194	34%
White, not Hispanic Females	46	10%	50	10%	54	11%	55	10%	59	10%
Black/African American, not Hispanic Males	116	25%	124	26%	131	25%	144	27%	159	28%
Black/African American, not Hispanic Females	80	18%	86	18%	90	18%	92	17%	95	16%
Hispanic/Latino Males	18	4%	18	4%	19	4%	20	4%	21	4%
Hispanic/Latina Females	14	3%	15	3%	16	3%	16	3%	17	3%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	22	5%	23	5%	25	5%	27	5%	29	5%
Total	456		485		514		542		576	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Youngstown Consortium 7

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	158	52%	167	52%	180	53%	194	53%	208	53%
Injection drug use (IDU)	20	7%	22	7%	22	6%	25	7%	25	6%
Male-to-male sexual contact and IDU	10	3%	11	3%	11	3%	11	3%	11	3%
Heterosexual contact	37	12%	39	12%	44	13%	47	13%	49	13%
Other/unknown	76	25%	80	25%	82	24%	87	24%	96	25%
Subtotal	301	100%	319	100%	339	100%	364	100%	389	100%
Female adult or adolescent										
Injection drug use	16	11%	16	10%	16	10%	17	10%	18	10%
Heterosexual contact	78	55%	85	56%	93	57%	94	57%	97	56%
Other/unknown	48	34%	52	34%	53	33%	54	33%	59	34%
Subtotal	142	100%	153	100%	162	100%	165	100%	174	100%
Child (<13 yrs at diagnosis)										
Perinatal	11	85%	11	85%	11	85%	11	85%	11	85%
Other/unknown	2	15%	2	15%	2	15%	2	15%	2	15%
Subtotal	13	100%	13	100%	13	100%	13	100%	13	100%
Total	456		485		514		542		576	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Youngstown Consortium 7

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	144	32%	152	31%	162	32%	175	32%	189	33%
Injection drug use (IDU) only	25	5%	27	6%	27	5%	30	6%	31	5%
Heterosexual contact only	115	25%	124	26%	137	27%	141	26%	146	25%
MSM & IDU	7	2%	8	2%	8	2%	8	1%	8	1%
IDU & Heterosexual contact	11	2%	11	2%	11	2%	12	2%	12	2%
MSM & Heterosexual contact	14	3%	15	3%	18	4%	19	4%	19	3%
MSM & IDU & Heterosexual contact	3	1%	3	1%	3	1%	3	1%	3	1%
Perinatal exposure	11	2%	11	2%	11	2%	11	2%	11	2%
Other/unknown	126	28%	134	28%	137	27%	143	26%	157	27%
Total	456		485		514		542		576	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B Canton
Consortium 8

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Canton Consortium 8

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	6.4	21	84%	17	81%	3	100%	1	100%
Females	*	4	16%	4	19%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	1	4%	1	5%	-	-	-	-
20-24	*	2	8%	2	10%	-	-	-	-
25-29	13.4	5	20%	5	24%	-	-	-	-
30-34	*	4	16%	2	10%	2	67%	-	-
35-39	*	1	4%	1	5%	-	-	-	-
40-44	*	4	16%	4	19%	-	-	-	-
45-49	11.1	5	20%	4	19%	-	-	1	100%
50-54	*	2	8%	2	10%	-	-	-	-
55-64	*	1	4%	-	-	1	33%	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	3.1	19	76%	17	81%	1	33%	1	100%
Black/African American, not Hispanic	*	4	16%	3	14%	1	33%	-	-
Hispanic/Latino	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	2	8%	1	5%	1	33%	-	-
Race/Sex									
White, not Hispanic Males	5.0	15	60%	13	62%	1	33%	1	100%
White, not Hispanic Females	*	4	16%	4	19%	-	-	-	-
Black/African American, not Hispanic Males	*	4	16%	3	14%	1	33%	-	-
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	2	8%	1	5%	1	33%	-	-
Total	3.7	25		21		3		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Canton Consortium 8

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	15	71%	14	82%	1	33%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	10%	-	-	1	33%	1	100%
Heterosexual contact	1	5%	1	6%	-	-	-	-
Other/unknown	3	14%	2	12%	1	33%	-	-
Subtotal	21	100%	17	100%	3	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	1	25%	1	25%	-	-	-	-
Other/unknown	3	75%	3	75%	-	-	-	-
Subtotal	4	100%	4	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	25		21		3		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Canton Consortium 8

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	15	60%	14	67%	1	33%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	2	8%	2	10%	-	-	-	-
MSM & IDU	1	4%	-	-	1	33%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	4%	-	-	-	-	1	100%
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	6	24%	5	24%	1	33%	-	-
Total	25		21		3		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B
Canton Consortium 8**

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	11	73%	3	75%	-	-	-	-	-	-	1	50%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	7%	1	25%	-	-	-	-	-	-	-	-
Heterosexual contact	1	7%	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	13%	-	-	-	-	-	-	-	-	1	50%
Subtotal	15	100%	4	100%	-	-	-	-	-	-	2	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	25%	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	75%	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	19		4		0		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Canton Consortium 8

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	11	58%	3	75%	-	-	-	-	-	-	1	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	11%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	1	25%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	26%	-	-	-	-	-	-	-	-	1	50%
Total	19		4		0		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Canton Consortium 8**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	6.7	22	88%	8.2	27	84%	9.2	30	88%	7.9	26	84%
Females	*	3	12%	1.5	5	16%	*	4	12%	1.5	5	16%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	3	12%	*	1	3%	*	4	12%	*	2	6%
20-24	18.9	8	32%	26.0	11	34%	*	4	12%	21.3	9	29%
25-29	*	1	4%	*	1	3%	*	3	9%	16.1	6	19%
30-34	*	1	4%	*	3	9%	*	4	12%	*	2	6%
35-39	*	3	12%	*	4	13%	13.7	5	15%	*	2	6%
40-44	*	3	12%	*	3	9%	14.3	6	18%	*	1	3%
45-49	*	3	12%	*	4	13%	*	1	3%	*	2	6%
50-54	*	2	8%	*	2	6%	*	4	12%	*	2	6%
55-64	*	1	4%	*	2	6%	*	3	9%	*	4	13%
65+	*	-	-	*	1	3%	*	-	-	*	1	3%
Race/Ethnicity												
White, not Hispanic	2.1	13	52%	3.6	22	69%	3.4	21	62%	3.3	20	65%
Black/African American, not Hispanic	31.7	10	40%	22.2	7	22%	31.7	10	29%	28.5	9	29%
Hispanic/Latino	*	1	4%	*	-	-	*	1	3%	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	1	3%	*	-	-
Unknown	*	1	4%	*	3	9%	*	1	3%	*	2	6%
Race/Sex												
White, not Hispanic Males	4.0	12	48%	6.7	20	63%	6.0	18	53%	5.7	17	55%
White, not Hispanic Females	*	1	4%	*	2	6%	*	3	9%	*	3	10%
Black/African American, not Hispanic Males	51.0	8	32%	31.9	5	16%	57.4	9	26%	44.6	7	23%
Black/African American, not Hispanic Females	*	2	8%	*	2	6%	*	1	3%	*	2	6%
Hispanic/Latino Males	*	1	4%	*	-	-	*	1	3%	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	1	3%	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	3	9%	*	1	3%	*	2	6%
Total	3.7	25		4.8	32		5.1	34		4.6	31	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Canton Consortium 8

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	19	86%	18	67%	22	73%	20	77%
Injection drug use (IDU)	1	5%	2	7%	-	-	1	4%
Male-to-male sexual contact and IDU	-	-	1	4%	2	7%	1	4%
Heterosexual contact	-	-	3	11%	3	10%	1	4%
Other/unknown	2	9%	3	11%	3	10%	3	12%
Subtotal	22	100%	27	100%	30	100%	26	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	2	40%
Heterosexual contact	3	100%	4	80%	2	50%	3	60%
Other/unknown	-	-	1	20%	2	50%	-	-
Subtotal	3	100%	5	100%	4	100%	5	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	25		32		34		31	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Canton Consortium 8

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	17	68%	17	53%	19	56%	18	58%
Injection drug use (IDU) only	-	-	2	6%	-	-	1	3%
Heterosexual contact only	3	12%	7	22%	5	15%	4	13%
MSM & IDU	-	-	1	3%	1	3%	-	-
IDU & Heterosexual contact	1	4%	-	-	-	-	2	6%
MSM & Heterosexual contact	2	8%	1	3%	3	9%	2	6%
MSM & IDU & Heterosexual contact	-	-	-	-	1	3%	1	3%
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	2	8%	4	13%	5	15%	3	10%
Total	25		32		34		31	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Canton Consortium 8

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	12	92%	18	82%	21	88%	19	90%	17	81%
Females	1	8%	4	18%	3	13%	2	10%	4	19%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	23%	1	5%	4	17%	2	10%	1	5%
20-24	6	46%	11	50%	3	13%	8	38%	2	10%
25-29	1	8%	1	5%	1	4%	4	19%	5	24%
30-34	-	-	2	9%	4	17%	2	10%	2	10%
35-39	1	8%	2	9%	4	17%	1	5%	1	5%
40-44	-	-	-	-	2	8%	1	5%	4	19%
45-49	1	8%	3	14%	1	4%	-	-	4	19%
50-54	-	-	-	-	4	17%	1	5%	2	10%
55-64	1	8%	1	5%	1	4%	2	10%	-	-
65+	-	-	1	5%	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	6	46%	15	68%	16	67%	17	81%	17	81%
Black/African American, not Hispanic	6	46%	5	23%	5	21%	3	14%	3	14%
Hispanic/Latino	1	8%	-	-	1	4%	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	1	4%	-	-	-	-
Unknown	-	-	2	9%	1	4%	1	5%	1	5%
Race/Sex										
White, not Hispanic Males	6	46%	13	59%	13	54%	15	71%	13	62%
White, not Hispanic Females	-	-	2	9%	3	13%	2	10%	4	19%
Black/African American, not Hispanic Males	5	38%	4	18%	5	21%	3	14%	3	14%
Black/African American, not Hispanic Females	1	8%	1	5%	-	-	-	-	-	-
Hispanic/Latino Males	1	8%	-	-	1	4%	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	1	4%	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	9%	1	4%	1	5%	1	5%
Total	13		22		24		21		21	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Canton Consortium 8

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	10	83%	15	83%	14	67%	15	79%	14	82%
Injection drug use (IDU)	1	8%	1	6%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	2	10%	1	5%	-	-
Heterosexual contact	-	-	1	6%	2	10%	-	-	1	6%
Other/unknown	1	8%	1	6%	3	14%	3	16%	2	12%
Subtotal	12	100%	18	100%	21	100%	19	100%	17	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	100%	4	100%	2	67%	2	100%	1	25%
Other/unknown	-	-	-	-	1	33%	-	-	3	75%
Subtotal	1	100%	4	100%	3	100%	2	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	13		22		24		21		21	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Canton Consortium 8

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	9	69%	14	64%	13	54%	14	67%	14	67%
Injection drug use (IDU) only	-	-	1	5%	-	-	-	-	-	-
Heterosexual contact only	1	8%	5	23%	4	17%	2	10%	2	10%
MSM & IDU	-	-	-	-	1	4%	-	-	-	-
IDU & Heterosexual contact	1	8%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	8%	1	5%	1	4%	1	5%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	1	4%	1	5%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	8%	1	5%	4	17%	3	14%	5	24%
Total	13		22		24		21		21	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Canton Consortium 8

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	9	75%	12	75%	9	69%	13	76%	5	100%	373	83%
Females	3	25%	4	25%	4	31%	4	24%	-	-	74	17%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	6	1%
20-24	1	8%	2	13%	1	8%	3	18%	-	-	46	10%
25-29	-	-	-	-	3	23%	3	18%	-	-	68	15%
30-34	1	8%	4	25%	2	15%	1	6%	2	40%	89	20%
35-39	2	17%	1	6%	1	8%	1	6%	1	20%	74	17%
40-44	3	25%	3	19%	4	31%	2	12%	-	-	65	15%
45-49	3	25%	2	13%	-	-	3	18%	1	20%	38	9%
50-54	2	17%	2	13%	-	-	1	6%	-	-	22	5%
55-64	-	-	2	13%	2	15%	2	12%	1	20%	25	6%
65+	-	-	-	-	-	-	1	6%	-	-	8	2%
Race/Ethnicity												
White, not Hispanic	9	75%	13	81%	5	38%	6	35%	2	40%	329	74%
Black/African American, not Hispanic	3	25%	2	13%	7	54%	9	53%	2	40%	102	23%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	10	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	6%	1	8%	2	12%	1	20%	6	1%
Race/Sex												
White, not Hispanic Males	7	58%	9	56%	5	38%	5	29%	2	40%	285	64%
White, not Hispanic Females	2	17%	4	25%	-	-	1	6%	-	-	44	10%
Black/African American, not Hispanic Males	2	17%	2	13%	3	23%	6	35%	2	40%	74	17%
Black/African American, not Hispanic Females	1	8%	-	-	4	31%	3	18%	-	-	28	6%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	8	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	6%	1	8%	2	12%	1	20%	6	1%
Total	12		16		13		17		5		447	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Canton Consortium 8

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	8	89%	4	33%	8	89%	9	69%	2	40%	262	71%
Injection drug use (IDU)	-	-	1	8%	-	-	1	8%	-	-	15	4%
Male-to-male sexual contact and IDU	-	-	1	8%	-	-	-	-	2	40%	19	5%
Heterosexual contact	-	-	2	17%	-	-	2	15%	-	-	20	5%
Other/unknown	1	11%	4	33%	1	11%	1	8%	1	20%	55	15%
Subtotal	9	100%	12	100%	9	100%	13	100%	5	100%	371	100%
Female adult or adolescent												
Injection drug use	-	-	1	25%	-	-	2	50%	-	-	14	20%
Heterosexual contact	3	100%	3	75%	1	25%	2	50%	-	-	41	59%
Other/unknown	-	-	-	-	3	75%	-	-	-	-	15	21%
Subtotal	3	100%	4	100%	4	100%	4	100%	-	-	70	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	12		16		13		17		5		447	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Canton Consortium 8

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	7	58%	4	25%	7	54%	7	41%	2	40%	251	56%
Injection drug use (IDU) only	-	-	1	6%	-	-	1	6%	-	-	16	4%
Heterosexual contact only	3	25%	5	31%	1	8%	4	24%	-	-	61	14%
MSM & IDU	-	-	1	6%	-	-	-	-	1	20%	13	3%
IDU & Heterosexual contact	-	-	1	6%	-	-	2	12%	-	-	13	3%
MSM & Heterosexual contact	1	8%	-	-	1	8%	2	12%	-	-	11	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	20%	6	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	6	1%
Other/unknown	1	8%	4	25%	4	31%	1	6%	1	20%	70	16%
Total	12		16		13		17		5		447	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Canton Consortium 8

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Canton Consortium 8

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	107.8	353	81%	191	81%	162	81%
Females	24.6	84	19%	45	19%	39	19%
Age at end of year							
<13	4.5	5	1%	3	1%	2	1%
13-14	*	-	-	-	-	-	-
15-19	41.6	19	4%	15	6%	4	2%
20-24	177.3	75	17%	51	22%	24	12%
25-29	168.8	63	14%	27	11%	36	18%
30-34	210.6	80	18%	41	17%	39	19%
35-39	180.7	66	15%	38	16%	28	14%
40-44	131.0	55	13%	22	9%	33	16%
45-49	69.1	31	7%	16	7%	15	7%
50-54	43.3	22	5%	12	5%	10	5%
55-64	19.4	18	4%	10	4%	8	4%
65+	*	3	1%	1	<1%	2	1%
Race/Ethnicity							
White, not Hispanic	49.3	301	69%	171	72%	130	65%
Black/African American, not Hispanic	351.6	111	25%	53	22%	58	29%
Hispanic/Latino	108.4	12	3%	4	2%	8	4%
Asian/Pacific Islander	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	10	2%	6	3%	4	2%
Race/Sex							
White, not Hispanic Males	84.0	250	57%	141	60%	109	54%
White, not Hispanic Females	16.3	51	12%	30	13%	21	10%
Black/African American, not Hispanic Males	516.3	81	19%	39	17%	42	21%
Black/African American, not Hispanic Females	188.9	30	7%	14	6%	16	8%
Hispanic/Latino Males	166.4	10	2%	4	2%	6	3%
Hispanic/Latina Females	*	2	<1%	-	-	2	1%
Asian/Pacific Islander Males	*	1	<1%	1	<1%	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	10	2%	6	3%	4	2%
Total	65.3	437		236		201	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Canton Consortium 8

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	252	71%	132	69%	120	74%
Injection drug use (IDU)	7	2%	5	3%	2	1%
Male-to-male sexual contact and IDU	12	3%	5	3%	7	4%
Heterosexual contact	20	6%	7	4%	13	8%
Other/unknown	62	18%	42	22%	20	12%
Subtotal	353	100%	191	100%	162	100%
Female adult or adolescent						
Injection drug use	8	10%	4	10%	4	11%
Heterosexual contact	48	61%	23	55%	25	68%
Other/unknown	23	29%	15	36%	8	22%
Subtotal	79	100%	42	100%	37	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	80%	2	67%	2	100%
Other/unknown	1	20%	1	33%	-	-
Subtotal	5	100%	3	100%	2	100%
Total	437		236		201	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Canton Consortium 8

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	240	55%	127	54%	113	56%
Injection drug use (IDU) only	6	1%	3	1%	3	1%
Heterosexual contact only	68	16%	30	13%	38	19%
MSM & IDU	7	2%	3	1%	4	2%
IDU & Heterosexual contact	9	2%	6	3%	3	1%
MSM & Heterosexual contact	12	3%	5	2%	7	3%
MSM & IDU & Heterosexual contact	5	1%	2	1%	3	1%
Perinatal exposure	4	1%	2	1%	2	1%
Other/unknown	86	20%	58	25%	28	14%
Total	437		236		201	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Canton Consortium 8

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	190	76%	51	63%	6	60%	1	100%	-	-	4	44%
Injection drug use (IDU)	4	2%	3	4%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	8	3%	4	5%	-	-	-	-	-	-	-	-
Heterosexual contact	7	3%	7	9%	2	20%	-	-	1	50%	3	33%
Other/unknown	41	16%	16	20%	2	20%	-	-	1	50%	2	22%
Subtotal	250	100%	81	100%	10	100%	1	100%	2	100%	9	100%
Female adult or adolescent												
Injection drug use	5	10%	3	11%	-	-	-	-	-	-	-	-
Heterosexual contact	29	59%	16	59%	2	100%	-	-	-	-	1	100%
Other/unknown	15	31%	8	30%	-	-	-	-	-	-	-	-
Subtotal	49	100%	27	100%	2	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	2	67%	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	33%	-	-	-	-	-	-	-	-
Subtotal	2	100%	3	100%	-	-	-	-	-	-	-	-
Total	301		111		12		1		2		10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Canton Consortium 8

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	183	61%	47	42%	6	50%	1	100%	-	-	3	30%
Injection drug use (IDU) only	3	1%	3	3%	-	-	-	-	-	-	-	-
Heterosexual contact only	36	12%	23	21%	4	33%	-	-	1	50%	4	40%
MSM & IDU	5	2%	2	2%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	6	2%	3	3%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	7	2%	4	4%	-	-	-	-	-	-	1	10%
MSM & IDU & Heterosexual contact	3	1%	2	2%	-	-	-	-	-	-	-	-
Perinatal exposure	2	1%	2	2%	-	-	-	-	-	-	-	-
Other/unknown	56	19%	25	23%	2	17%	-	-	1	50%	2	20%
Total	301		111		12		1		2		10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Canton Consortium 8

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	253	79%	274	80%	299	80%	328	81%	353	81%
Females	67	21%	70	20%	75	20%	79	19%	84	19%
Age at end of year										
<13	5	2%	5	1%	5	1%	5	1%	5	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	9	3%	12	3%	13	3%	17	4%	19	4%
20-24	42	13%	51	15%	62	17%	66	16%	75	17%
25-29	53	17%	53	15%	54	14%	57	14%	63	14%
30-34	70	22%	71	21%	74	20%	78	19%	80	18%
35-39	52	16%	55	16%	59	16%	64	16%	66	15%
40-44	42	13%	45	13%	48	13%	54	13%	55	13%
45-49	22	7%	25	7%	28	7%	29	7%	31	7%
50-54	14	4%	15	4%	16	4%	20	5%	22	5%
55-64	10	3%	11	3%	13	3%	15	4%	18	4%
65+	1	<1%	1	<1%	2	1%	2	<1%	3	1%
Race/Ethnicity										
White, not Hispanic	228	71%	241	70%	261	70%	281	69%	301	69%
Black/African American, not Hispanic	76	24%	85	25%	92	25%	102	25%	111	25%
Hispanic/Latino	10	3%	11	3%	11	3%	12	3%	12	3%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
Unknown	4	1%	5	1%	8	2%	9	2%	10	2%
Race/Sex										
White, not Hispanic Males	186	58%	198	58%	216	58%	233	57%	250	57%
White, not Hispanic Females	42	13%	43	13%	45	12%	48	12%	51	12%
Black/African American, not Hispanic Males	53	17%	60	17%	65	17%	74	18%	81	19%
Black/African American, not Hispanic Females	23	7%	25	7%	27	7%	28	7%	30	7%
Hispanic/Latino Males	8	3%	9	3%	9	2%	10	2%	10	2%
Hispanic/Latina Females	2	1%	2	1%	2	1%	2	<1%	2	<1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	4	1%	5	1%	8	2%	9	2%	10	2%
Total	320		344		374		407		437	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Canton Consortium 8

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	176	70%	194	71%	211	71%	232	71%	252	71%
Injection drug use (IDU)	5	2%	6	2%	7	2%	7	2%	7	2%
Male-to-male sexual contact and IDU	8	3%	8	3%	9	3%	11	3%	12	3%
Heterosexual contact	13	5%	13	5%	16	5%	19	6%	20	6%
Other/unknown	51	20%	53	19%	56	19%	59	18%	62	18%
Subtotal	253	100%	274	100%	299	100%	328	100%	353	100%
Female adult or adolescent										
Injection drug use	6	10%	6	9%	6	9%	6	8%	8	10%
Heterosexual contact	36	58%	39	60%	43	61%	45	61%	48	61%
Other/unknown	20	32%	20	31%	21	30%	23	31%	23	29%
Subtotal	62	100%	65	100%	70	100%	74	100%	79	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	4	80%	4	80%	4	80%
Other/unknown	1	20%	1	20%	1	20%	1	20%	1	20%
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	320		344		374		407		437	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Canton Consortium 8

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	171	53%	187	54%	203	54%	222	55%	240	55%
Injection drug use (IDU) only	4	1%	4	1%	5	1%	5	1%	6	1%
Heterosexual contact only	49	15%	52	15%	59	16%	64	16%	68	16%
MSM & IDU	5	2%	5	1%	6	2%	7	2%	7	2%
IDU & Heterosexual contact	7	2%	8	2%	8	2%	8	2%	9	2%
MSM & Heterosexual contact	5	2%	7	2%	8	2%	10	2%	12	3%
MSM & IDU & Heterosexual contact	3	1%	3	1%	3	1%	4	1%	5	1%
Perinatal exposure	4	1%	4	1%	4	1%	4	1%	4	1%
Other/unknown	72	23%	74	22%	78	21%	83	20%	86	20%
Total	320		344		374		407		437	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B Lima
Consortium 9A

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Lima Consortium 9A

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	5.1	13	87%	8	80%	3	100%	2	100%
Females	*	2	13%	2	20%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	1	7%	-	-	1	33%	-	-
20-24	*	1	7%	1	10%	-	-	-	-
25-29	*	1	7%	1	10%	-	-	-	-
30-34	*	2	13%	2	20%	-	-	-	-
35-39	*	1	7%	1	10%	-	-	-	-
40-44	*	4	27%	1	10%	1	33%	2	100%
45-49	*	1	7%	-	-	1	33%	-	-
50-54	*	3	20%	3	30%	-	-	-	-
55-64	*	1	7%	1	10%	-	-	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	1.5	7	47%	4	40%	1	33%	2	100%
Black/African American, not Hispanic	28.7	5	33%	4	40%	1	33%	-	-
Hispanic/Latino	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	3	20%	2	20%	1	33%	-	-
Race/Sex									
White, not Hispanic Males	3.0	7	47%	4	40%	1	33%	2	100%
White, not Hispanic Females	*	-	-	-	-	-	-	-	-
Black/African American, not Hispanic Males	*	3	20%	2	20%	1	33%	-	-
Black/African American, not Hispanic Females	*	2	13%	2	20%	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	3	20%	2	20%	1	33%	-	-
Total	2.9	15		10		3		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Lima Consortium 9A

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	7	54%	4	50%	3	100%	-	-
Injection drug use (IDU)	2	15%	2	25%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	8%	1	13%	-	-	-	-
Other/unknown	3	23%	1	13%	-	-	2	100%
Subtotal	13	100%	8	100%	3	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	2	100%	2	100%	-	-	-	-
Subtotal	2	100%	2	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	15		10		3		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Lima Consortium 9A

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	4	27%	2	20%	2	67%	-	-
Injection drug use (IDU) only	2	13%	2	20%	-	-	-	-
Heterosexual contact only	1	7%	1	10%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	3	20%	2	20%	1	33%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	5	33%	3	30%	-	-	2	100%
Total	15		10		3		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Lima Consortium 9A

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	14	82%	2	100%	-	-	-	-	-	-	1	33%
Injection drug use (IDU)	1	6%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	12%	-	-	-	-	-	-	-	-	2	67%
Subtotal	17	100%	2	100%	-	-	-	-	-	-	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	33%	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	67%	3	100%	-	-	-	-	-	-	-	-
Subtotal	6	100%	3	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	23		5		0		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Lima Consortium 9A

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	61%	2	40%	-	-	-	-	-	-	1	33%
Injection drug use (IDU) only	1	4%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	9%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	26%	3	60%	-	-	-	-	-	-	2	67%
Total	23		5		0		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Lima Consortium 9A**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	4.7	12	75%	3.5	9	90%	7.4	19	86%	5.9	15	88%
Females	*	4	25%	*	1	10%	*	3	14%	*	2	12%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	1	6%	*	1	10%	*	4	18%	15.0	5	29%
25-29	*	3	19%	17.4	5	50%	*	4	18%	*	1	6%
30-34	*	3	19%	*	-	-	*	2	9%	*	3	18%
35-39	*	1	6%	*	1	10%	*	1	5%	*	3	18%
40-44	*	1	6%	*	1	10%	*	2	9%	15.6	5	29%
45-49	14.2	5	31%	*	1	10%	*	4	18%	*	-	-
50-54	*	1	6%	*	-	-	*	3	14%	*	-	-
55-64	*	1	6%	*	1	10%	*	2	9%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity												
White, not Hispanic	2.3	11	69%	1.5	7	70%	4.0	19	86%	2.5	12	71%
Black/African American, not Hispanic	*	3	19%	*	3	30%	*	2	9%	*	3	18%
Hispanic/Latino	*	2	13%	*	-	-	*	-	-	*	1	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	1	6%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	1	5%	*	-	-
Race/Sex												
White, not Hispanic Males	3.9	9	56%	2.6	6	60%	6.8	16	73%	4.7	11	65%
White, not Hispanic Females	*	2	13%	*	1	10%	*	3	14%	*	1	6%
Black/African American, not Hispanic Males	*	2	13%	*	3	30%	*	2	9%	*	2	12%
Black/African American, not Hispanic Females	*	1	6%	*	-	-	*	-	-	*	1	6%
Hispanic/Latino Males	*	1	6%	*	-	-	*	-	-	*	1	6%
Hispanic/Latina Females	*	1	6%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	6%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	1	5%	*	-	-
Total	3.1	16		1.9	10		4.3	22		3.3	17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Lima Consortium 9A

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	5	42%	8	89%	7	37%	8	53%
Injection drug use (IDU)	2	17%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	5%	-	-
Heterosexual contact	3	25%	-	-	1	5%	-	-
Other/unknown	2	17%	1	11%	10	53%	7	47%
Subtotal	12	100%	9	100%	19	100%	15	100%
Female adult or adolescent								
Injection drug use	2	50%	1	100%	-	-	-	-
Heterosexual contact	1	25%	-	-	3	100%	1	50%
Other/unknown	1	25%	-	-	-	-	1	50%
Subtotal	4	100%	1	100%	3	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	16		10		22		17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Lima Consortium 9A

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	4	25%	6	60%	7	32%	8	47%
Injection drug use (IDU) only	2	13%	-	-	-	-	-	-
Heterosexual contact only	4	25%	-	-	4	18%	1	6%
MSM & IDU	-	-	-	-	1	5%	-	-
IDU & Heterosexual contact	2	13%	1	10%	-	-	-	-
MSM & Heterosexual contact	1	6%	2	20%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	3	19%	1	10%	10	45%	8	47%
Total	16		10		22		17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Lima Consortium 9A

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	4	80%	5	100%	12	100%	12	92%	8	80%
Females	1	20%	-	-	-	-	1	8%	2	20%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	-	-	1	20%	4	33%	5	38%	1	10%
25-29	-	-	3	60%	1	8%	1	8%	1	10%
30-34	1	20%	-	-	-	-	2	15%	2	20%
35-39	1	20%	1	20%	1	8%	2	15%	1	10%
40-44	-	-	-	-	2	17%	3	23%	1	10%
45-49	3	60%	-	-	2	17%	-	-	-	-
50-54	-	-	-	-	1	8%	-	-	3	30%
55-64	-	-	-	-	1	8%	-	-	1	10%
65+	-	-	-	-	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	3	60%	4	80%	10	83%	9	69%	4	40%
Black/African American, not Hispanic	-	-	1	20%	2	17%	2	15%	4	40%
Hispanic/Latino	2	40%	-	-	-	-	1	8%	-	-
Asian/Pacific Islander	-	-	-	-	-	-	1	8%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	2	20%
Race/Sex										
White, not Hispanic Males	3	60%	4	80%	10	83%	8	62%	4	40%
White, not Hispanic Females	-	-	-	-	-	-	1	8%	-	-
Black/African American, not Hispanic Males	-	-	1	20%	2	17%	2	15%	2	20%
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	2	20%
Hispanic/Latino Males	1	20%	-	-	-	-	1	8%	-	-
Hispanic/Latina Females	1	20%	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	1	8%	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	2	20%
Total	5		5		12		13		10	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Lima Consortium 9A

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1	25%	4	80%	6	50%	6	50%	4	50%
Injection drug use (IDU)	1	25%	-	-	-	-	-	-	2	25%
Male-to-male sexual contact and IDU	-	-	-	-	1	8%	-	-	-	-
Heterosexual contact	2	50%	-	-	-	-	-	-	1	13%
Other/unknown	-	-	1	20%	5	42%	6	50%	1	13%
Subtotal	4	100%	5	100%	12	100%	12	100%	8	100%
Female adult or adolescent										
Injection drug use	1	100%	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	100%
Subtotal	1	100%	-	-	-	-	1	100%	2	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	5		5		12		13		10	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Lima Consortium 9A

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	1	20%	2	40%	6	50%	6	46%	2	20%
Injection drug use (IDU) only	1	20%	-	-	-	-	-	-	2	20%
Heterosexual contact only	2	40%	-	-	-	-	1	8%	1	10%
MSM & IDU	-	-	-	-	1	8%	-	-	-	-
IDU & Heterosexual contact	1	20%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	2	40%	-	-	-	-	2	20%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	20%	5	42%	6	46%	3	30%
Total	5		5		12		13		10	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Lima Consortium 9A

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	7	64%	5	42%	11	85%	6	75%	9	90%	272	82%
Females	4	36%	7	58%	2	15%	2	25%	1	10%	58	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	5	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	-	-	2	20%	7	2%
20-24	1	9%	2	17%	-	-	-	-	-	-	20	6%
25-29	2	18%	3	25%	4	31%	3	38%	1	10%	53	16%
30-34	3	27%	1	8%	2	15%	2	25%	-	-	57	17%
35-39	-	-	2	17%	2	15%	1	13%	1	10%	58	18%
40-44	1	9%	1	8%	1	8%	1	13%	4	40%	48	15%
45-49	2	18%	2	17%	2	15%	-	-	2	20%	35	11%
50-54	1	9%	-	-	2	15%	-	-	-	-	25	8%
55-64	1	9%	1	8%	-	-	1	13%	-	-	16	5%
65+	-	-	-	-	-	-	-	-	-	-	5	2%
Race/Ethnicity												
White, not Hispanic	7	64%	7	58%	11	85%	6	75%	7	70%	267	81%
Black/African American, not Hispanic	4	36%	4	33%	-	-	2	25%	2	20%	47	14%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	9	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	8%	2	15%	-	-	1	10%	6	2%
Race/Sex												
White, not Hispanic Males	5	45%	3	25%	9	69%	5	63%	7	70%	232	70%
White, not Hispanic Females	2	18%	4	33%	2	15%	1	13%	-	-	35	11%
Black/African American, not Hispanic Males	2	18%	2	17%	-	-	1	13%	1	10%	29	9%
Black/African American, not Hispanic Females	2	18%	2	17%	-	-	1	13%	1	10%	18	5%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	6	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	3	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	8%	2	15%	-	-	1	10%	6	2%
Total	11		12		13		8		10		330	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Lima Consortium 9A

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	3	43%	4	80%	6	55%	4	67%	5	56%	182	67%
Injection drug use (IDU)	1	14%	1	20%	-	-	-	-	-	-	10	4%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	1	11%	11	4%
Heterosexual contact	1	14%	-	-	1	9%	-	-	-	-	18	7%
Other/unknown	2	29%	-	-	4	36%	2	33%	3	33%	51	19%
Subtotal	7	100%	5	100%	11	100%	6	100%	9	100%	272	100%
Female adult or adolescent												
Injection drug use	1	25%	1	14%	-	-	-	-	-	-	8	15%
Heterosexual contact	2	50%	4	57%	2	100%	1	50%	1	100%	38	72%
Other/unknown	1	25%	2	29%	-	-	1	50%	-	-	7	13%
Subtotal	4	100%	7	100%	2	100%	2	100%	1	100%	53	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	20%
Subtotal	-	-	-	-	-	-	-	-	-	-	5	100%
Total	11		12		13		8		10		330	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Lima Consortium 9A

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	2	18%	4	33%	6	46%	4	50%	4	40%	169	51%
Injection drug use (IDU) only	1	9%	1	8%	-	-	-	-	-	-	9	3%
Heterosexual contact only	3	27%	4	33%	3	23%	1	13%	1	10%	56	17%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	8	2%
IDU & Heterosexual contact	1	9%	1	8%	-	-	-	-	-	-	9	3%
MSM & Heterosexual contact	1	9%	-	-	-	-	-	-	1	10%	13	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	10%	3	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	4	1%
Other/unknown	3	27%	2	17%	4	31%	3	38%	3	30%	59	18%
Total	11		12		13		8		10		330	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Lima Consortium 9A

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Lima Consortium 9A

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	84.7	217	76%	106	76%	111	76%
Females	26.6	69	24%	33	24%	36	24%
Age at end of year							
<13	*	4	1%	-	-	4	3%
13-14	*	1	<1%	-	-	1	1%
15-19	24.7	9	3%	5	4%	4	3%
20-24	140.7	47	16%	28	20%	19	13%
25-29	173.9	50	17%	20	14%	30	20%
30-34	169.6	51	18%	25	18%	26	18%
35-39	147.9	43	15%	23	17%	20	14%
40-44	118.2	38	13%	22	16%	16	11%
45-49	68.1	24	8%	9	6%	15	10%
50-54	33.1	13	5%	3	2%	10	7%
55-64	7.3	5	2%	3	2%	2	1%
65+	*	1	<1%	1	1%	-	-
Race/Ethnicity							
White, not Hispanic	43.6	206	72%	101	73%	105	71%
Black/African American, not Hispanic	338.9	59	21%	28	20%	31	21%
Hispanic/Latino	97.3	13	5%	8	6%	5	3%
Asian/Pacific Islander	*	2	1%	1	1%	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	6	2%	1	1%	5	3%
Race/Sex							
White, not Hispanic Males	70.2	164	57%	81	58%	83	56%
White, not Hispanic Females	17.6	42	15%	20	14%	22	15%
Black/African American, not Hispanic Males	396.4	37	13%	17	12%	20	14%
Black/African American, not Hispanic Females	272.5	22	8%	11	8%	11	7%
Hispanic/Latino Males	127.7	9	3%	6	4%	3	2%
Hispanic/Latina Females	*	4	1%	2	1%	2	1%
Asian/Pacific Islander Males	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	6	2%	1	1%	5	3%
Total	55.5	286		139		147	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Lima Consortium 9A

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	137	63%	64	60%	73	66%
Injection drug use (IDU)	7	3%	4	4%	3	3%
Male-to-male sexual contact and IDU	11	5%	7	7%	4	4%
Heterosexual contact	16	7%	5	5%	11	10%
Other/unknown	45	21%	26	25%	19	17%
Subtotal	216	100%	106	100%	110	100%
Female adult or adolescent						
Injection drug use	8	12%	3	9%	5	15%
Heterosexual contact	42	64%	19	58%	23	70%
Other/unknown	16	24%	11	33%	5	15%
Subtotal	66	100%	33	100%	33	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	100%	-	-	4	100%
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	-	-	4	100%
Total	286		139		147	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Lima Consortium 9A

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	122	43%	58	42%	64	44%
Injection drug use (IDU) only	7	2%	4	3%	3	2%
Heterosexual contact only	58	20%	24	17%	34	23%
MSM & IDU	9	3%	6	4%	3	2%
IDU & Heterosexual contact	8	3%	3	2%	5	3%
MSM & Heterosexual contact	15	5%	6	4%	9	6%
MSM & IDU & Heterosexual contact	2	1%	1	1%	1	1%
Perinatal exposure	4	1%	-	-	4	3%
Other/unknown	61	21%	37	27%	24	16%
Total	286		139		147	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Lima Consortium 9A

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	111	68%	20	56%	1	11%	1	50%	-	-	4	80%
Injection drug use (IDU)	2	1%	4	11%	1	11%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	9	5%	1	3%	1	11%	-	-	-	-	-	-
Heterosexual contact	8	5%	4	11%	4	44%	-	-	-	-	-	-
Other/unknown	34	21%	7	19%	2	22%	1	50%	-	-	1	20%
Subtotal	164	100%	36	100%	9	100%	2	100%	-	-	5	100%
Female adult or adolescent												
Injection drug use	4	10%	4	18%	-	-	-	-	-	-	-	-
Heterosexual contact	27	68%	12	55%	3	100%	-	-	-	-	-	-
Other/unknown	9	23%	6	27%	-	-	-	-	-	-	1	100%
Subtotal	40	100%	22	100%	3	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	1	100%	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	1	100%	1	100%	-	-	-	-	-	-
Total	206		59		13		2		0		6	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Lima Consortium 9A

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	98	48%	18	31%	1	8%	1	50%	-	-	4	67%
Injection drug use (IDU) only	2	1%	4	7%	1	8%	-	-	-	-	-	-
Heterosexual contact only	35	17%	16	27%	7	54%	-	-	-	-	-	-
MSM & IDU	7	3%	1	2%	1	8%	-	-	-	-	-	-
IDU & Heterosexual contact	4	2%	4	7%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	13	6%	2	3%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	2	1%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	2	1%	1	2%	1	8%	-	-	-	-	-	-
Other/unknown	43	21%	13	22%	2	15%	1	50%	-	-	2	33%
Total	206		59		13		2		0		6	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Lima Consortium 9A

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	166	73%	176	74%	183	74%	202	75%	217	76%
Females	61	27%	63	26%	64	26%	67	25%	69	24%
Age at end of year										
<13	4	2%	4	2%	4	2%	4	1%	4	1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	9	4%	9	4%	9	4%	9	3%	9	3%
20-24	36	16%	37	15%	38	15%	42	16%	47	16%
25-29	37	16%	40	17%	45	18%	49	18%	50	17%
30-34	44	19%	46	19%	46	19%	48	18%	51	18%
35-39	37	16%	38	16%	39	16%	40	15%	43	15%
40-44	30	13%	31	13%	31	13%	33	12%	38	13%
45-49	17	7%	19	8%	20	8%	24	9%	24	8%
50-54	9	4%	10	4%	10	4%	13	5%	13	5%
55-64	2	1%	3	1%	3	1%	5	2%	5	2%
65+	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Race/Ethnicity										
White, not Hispanic	159	70%	169	71%	175	71%	194	72%	206	72%
Black/African American, not Hispanic	51	22%	52	22%	54	22%	56	21%	59	21%
Hispanic/Latino	11	5%	12	5%	12	5%	12	4%	13	5%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	5	2%	5	2%	5	2%	6	2%	6	2%
Race/Sex										
White, not Hispanic Males	124	55%	132	55%	137	55%	153	57%	164	57%
White, not Hispanic Females	35	15%	37	15%	38	15%	41	15%	42	15%
Black/African American, not Hispanic Males	30	13%	31	13%	33	13%	35	13%	37	13%
Black/African American, not Hispanic Females	21	9%	21	9%	21	9%	21	8%	22	8%
Hispanic/Latino Males	7	3%	8	3%	8	3%	8	3%	9	3%
Hispanic/Latina Females	4	2%	4	2%	4	2%	4	1%	4	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	5	2%	5	2%	5	2%	6	2%	6	2%
Total	227		239		247		269		286	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Lima Consortium 9A

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	111	67%	116	66%	122	67%	129	64%	137	63%
Injection drug use (IDU)	7	4%	7	4%	7	4%	7	3%	7	3%
Male-to-male sexual contact and IDU	10	6%	10	6%	10	5%	11	5%	11	5%
Heterosexual contact	12	7%	15	9%	15	8%	16	8%	16	7%
Other/unknown	25	15%	27	15%	28	15%	38	19%	45	21%
Subtotal	165	100%	175	100%	182	100%	201	100%	216	100%
Female adult or adolescent										
Injection drug use	6	10%	7	12%	8	13%	8	13%	8	12%
Heterosexual contact	37	64%	38	63%	38	62%	41	64%	42	64%
Other/unknown	15	26%	15	25%	15	25%	15	23%	16	24%
Subtotal	58	100%	60	100%	61	100%	64	100%	66	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	100%	4	100%	4	100%	4	100%	4	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	4	100%	4	100%	4	100%	4	100%
Total	227		239		247		269		286	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Lima Consortium 9A

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	99	44%	103	43%	107	43%	114	42%	122	43%
Injection drug use (IDU) only	7	3%	7	3%	7	3%	7	3%	7	2%
Heterosexual contact only	49	22%	53	22%	53	21%	57	21%	58	20%
MSM & IDU	8	4%	8	3%	8	3%	9	3%	9	3%
IDU & Heterosexual contact	6	3%	7	3%	8	3%	8	3%	8	3%
MSM & Heterosexual contact	12	5%	13	5%	15	6%	15	6%	15	5%
MSM & IDU & Heterosexual contact	2	1%	2	1%	2	1%	2	1%	2	1%
Perinatal exposure	4	2%	4	2%	4	2%	4	1%	4	1%
Other/unknown	40	18%	42	18%	43	17%	53	20%	61	21%
Total	227		239		247		269		286	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Mansfield Consortium 9B

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Mansfield Consortium 9B

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	5.4	16	89%	10	91%	4	80%	2	67%
Females	*	2	11%	1	9%	1	20%	1	33%
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	*	2	11%	1	9%	1	20%	-	-
25-29	*	3	17%	3	27%	-	-	-	-
30-34	*	4	22%	3	27%	1	20%	-	-
35-39	*	1	6%	1	9%	-	-	-	-
40-44	*	3	17%	1	9%	1	20%	1	33%
45-49	*	1	6%	1	9%	-	-	-	-
50-54	*	1	6%	-	-	1	20%	-	-
55-64	*	3	17%	1	9%	1	20%	2	67%
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	2.8	15	83%	10	91%	4	80%	1	33%
Black/African American, not Hispanic	*	1	6%	-	-	-	-	2	67%
Hispanic/Latino	*	1	6%	-	-	1	20%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	6%	1	9%	-	-	-	-
Race/Sex									
White, not Hispanic Males	4.9	13	72%	9	82%	3	60%	1	33%
White, not Hispanic Females	*	2	11%	1	9%	1	20%	-	-
Black/African American, not Hispanic Males	*	1	6%	-	-	-	-	1	33%
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	1	33%
Hispanic/Latino Males	*	1	6%	-	-	1	20%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	6%	1	9%	-	-	-	-
Total	3.0	18		11		5		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Mansfield Consortium 9B

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	11	69%	8	80%	3	75%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	5	31%	2	20%	1	25%	2	100%
Subtotal	16	100%	10	100%	4	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	1	50%	1	100%	-	-	-	-
Other/unknown	1	50%	-	-	1	100%	1	100%
Subtotal	2	100%	1	100%	1	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	18		11		5		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Mansfield Consortium 9B

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	11	61%	8	73%	3	60%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	1	6%	1	9%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	6	33%	2	18%	2	40%	3	100%
Total	18		11		5		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B
Mansfield Consortium 9B**

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	9	69%	-	-	1	100%	-	-	-	-	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	31%	1	100%	-	-	-	-	-	-	-	-
Subtotal	13	100%	1	100%	1	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	50%	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	15		1		1		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Mansfield Consortium 9B

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	9	60%	-	-	1	100%	-	-	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	7%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	33%	1	100%	-	-	-	-	-	-	-	-
Total	15		1		1		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Mansfield Consortium 9B**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	3.0	9	75%	4.0	12	86%	2.7	8	80%	4.4	13	76%
Females	*	3	25%	*	2	14%	*	2	20%	*	4	24%
Age at diagnosis (yr)												
<13	*	-	-	*	1	7%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	8%	*	2	14%	*	2	20%	*	-	-
20-24	*	1	8%	*	2	14%	*	3	30%	13.2	5	29%
25-29	*	-	-	*	1	7%	*	2	20%	*	3	18%
30-34	*	2	17%	*	2	14%	*	2	20%	*	1	6%
35-39	*	2	17%	*	1	7%	*	1	10%	*	1	6%
40-44	*	2	17%	*	3	21%	*	-	-	*	-	-
45-49	*	2	17%	*	1	7%	*	-	-	*	3	18%
50-54	*	1	8%	*	1	7%	*	-	-	*	-	-
55-64	*	1	8%	*	-	-	*	-	-	*	3	18%
65+	*	-	-	*	-	-	*	-	-	*	1	6%
Race/Ethnicity												
White, not Hispanic	1.5	8	67%	2.0	11	79%	1.3	7	70%	2.0	11	65%
Black/African American, not Hispanic	*	3	25%	*	2	14%	*	3	30%	*	-	-
Hispanic/Latino	*	-	-	*	1	7%	*	-	-	*	4	24%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	1	6%
American Indian/Alaska Native	*	1	8%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	1	6%
Race/Sex												
White, not Hispanic Males	2.6	7	58%	3.4	9	64%	2.3	6	60%	3.0	8	47%
White, not Hispanic Females	*	1	8%	*	2	14%	*	1	10%	*	3	18%
Black/African American, not Hispanic Males	*	2	17%	*	2	14%	*	2	20%	*	-	-
Black/African American, not Hispanic Females	*	1	8%	*	-	-	*	1	10%	*	-	-
Hispanic/Latino Males	*	-	-	*	1	7%	*	-	-	*	4	24%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	1	6%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	1	8%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	1	6%
Total	2.0	12		2.4	14		1.7	10		2.9	17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Mansfield Consortium 9B

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	5	56%	8	67%	7	88%	10	77%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	11%	1	8%	-	-	3	23%
Other/unknown	3	33%	3	25%	1	13%	-	-
Subtotal	9	100%	12	100%	8	100%	13	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	1	25%
Heterosexual contact	3	100%	1	100%	1	50%	3	75%
Other/unknown	-	-	-	-	1	50%	-	-
Subtotal	3	100%	1	100%	2	100%	4	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-
Total	12		14		10		17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Mansfield Consortium 9B

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	4	33%	8	57%	7	70%	10	59%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	33%	2	14%	1	10%	6	35%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	1	6%
MSM & Heterosexual contact	1	8%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	3	25%	4	29%	2	20%	-	-
Total	12		14		10		17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Mansfield Consortium 9B

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	3	75%	7	78%	6	75%	12	86%	10	91%
Females	1	25%	2	22%	2	25%	2	14%	1	9%
Age at HIV diagnosis (yr)										
<13	-	-	1	11%	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	25%	1	11%	1	13%	-	-	-	-
20-24	1	25%	1	11%	3	38%	4	29%	1	9%
25-29	-	-	1	11%	2	25%	2	14%	3	27%
30-34	-	-	2	22%	2	25%	1	7%	3	27%
35-39	1	25%	-	-	-	-	-	-	1	9%
40-44	-	-	1	11%	-	-	-	-	1	9%
45-49	-	-	1	11%	-	-	3	21%	1	9%
50-54	1	25%	1	11%	-	-	-	-	-	-
55-64	-	-	-	-	-	-	3	21%	1	9%
65+	-	-	-	-	-	-	1	7%	-	-
Race/Ethnicity										
White, not Hispanic	3	75%	8	89%	5	63%	8	57%	10	91%
Black/African American, not Hispanic	1	25%	1	11%	3	38%	-	-	-	-
Hispanic/Latino	-	-	-	-	-	-	4	29%	-	-
Asian/Pacific Islander	-	-	-	-	-	-	1	7%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	1	7%	1	9%
Race/Sex										
White, not Hispanic Males	2	50%	6	67%	4	50%	7	50%	9	82%
White, not Hispanic Females	1	25%	2	22%	1	13%	1	7%	1	9%
Black/African American, not Hispanic Males	1	25%	1	11%	2	25%	-	-	-	-
Black/African American, not Hispanic Females	-	-	-	-	1	13%	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-	4	29%	-	-
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	1	7%	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	1	7%	1	9%
Total	4		9		8		14		11	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Mansfield Consortium 9B

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	2	67%	4	57%	5	83%	9	75%	8	80%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	14%	-	-	3	25%	-	-
Other/unknown	1	33%	2	29%	1	17%	-	-	2	20%
Subtotal	3	100%	7	100%	6	100%	12	100%	10	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	100%	1	100%	1	50%	2	100%	1	100%
Other/unknown	-	-	-	-	1	50%	-	-	-	-
Subtotal	1	100%	1	100%	2	100%	2	100%	1	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-
Total	4		9		8		14		11	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Mansfield Consortium 9B

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	2	50%	4	44%	5	63%	9	64%	8	73%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	25%	2	22%	1	13%	5	36%	1	9%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	25%	3	33%	2	25%	-	-	2	18%
Total	4		9		8		14		11	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Mansfield Consortium 9B

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	11	79%	7	88%	3	60%	6	75%	7	78%	361	85%
Females	3	21%	1	13%	2	40%	2	25%	2	22%	62	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	7	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	2	25%	1	11%	10	2%
20-24	2	14%	1	13%	1	20%	1	13%	1	11%	33	8%
25-29	1	7%	1	13%	1	20%	-	-	1	11%	73	17%
30-34	1	7%	-	-	2	40%	1	13%	1	11%	71	17%
35-39	3	21%	2	25%	1	20%	1	13%	-	-	91	22%
40-44	4	29%	3	38%	-	-	2	25%	2	22%	62	15%
45-49	2	14%	1	13%	-	-	-	-	-	-	37	9%
50-54	-	-	-	-	-	-	1	13%	1	11%	14	3%
55-64	1	7%	-	-	-	-	-	-	2	22%	19	4%
65+	-	-	-	-	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	9	64%	5	63%	4	80%	7	88%	7	78%	351	83%
Black/African American, not Hispanic	4	29%	2	25%	1	20%	1	13%	1	11%	56	13%
Hispanic/Latino	-	-	1	13%	-	-	-	-	1	11%	12	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	7%	-	-	-	-	-	-	-	-	2	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	2	<1%
Race/Sex												
White, not Hispanic Males	8	57%	4	50%	3	60%	6	75%	5	56%	306	72%
White, not Hispanic Females	1	7%	1	13%	1	20%	1	13%	2	22%	45	11%
Black/African American, not Hispanic Males	3	21%	2	25%	-	-	-	-	1	11%	43	10%
Black/African American, not Hispanic Females	1	7%	-	-	1	20%	1	13%	-	-	13	3%
Hispanic/Latino Males	-	-	1	13%	-	-	-	-	1	11%	10	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	1	7%	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	2	<1%
Total	14		8		5		8		9		423	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Mansfield Consortium 9B

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	82%	4	57%	3	100%	6	100%	4	57%	246	69%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	20	6%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	30	8%
Heterosexual contact	-	-	1	14%	-	-	-	-	-	-	15	4%
Other/unknown	2	18%	2	29%	-	-	-	-	3	43%	46	13%
Subtotal	11	100%	7	100%	3	100%	6	100%	7	100%	357	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	1	50%	9	15%
Heterosexual contact	3	100%	1	100%	-	-	2	100%	-	-	39	66%
Other/unknown	-	-	-	-	2	100%	-	-	1	50%	11	19%
Subtotal	3	100%	1	100%	2	100%	2	100%	2	100%	59	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	71%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	29%
Subtotal	-	-	-	-	-	-	-	-	-	-	7	100%
Total	14		8		5		8		9		423	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Mansfield Consortium 9B

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	8	57%	3	38%	2	40%	6	75%	4	44%	237	56%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	15	4%
Heterosexual contact only	3	21%	2	25%	-	-	2	25%	-	-	54	13%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	25	6%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	11%	14	3%
MSM & Heterosexual contact	1	7%	1	13%	1	20%	-	-	-	-	9	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	5	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	2	14%	2	25%	2	40%	-	-	4	44%	59	14%
Total	14		8		5		8		9		423	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Mansfield Consortium 9B

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Mansfield Consortium 9B

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	78.2	232	77%	94	72%	138	80%
Females	23.8	71	23%	36	28%	35	20%
Age at end of year							
<13	6.3	6	2%	4	3%	2	1%
13-14	*	2	1%	2	2%	-	-
15-19	34.6	14	5%	8	6%	6	3%
20-24	118.9	45	15%	26	20%	19	11%
25-29	142.5	47	16%	20	15%	27	16%
30-34	121.3	42	14%	20	15%	22	13%
35-39	188.8	64	21%	23	18%	41	24%
40-44	95.4	36	12%	7	5%	29	17%
45-49	58.9	24	8%	9	7%	15	9%
50-54	19.8	9	3%	4	3%	5	3%
55-64	13.3	11	4%	6	5%	5	3%
65+	*	2	1%	-	-	2	1%
Race/Ethnicity							
White, not Hispanic	40.7	220	73%	95	73%	125	72%
Black/African American, not Hispanic	228.9	58	19%	23	18%	35	20%
Hispanic/Latino	104.9	16	5%	8	6%	8	5%
Asian/Pacific Islander	*	1	<1%	1	1%	-	-
American Indian/Alaska Native	*	1	<1%	-	-	1	1%
Unknown	*	7	2%	3	2%	4	2%
Race/Sex							
White, not Hispanic Males	65.7	175	58%	72	55%	103	60%
White, not Hispanic Females	16.4	45	15%	23	18%	22	13%
Black/African American, not Hispanic Males	250.8	39	13%	12	9%	27	16%
Black/African American, not Hispanic Females	194.2	19	6%	11	8%	8	5%
Hispanic/Latino Males	172.8	14	5%	8	6%	6	3%
Hispanic/Latina Females	*	2	1%	-	-	2	1%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	1%
Unknown	*	7	2%	3	2%	4	2%
Total	51.0	303		130		173	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Mansfield Consortium 9B

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	155	68%	54	59%	101	74%
Injection drug use (IDU)	6	3%	1	1%	5	4%
Male-to-male sexual contact and IDU	14	6%	3	3%	11	8%
Heterosexual contact	16	7%	7	8%	9	7%
Other/unknown	37	16%	26	29%	11	8%
Subtotal	228	100%	91	100%	137	100%
Female adult or adolescent						
Injection drug use	4	6%	-	-	4	12%
Heterosexual contact	49	71%	26	74%	23	68%
Other/unknown	16	23%	9	26%	7	21%
Subtotal	69	100%	35	100%	34	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	67%	2	50%	2	100%
Other/unknown	2	33%	2	50%	-	-
Subtotal	6	100%	4	100%	2	100%
Total	303		130		173	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Mansfield Consortium 9B

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	148	49%	53	41%	95	55%
Injection drug use (IDU) only	3	1%	1	1%	2	1%
Heterosexual contact only	65	21%	33	25%	32	18%
MSM & IDU	12	4%	3	2%	9	5%
IDU & Heterosexual contact	7	2%	-	-	7	4%
MSM & Heterosexual contact	7	2%	1	1%	6	3%
MSM & IDU & Heterosexual contact	2	1%	-	-	2	1%
Perinatal exposure	4	1%	2	2%	2	1%
Other/unknown	55	18%	37	28%	18	10%
Total	303		130		173	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Mansfield Consortium 9B

Living with a Diagnosis of HIV Infection in 2011

Transmission Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	123	72%	22	56%	8	57%	-	-	-	-	2	50%
Injection drug use (IDU)	5	3%	-	-	1	7%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	10	6%	4	10%	-	-	-	-	-	-	-	-
Heterosexual contact	7	4%	5	13%	3	21%	-	-	-	-	1	25%
Other/unknown	26	15%	8	21%	2	14%	-	-	-	-	1	25%
Subtotal	171	100%	39	100%	14	100%	-	-	-	-	4	100%
Female adult or adolescent												
Injection drug use	3	7%	-	-	-	-	-	-	-	-	1	33%
Heterosexual contact	28	64%	16	89%	1	50%	1	100%	1	100%	2	67%
Other/unknown	13	30%	2	11%	1	50%	-	-	-	-	-	-
Subtotal	44	100%	18	100%	2	100%	1	100%	1	100%	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	60%	1	100%	-	-	-	-	-	-	-	-
Other/unknown	2	40%	-	-	-	-	-	-	-	-	-	-
Subtotal	5	100%	1	100%	-	-	-	-	-	-	-	-
Total	220		58		16		1		1		7	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Mansfield Consortium 9B

Living with a Diagnosis of HIV Infection in 2011

Exposure Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	116	53%	22	38%	8	50%	-	-	-	-	2	29%
Injection drug use (IDU) only	3	1%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	35	16%	21	36%	4	25%	1	100%	1	100%	3	43%
MSM & IDU	9	4%	3	5%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	5	2%	-	-	1	6%	-	-	-	-	1	14%
MSM & Heterosexual contact	7	3%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	<1%	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	3	1%	1	2%	-	-	-	-	-	-	-	-
Other/unknown	41	19%	10	17%	3	19%	-	-	-	-	1	14%
Total	220		58		16		1		1		7	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Mansfield Consortium 9B

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	191	77%	200	76%	212	77%	220	77%	232	77%
Females	58	23%	63	24%	65	23%	67	23%	71	23%
Age at end of year										
<13	5	2%	5	2%	6	2%	6	2%	6	2%
13-14	2	1%	2	1%	2	1%	2	1%	2	1%
15-19	9	4%	10	4%	12	4%	14	5%	14	5%
20-24	33	13%	35	13%	37	13%	40	14%	45	15%
25-29	40	16%	41	16%	42	15%	44	15%	47	16%
30-34	34	14%	37	14%	39	14%	41	14%	42	14%
35-39	59	24%	61	23%	62	22%	63	22%	64	21%
40-44	32	13%	33	13%	36	13%	36	13%	36	12%
45-49	18	7%	20	8%	21	8%	21	7%	24	8%
50-54	7	3%	8	3%	9	3%	9	3%	9	3%
55-64	7	3%	8	3%	8	3%	8	3%	11	4%
65+	2	1%	2	1%	2	1%	2	1%	2	1%
Race/Ethnicity										
White, not Hispanic	184	74%	192	73%	203	73%	210	73%	220	73%
Black/African American, not Hispanic	49	20%	53	20%	55	20%	58	20%	58	19%
Hispanic/Latino	11	4%	11	4%	12	4%	12	4%	16	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	5	2%	6	2%	6	2%	6	2%	7	2%
Race/Sex										
White, not Hispanic Males	146	59%	153	58%	162	58%	168	59%	175	58%
White, not Hispanic Females	38	15%	39	15%	41	15%	42	15%	45	15%
Black/African American, not Hispanic Males	33	13%	35	13%	37	13%	39	14%	39	13%
Black/African American, not Hispanic Females	16	6%	18	7%	18	6%	19	7%	19	6%
Hispanic/Latino Males	9	4%	9	3%	10	4%	10	3%	14	5%
Hispanic/Latina Females	2	1%	2	1%	2	1%	2	1%	2	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	5	2%	6	2%	6	2%	6	2%	7	2%
Total	249		263		277		287		303	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Mansfield Consortium 9B

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	125	67%	131	67%	139	67%	146	68%	155	68%
Injection drug use (IDU)	6	3%	6	3%	6	3%	6	3%	6	3%
Male-to-male sexual contact and IDU	14	7%	14	7%	14	7%	14	6%	14	6%
Heterosexual contact	11	6%	12	6%	13	6%	13	6%	16	7%
Other/unknown	31	17%	33	17%	36	17%	37	17%	37	16%
Subtotal	187	100%	196	100%	208	100%	216	100%	228	100%
Female adult or adolescent										
Injection drug use	3	5%	3	5%	3	5%	3	5%	4	6%
Heterosexual contact	39	68%	44	71%	45	71%	46	71%	49	71%
Other/unknown	15	26%	15	24%	15	24%	16	25%	16	23%
Subtotal	57	100%	62	100%	63	100%	65	100%	69	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	4	67%	4	67%	4	67%
Other/unknown	1	20%	1	20%	2	33%	2	33%	2	33%
Subtotal	5	100%	5	100%	6	100%	6	100%	6	100%
Total	249		263		277		287		303	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Mansfield Consortium 9B

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	119	48%	124	47%	132	48%	139	48%	148	49%
Injection drug use (IDU) only	3	1%	3	1%	3	1%	3	1%	3	1%
Heterosexual contact only	50	20%	56	21%	58	21%	59	21%	65	21%
MSM & IDU	12	5%	12	5%	12	4%	12	4%	12	4%
IDU & Heterosexual contact	6	2%	6	2%	6	2%	6	2%	7	2%
MSM & Heterosexual contact	6	2%	7	3%	7	3%	7	2%	7	2%
MSM & IDU & Heterosexual contact	2	1%	2	1%	2	1%	2	1%	2	1%
Perinatal exposure	4	2%	4	2%	4	1%	4	1%	4	1%
Other/unknown	47	19%	49	19%	53	19%	55	19%	55	18%
Total	249		263		277		287		303	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV and/or AIDS
Reported in Ryan White Part B
Southeast Consortium 9C

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by selected characteristics, Part B Southeast Consortium 9C

Characteristic	2012 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	5.2	22	71%	12	63%	5	83%	5	83%	
Females	2.1	9	29%	7	37%	1	17%	1	17%	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	4	13%	3	16%	-	-	1	17%	
25-29	*	4	13%	2	11%	1	17%	1	17%	
30-34	*	4	13%	2	11%	1	17%	1	17%	
35-39	*	3	10%	2	11%	1	17%	-	-	
40-44	*	2	6%	1	5%	1	17%	-	-	
45-49	10.6	6	19%	4	21%	1	17%	1	17%	
50-54	7.9	5	16%	2	11%	1	17%	2	33%	
55-64	*	3	10%	3	16%	-	-	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity										
White, not Hispanic	2.9	23	74%	16	84%	2	33%	5	83%	
Black/African American, not Hispanic	23.8	5	16%	2	11%	2	33%	1	17%	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	3	10%	1	5%	2	33%	-	-	
Race/Sex										
White, not Hispanic Males	4.3	17	55%	11	58%	2	33%	4	67%	
White, not Hispanic Females	1.5	6	19%	5	26%	-	-	1	17%	
Black/African American, not Hispanic Males	*	2	6%	-	-	1	17%	1	17%	
Black/African American, not Hispanic Females	*	3	10%	2	11%	1	17%	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	3	10%	1	5%	2	33%	-	-	
Total	3.7	31		19		6		6		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	17	77%	9	75%	3	60%	5	100%
Injection drug use (IDU)	1	5%	-	-	1	20%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	4	18%	3	25%	1	20%	-	-
Subtotal	22	100%	12	100%	5	100%	5	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	2	22%	2	29%	-	-	-	-
Other/unknown	7	78%	5	71%	1	100%	1	100%
Subtotal	9	100%	7	100%	1	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	31		19		6		6	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by disease status at initial diagnosis and by exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	2012 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	17	55%	9	47%	3	50%	5	83%
Injection drug use (IDU) only	1	3%	-	-	1	17%	-	-
Heterosexual contact only	2	6%	2	11%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	11	35%	8	42%	2	33%	1	17%
Total	31		19		6		6	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	14	82%	2	100%	-	-	-	-	-	-	1	33%
Injection drug use (IDU)	1	6%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	12%	-	-	-	-	-	-	-	-	2	67%
Subtotal	17	100%	2	100%	-	-	-	-	-	-	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	33%	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	67%	3	100%	-	-	-	-	-	-	-	-
Subtotal	6	100%	3	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	23		5		0		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by race/ethnicity and exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	2012 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	61%	2	40%	-	-	-	-	-	-	1	33%
Injection drug use (IDU) only	1	4%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	9%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	26%	3	60%	-	-	-	-	-	-	2	67%
Total	23		5		0		0		0		3	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

**Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by selected characteristics,
Part B Southeast Consortium 9C**

Diagnosis of HIV Infection by Year

Characteristic	2008			2009			2010			2011		
	Rate ^a	No.	%									
Sex												
Males	4.3	18	82%	5.7	24	86%	4.0	17	81%	3.8	16	67%
Females	*	4	18%	*	4	14%	*	4	19%	1.9	8	33%
Age at diagnosis (yr)												
<13	*	-	-	*	1	4%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	5%	*	-	-	*	-	-	*	1	4%
20-24	*	2	9%	*	3	11%	*	2	10%	13.0	8	33%
25-29	*	2	9%	*	4	14%	*	3	14%	*	1	4%
30-34	16.0	8	36%	*	4	14%	10.0	5	24%	*	-	-
35-39	*	4	18%	12.4	6	21%	*	4	19%	10.3	5	21%
40-44	*	2	9%	*	3	11%	9.2	5	24%	*	-	-
45-49	*	-	-	*	2	7%	*	1	5%	*	4	17%
50-54	*	2	9%	*	2	7%	*	1	5%	*	2	8%
55-64	*	-	-	*	3	11%	*	-	-	*	2	8%
65+	*	1	5%	*	-	-	*	-	-	*	1	4%
Race/Ethnicity												
White, not Hispanic	2.3	18	82%	2.6	21	75%	2.1	17	81%	2.9	23	96%
Black/African American, not Hispanic	*	3	14%	*	3	11%	*	2	10%	*	1	4%
Hispanic/Latino	*	-	-	*	4	14%	*	1	5%	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	5%	*	-	-	*	1	5%	*	-	-
Race/Sex												
White, not Hispanic Males	3.8	15	68%	4.6	18	64%	3.8	15	71%	3.8	15	63%
White, not Hispanic Females	*	3	14%	*	3	11%	*	2	10%	2.0	8	33%
Black/African American, not Hispanic Males	*	2	9%	*	3	11%	*	-	-	*	1	4%
Black/African American, not Hispanic Females	*	1	5%	*	-	-	*	2	10%	*	-	-
Hispanic/Latino Males	*	-	-	*	3	11%	*	1	5%	*	-	-
Hispanic/Latina Females	*	-	-	*	1	4%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	5%	*	-	-	*	1	5%	*	-	-
Total	2.6	22		3.3	28		2.5	21		2.8	24	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by transmission category, Part B Southeast Consortium 9C

Diagnosis of HIV Infection by Year

Transmission Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	17	94%	17	74%	14	82%	11	69%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	4%	1	6%	-	-
Heterosexual contact	-	-	1	4%	-	-	1	6%
Other/unknown	1	6%	4	17%	2	12%	4	25%
Subtotal	18	100%	23	100%	17	100%	16	100%
Female adult or adolescent								
Injection drug use	1	25%	-	-	-	-	2	25%
Heterosexual contact	2	50%	3	75%	1	25%	4	50%
Other/unknown	1	25%	1	25%	3	75%	2	25%
Subtotal	4	100%	4	100%	4	100%	8	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-
Total	22		28		21		24	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2013, by year of diagnosis and by exposure category, Part B Southeast Consortium 9C

Diagnosis of HIV Infection by Year

Exposure Category ^a	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	17	77%	17	61%	14	67%	11	46%
Injection drug use (IDU) only	1	5%	-	-	-	-	1	4%
Heterosexual contact only	2	9%	4	14%	1	5%	5	21%
MSM & IDU	-	-	1	4%	1	5%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	1	4%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	4%	-	-	-	-
Other/unknown	2	9%	5	18%	5	24%	6	25%
Total	22		28		21		24	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2008-2011) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Part B Southeast Consortium 9C

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	13	81%	14	93%	11	92%	10	67%	12	63%
Females	3	19%	1	7%	1	8%	5	33%	7	37%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	6%	-	-	-	-	1	7%	-	-
20-24	1	6%	2	13%	1	8%	7	47%	3	16%
25-29	1	6%	3	20%	2	17%	1	7%	2	11%
30-34	6	38%	2	13%	4	33%	-	-	2	11%
35-39	2	13%	4	27%	3	25%	3	20%	2	11%
40-44	2	13%	1	7%	2	17%	-	-	1	5%
45-49	-	-	1	7%	-	-	1	7%	4	21%
50-54	2	13%	1	7%	-	-	1	7%	2	11%
55-64	-	-	1	7%	-	-	-	-	3	16%
65+	1	6%	-	-	-	-	1	7%	-	-
Race/Ethnicity										
White, not Hispanic	12	75%	11	73%	9	75%	14	93%	16	84%
Black/African American, not Hispanic	3	19%	2	13%	1	8%	1	7%	2	11%
Hispanic/Latino	-	-	2	13%	1	8%	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	6%	-	-	1	8%	-	-	1	5%
Race/Sex										
White, not Hispanic Males	10	63%	10	67%	9	75%	9	60%	11	58%
White, not Hispanic Females	2	13%	1	7%	-	-	5	33%	5	26%
Black/African American, not Hispanic Males	2	13%	2	13%	-	-	1	7%	-	-
Black/African American, not Hispanic Females	1	6%	-	-	1	8%	-	-	2	11%
Hispanic/Latino Males	-	-	2	13%	1	8%	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	6%	-	-	1	8%	-	-	1	5%
Total	16		15		12		15		19	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	12	92%	11	79%	9	82%	8	80%	9	75%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	9%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	1	10%	-	-
Other/unknown	1	8%	3	21%	1	9%	1	10%	3	25%
Subtotal	13	100%	14	100%	11	100%	10	100%	12	100%
Female adult or adolescent										
Injection drug use	1	33%	-	-	-	-	2	40%	-	-
Heterosexual contact	1	33%	1	100%	-	-	1	20%	2	29%
Other/unknown	1	33%	-	-	1	100%	2	40%	5	71%
Subtotal	3	100%	1	100%	1	100%	5	100%	7	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	16		15		12		15		19	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	75%	11	73%	9	75%	8	53%	9	47%
Injection drug use (IDU) only	1	6%	-	-	-	-	1	7%	-	-
Heterosexual contact only	1	6%	1	7%	-	-	2	13%	2	11%
MSM & IDU	-	-	-	-	1	8%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	1	7%	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	13%	3	20%	2	17%	3	20%	8	42%
Total	16		15		12		15		19	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Part B Southeast Consortium 9C

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	6	60%	10	67%	8	73%	7	78%	13	76%	456	86%
Females	4	40%	5	33%	3	27%	2	22%	4	24%	75	14%
Age at AIDS diagnosis (yr)												
<13	-	-	1	7%	-	-	-	-	-	-	10	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	-	-	1	6%	6	1%
20-24	1	10%	5	33%	2	18%	2	22%	1	6%	34	6%
25-29	1	10%	1	7%	1	9%	-	-	2	12%	76	14%
30-34	5	50%	1	7%	2	18%	-	-	2	12%	109	21%
35-39	3	30%	2	13%	2	18%	2	22%	1	6%	97	18%
40-44	-	-	2	13%	2	18%	1	11%	1	6%	97	18%
45-49	-	-	1	7%	1	9%	1	11%	5	29%	45	8%
50-54	-	-	1	7%	1	9%	1	11%	3	18%	28	5%
55-64	-	-	1	7%	-	-	2	22%	1	6%	25	5%
65+	-	-	-	-	-	-	-	-	-	-	3	1%
Race/Ethnicity												
White, not Hispanic	10	100%	11	73%	10	91%	9	100%	12	71%	481	91%
Black/African American, not Hispanic	-	-	1	7%	1	9%	-	-	3	18%	35	7%
Hispanic/Latino	-	-	2	13%	-	-	-	-	-	-	6	1%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	7%	-	-	-	-	2	12%	7	1%
Race/Sex												
White, not Hispanic Males	6	60%	7	47%	8	73%	7	78%	9	53%	417	79%
White, not Hispanic Females	4	40%	4	27%	2	18%	2	22%	3	18%	64	12%
Black/African American, not Hispanic Males	-	-	1	7%	-	-	-	-	2	12%	25	5%
Black/African American, not Hispanic Females	-	-	-	-	1	9%	-	-	1	6%	10	2%
Hispanic/Latino Males	-	-	1	7%	-	-	-	-	-	-	5	1%
Hispanic/Latina Females	-	-	1	7%	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	7%	-	-	-	-	2	12%	7	1%
Total	10		15		11		9		17		531	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	4	67%	7	78%	6	75%	4	57%	11	85%	306	68%
Injection drug use (IDU)	1	17%	-	-	-	-	-	-	1	8%	17	4%
Male-to-male sexual contact and IDU	-	-	1	11%	-	-	-	-	-	-	22	5%
Heterosexual contact	1	17%	1	11%	-	-	-	-	-	-	22	5%
Other/unknown	-	-	-	-	2	25%	3	43%	1	8%	80	18%
Subtotal	6	100%	9	100%	8	100%	7	100%	13	100%	447	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	11	15%
Heterosexual contact	4	100%	4	80%	1	33%	2	100%	2	50%	43	58%
Other/unknown	-	-	1	20%	2	67%	-	-	2	50%	20	27%
Subtotal	4	100%	5	100%	3	100%	2	100%	4	100%	74	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	4	40%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	60%
Subtotal	-	-	1	100%	-	-	-	-	-	-	10	100%
Total	10		15		11		9		17		531	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported cases of AIDS as of June 30, 2013, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2008		2009		2010		2011		2012		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	4	40%	7	47%	6	55%	4	44%	10	59%	292	55%
Injection drug use (IDU) only	1	10%	-	-	-	-	-	-	1	6%	19	4%
Heterosexual contact only	5	50%	5	33%	1	9%	2	22%	2	12%	65	12%
MSM & IDU	-	-	1	7%	-	-	-	-	-	-	17	3%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	9	2%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	1	6%	14	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	5	1%
Perinatal exposure	-	-	1	7%	-	-	-	-	-	-	4	1%
Other/unknown	-	-	1	7%	4	36%	3	33%	3	18%	106	20%
Total	10		15		11		9		17		531	

Note: AIDS diagnoses by year (2008-2012) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2012.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in Ryan White
Part B Southeast Consortium 9C

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by selected characteristics, Part B Southeast Consortium 9C

Characteristic	Living with a diagnosis of HIV infection in 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	79.5	334	76%	146	72%	188	80%
Females	24.8	105	24%	58	28%	47	20%
Age at end of year							
<13	6.9	9	2%	3	1%	6	3%
13-14	*	3	1%	2	1%	1	<1%
15-19	13.8	8	2%	4	2%	4	2%
20-24	86.3	53	12%	32	16%	21	9%
25-29	135.7	65	15%	34	17%	31	13%
30-34	170.5	85	19%	34	17%	51	22%
35-39	177.4	86	20%	41	20%	45	19%
40-44	97.6	53	12%	18	9%	35	15%
45-49	63.5	36	8%	14	7%	22	9%
50-54	42.4	27	6%	16	8%	11	5%
55-64	9.5	11	3%	3	1%	8	3%
65+	*	3	1%	3	1%	-	-
Race/Ethnicity							
White, not Hispanic	46.5	369	84%	166	81%	203	86%
Black/African American, not Hispanic	219.3	46	10%	23	11%	23	10%
Hispanic/Latino	147.7	12	3%	7	3%	5	2%
Asian/Pacific Islander	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	11	3%	8	4%	3	1%
Race/Sex							
White, not Hispanic Males	72.9	286	65%	121	59%	165	70%
White, not Hispanic Females	20.7	83	19%	45	22%	38	16%
Black/African American, not Hispanic Males	228.1	31	7%	15	7%	16	7%
Black/African American, not Hispanic Females	203.2	15	3%	8	4%	7	3%
Hispanic/Latino Males	189.7	8	2%	5	2%	3	1%
Hispanic/Latina Females	*	4	1%	2	1%	2	1%
Asian/Pacific Islander Males	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	11	3%	8	4%	3	1%
Total	52.1	439		204		235	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	218	66%	89	61%	129	70%
Injection drug use (IDU)	14	4%	4	3%	10	5%
Male-to-male sexual contact and IDU	12	4%	4	3%	8	4%
Heterosexual contact	29	9%	10	7%	19	10%
Other/unknown	55	17%	38	26%	17	9%
Subtotal	328	100%	145	100%	183	100%
Female adult or adolescent						
Injection drug use	9	9%	6	11%	3	7%
Heterosexual contact	60	59%	25	45%	35	76%
Other/unknown	32	32%	24	44%	8	17%
Subtotal	101	100%	55	100%	46	100%
Child (<13 yrs at diagnosis)						
Perinatal	6	60%	3	75%	3	50%
Other/unknown	4	40%	1	25%	3	50%
Subtotal	10	100%	4	100%	6	100%
Total	439		204		235	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by current disease status and by exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	Living with a diagnosis of HIV infection in 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	202	46%	84	41%	118	50%
Injection drug use (IDU) only	17	4%	6	3%	11	5%
Heterosexual contact only	89	20%	35	17%	54	23%
MSM & IDU	11	3%	4	2%	7	3%
IDU & Heterosexual contact	6	1%	4	2%	2	1%
MSM & Heterosexual contact	16	4%	5	2%	11	5%
MSM & IDU & Heterosexual contact	1	<1%	-	-	1	<1%
Perinatal exposure	7	2%	4	2%	3	1%
Other/unknown	90	21%	62	30%	28	12%
Total	439		204		235	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by transmission category, Part B Southeast Consortium 9C

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	195	69%	12	40%	5	63%	-	-	-	-	6	75%
Injection drug use (IDU)	11	4%	3	10%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	12	4%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	24	9%	3	10%	1	13%	1	100%	-	-	-	-
Other/unknown	39	14%	12	40%	2	25%	-	-	-	-	2	25%
Subtotal	281	100%	30	100%	8	100%	1	100%	-	-	8	100%
Female adult or adolescent												
Injection drug use	8	10%	-	-	1	33%	-	-	-	-	-	-
Heterosexual contact	53	65%	5	36%	2	67%	-	-	-	-	-	-
Other/unknown	20	25%	9	64%	-	-	-	-	-	-	3	100%
Subtotal	81	100%	14	100%	3	100%	-	-	-	-	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	43%	2	100%	1	100%	-	-	-	-	-	-
Other/unknown	4	57%	-	-	-	-	-	-	-	-	-	-
Subtotal	7	100%	2	100%	1	100%	-	-	-	-	-	-
Total	369		46		12		1		0		11	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by race/ethnicity and by exposure category, Part B Southeast Consortium 9C

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2011											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	181	49%	10	22%	5	42%	-	-	-	-	6	55%
Injection drug use (IDU) only	13	4%	3	7%	1	8%	-	-	-	-	-	-
Heterosexual contact only	77	21%	8	17%	3	25%	1	100%	-	-	-	-
MSM & IDU	11	3%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	6	2%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	14	4%	2	4%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	<1%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	4	1%	2	4%	1	8%	-	-	-	-	-	-
Other/unknown	62	17%	21	46%	2	17%	-	-	-	-	5	45%
Total	369		46		12		1		0		11	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by selected characteristics, Part B Southeast Consortium 9C

Living with a Diagnosis of HIV Infection

Characteristic	2007		2008		2009		2010		2011	
	No.	%								
Sex										
Males	264	75%	280	76%	301	76%	319	77%	334	76%
Females	86	25%	89	24%	93	24%	97	23%	105	24%
Age at end of year										
<13	8	2%	8	2%	9	2%	9	2%	9	2%
13-14	3	1%	3	1%	3	1%	3	1%	3	1%
15-19	6	2%	7	2%	7	2%	7	2%	8	2%
20-24	39	11%	41	11%	43	11%	45	11%	53	12%
25-29	55	16%	57	15%	61	15%	64	15%	65	15%
30-34	69	20%	77	21%	80	20%	85	20%	85	19%
35-39	68	19%	71	19%	77	20%	82	20%	86	20%
40-44	45	13%	45	12%	48	12%	53	13%	53	12%
45-49	30	9%	30	8%	31	8%	32	8%	36	8%
50-54	20	6%	22	6%	24	6%	25	6%	27	6%
55-64	6	2%	6	2%	9	2%	9	2%	11	3%
65+	1	<1%	2	1%	2	1%	2	<1%	3	1%
Race/Ethnicity										
White, not Hispanic	296	85%	311	84%	329	84%	347	83%	369	84%
Black/African American, not Hispanic	37	11%	40	11%	43	11%	45	11%	46	10%
Hispanic/Latino	7	2%	7	2%	11	3%	12	3%	12	3%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	10	3%	10	3%	11	3%	11	3%
Race/Sex										
White, not Hispanic Males	228	65%	241	65%	256	65%	272	65%	286	65%
White, not Hispanic Females	68	19%	70	19%	73	19%	75	18%	83	19%
Black/African American, not Hispanic Males	25	7%	27	7%	30	8%	30	7%	31	7%
Black/African American, not Hispanic Females	12	3%	13	4%	13	3%	15	4%	15	3%
Hispanic/Latino Males	4	1%	4	1%	7	2%	8	2%	8	2%
Hispanic/Latina Females	3	1%	3	1%	4	1%	4	1%	4	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	10	3%	10	3%	11	3%	11	3%
Total	350		369		394		416		439	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by transmission category, Part B Southeast Consortium 9C

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	164	63%	179	65%	193	65%	207	66%	218	66%
Injection drug use (IDU)	14	5%	14	5%	14	5%	14	4%	14	4%
Male-to-male sexual contact and IDU	9	3%	9	3%	10	3%	12	4%	12	4%
Heterosexual contact	27	10%	27	10%	28	9%	28	9%	29	9%
Other/unknown	45	17%	46	17%	50	17%	52	17%	55	17%
Subtotal	259	100%	275	100%	295	100%	313	100%	328	100%
Female adult or adolescent										
Injection drug use	6	7%	7	8%	7	8%	7	8%	9	9%
Heterosexual contact	50	61%	52	61%	55	62%	56	60%	60	59%
Other/unknown	26	32%	26	31%	27	30%	30	32%	32	32%
Subtotal	82	100%	85	100%	89	100%	93	100%	101	100%
Child (<13 yrs at diagnosis)										
Perinatal	5	56%	5	56%	6	60%	6	60%	6	60%
Other/unknown	4	44%	4	44%	4	40%	4	40%	4	40%
Subtotal	9	100%	9	100%	10	100%	10	100%	10	100%
Total	350		369		394		416		439	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported persons living with a diagnosis of HIV infection as of June 30, 2013, by year and by exposure category, Part B Southeast Consortium 9C

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2007		2008		2009		2010		2011	
	No.	%								
Male-to-male sex (MSM) only	148	42%	163	44%	177	45%	191	46%	202	46%
Injection drug use (IDU) only	15	4%	16	4%	16	4%	16	4%	17	4%
Heterosexual contact only	77	22%	79	21%	83	21%	84	20%	89	20%
MSM & IDU	8	2%	8	2%	9	2%	11	3%	11	3%
IDU & Heterosexual contact	5	1%	5	1%	5	1%	5	1%	6	1%
MSM & Heterosexual contact	16	5%	16	4%	16	4%	16	4%	16	4%
MSM & IDU & Heterosexual contact	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Perinatal exposure	6	2%	6	2%	7	2%	7	2%	7	2%
Other/unknown	74	21%	75	20%	80	20%	85	20%	90	21%
Total	350		369		394		416		439	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.