

Ohio HIV Prevention Planning Groups Breakdown

Regional Advisory Groups

<p>Region 1</p> <ul style="list-style-type: none"> Defiance Fulton Henry Lucas Ottawa Sandusky Williams Wood 	<p>Region 7</p> <ul style="list-style-type: none"> Butler Darke Miami Montgomery Preble Warren 	<p>Region 8C</p> <ul style="list-style-type: none"> Adams Brown Clark Clermont Clinton Fayette Greene Highland Jackson Lawrence Madison Pickaway Pike Ross Scioto
<p>Region 2</p> <ul style="list-style-type: none"> Cuyahoga 	<p>Region 8A</p> <ul style="list-style-type: none"> Allen Auglaize Champaign Delaware Hancock Hardin Logan Marion Mercer Paulding Putnam Seneca Shelby Union Van Wert Wyandot 	<p>Region 8D</p> <ul style="list-style-type: none"> Licking Muskingum Guernsey Belmont Fairfield Perry Morgan Noble Monroe Hocking Athens Washington Vinton Meigs Gallia
<p>Region 3</p> <ul style="list-style-type: none"> Ashtabula Geauga Lake Portage Summit Trumbull 	<p>Region 8B</p> <ul style="list-style-type: none"> Ashland Coshocton Crawford Erie Holmes Huron Knox Lorain Medina Morrow Richland Wayne 	
<p>Region 4</p> <ul style="list-style-type: none"> Carroll Columbiana Harrison Jefferson Mahoning Stark Tuscarawas 		
<p>Region 5</p> <ul style="list-style-type: none"> Franklin 		
<p>Region 6</p> <ul style="list-style-type: none"> Hamilton 		

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 1

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 1

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	11.8	47	90%	39	89%	8	100%	-	-
Females	1.2	5	10%	5	11%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	8.9	5	10%	5	11%	-	-	-	-
20-24	21.1	13	25%	13	30%	-	-	-	-
25-29	15.1	8	15%	7	16%	1	13%	-	-
30-34	14.4	7	13%	5	11%	2	25%	-	-
35-39	*	-	-	-	-	-	-	-	-
40-44	14.0	7	13%	5	11%	2	25%	-	-
45-49	10.1	5	10%	4	9%	1	13%	-	-
50-54	*	2	4%	1	2%	1	13%	-	-
55-64	*	4	8%	3	7%	1	13%	-	-
65+	*	1	2%	1	2%	-	-	-	-
Race/Ethnicity									
White, not Hispanic	3.3	21	40%	16	36%	5	63%	-	-
Black/African American, not Hispanic	25.6	23	44%	22	50%	1	13%	-	-
Hispanic/Latino	11.0	6	12%	4	9%	2	25%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	2	4%	2	5%	-	-	-	-
Race/Sex									
White, not Hispanic Males	6.1	19	37%	14	32%	5	63%	-	-
White, not Hispanic Females	*	2	4%	2	5%	-	-	-	-
Black/African American, not Hispanic Males	49.0	21	40%	20	45%	1	13%	-	-
Black/African American, not Hispanic Females	*	2	4%	2	5%	-	-	-	-
Hispanic/Latino Males	18.2	5	10%	3	7%	2	25%	-	-
Hispanic/Latina Females	*	1	2%	1	2%	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	2	4%	2	5%	-	-	-	-
Total	6.4	52		44		8			

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	37	79%	30	77%	7	88%	-	-
Injection drug use (IDU)	1	2%	1	3%	-	-	-	-
Male-to-male sexual contact and IDU	1	2%	1	3%	-	-	-	-
Heterosexual contact	5	11%	5	13%	-	-	-	-
Other/unknown	3	6%	2	5%	1	13%	-	-
Subtotal	47	100%	39	100%	8	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	4	80%	4	80%	-	-	-	-
Other/unknown	1	20%	1	20%	-	-	-	-
Subtotal	5	100%	5	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	52		44		8		0	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	36	69%	29	66%	7	88%	-	-
Injection drug use (IDU) only	1	2%	1	2%	-	-	-	-
Heterosexual contact only	9	17%	9	20%	-	-	-	-
MSM & IDU	1	2%	1	2%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	2%	1	2%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	4	8%	3	7%	1	13%	-	-
Total	52		44		8		-	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	14	74%	16	76%	5	100%	-	-	-	-	2	100%
Injection drug use (IDU)	-	-	1	5%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	11%	3	14%	-	-	-	-	-	-	-	-
Other/unknown	2	11%	1	5%	-	-	-	-	-	-	-	-
Subtotal	19	100%	21	100%	5	100%	-	-	-	-	2	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	2	100%	2	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	21		23		6		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	13	62%	16	70%	5	83%	-	-	-	-	2	100%
Injection drug use (IDU) only	-	-	1	4%	-	-	-	-	-	-	-	-
Heterosexual contact only	4	19%	5	22%	-	-	-	-	-	-	-	-
MSM & IDU	1	5%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	10%	1	4%	1	17%	-	-	-	-	-	-
Total	21		23		6		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 1

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	8.3	33	83%	11.3	45	82%	12.8	51	84%	9.6	38	90%
Females	1.7	7	18%	2.4	10	18%	2.4	10	16%	*	4	10%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	1	2%	*	1	2%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	5%	*	4	7%	*	-	-	*	4	10%
20-24	13.0	8	20%	17.8	11	20%	32.4	20	33%	16.2	10	24%
25-29	*	1	3%	13.2	7	13%	15.1	8	13%	13.2	7	17%
30-34	14.4	7	18%	12.3	6	11%	18.5	9	15%	*	2	5%
35-39	22.0	10	25%	19.8	9	16%	11.0	5	8%	*	3	7%
40-44	*	2	5%	12.0	6	11%	18.0	9	15%	*	2	5%
45-49	*	3	8%	*	4	7%	*	4	7%	10.1	5	12%
50-54	*	3	8%	8.6	5	9%	*	2	3%	*	1	2%
55-64	*	4	10%	*	3	5%	*	2	3%	4.5	5	12%
65+	*	-	-	*	-	-	*	1	2%	*	2	5%
Race/Ethnicity												
White, not Hispanic	3.4	22	55%	3.6	23	42%	5.3	34	56%	2.8	18	43%
Black/African American, not Hispanic	14.5	13	33%	31.2	28	51%	23.4	21	34%	23.4	21	50%
Hispanic/Latino	*	2	5%	*	2	4%	*	3	5%	*	3	7%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	8%	*	2	4%	*	3	5%	*	-	-
Race/Sex												
White, not Hispanic Males	6.1	19	48%	7.0	22	40%	8.9	28	46%	4.8	15	36%
White, not Hispanic Females	*	3	8%	*	1	2%	1.8	6	10%	*	3	7%
Black/African American, not Hispanic Males	21.0	9	23%	44.3	19	35%	42.0	18	30%	46.7	20	48%
Black/African American, not Hispanic Females	*	4	10%	19.2	9	16%	*	3	5%	*	1	2%
Hispanic/Latino Males	*	2	5%	*	2	4%	*	2	3%	*	3	7%
Hispanic/Latina Females	*	-	-	*	-	-	*	1	2%	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	8%	*	2	4%	*	3	5%	*	-	-
Total	4.9	40		6.8	55		7.5	61		5.2	42	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 1

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	28	85%	32	71%	36	71%	32	86%
Injection drug use (IDU)	-	-	1	2%	1	2%	1	3%
Male-to-male sexual contact and IDU	1	3%	2	4%	4	8%	1	3%
Heterosexual contact	3	9%	4	9%	-	-	1	3%
Other/unknown	1	3%	6	13%	10	20%	2	5%
Subtotal	33	100%	45	100%	51	100%	37	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	1	11%	-	-
Heterosexual contact	7	100%	10	100%	7	78%	4	100%
Other/unknown	-	-	-	-	1	11%	-	-
Subtotal	7	100%	10	100%	9	100%	4	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%
Total	40		55		61		42	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 1

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	27	68%	31	56%	35	57%	30	71%
Injection drug use (IDU) only	-	-	1	2%	-	-	1	2%
Heterosexual contact only	10	25%	14	25%	7	11%	5	12%
MSM & IDU	1	3%	1	2%	4	7%	1	2%
IDU & Heterosexual contact	-	-	-	-	2	3%	-	-
MSM & Heterosexual contact	1	3%	1	2%	1	2%	2	5%
MSM & IDU & Heterosexual contact	-	-	1	2%	-	-	-	-
Perinatal exposure	-	-	-	-	1	2%	1	2%
Other/unknown	1	3%	6	11%	11	18%	2	5%
Total	40		55		61		42	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 1

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	20	83%	27	82%	40	87%	28	90%	39	89%
Females	4	17%	6	18%	6	13%	3	10%	5	11%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	1	2%	1	3%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	2	8%	3	9%	-	-	4	13%	5	11%
20-24	7	29%	10	30%	18	39%	6	19%	13	30%
25-29	1	4%	6	18%	7	15%	5	16%	7	16%
30-34	2	8%	4	12%	7	15%	1	3%	5	11%
35-39	7	29%	3	9%	4	9%	1	3%	-	-
40-44	1	4%	3	9%	5	11%	2	6%	5	11%
45-49	-	-	2	6%	2	4%	5	16%	4	9%
50-54	1	4%	1	3%	1	2%	1	3%	1	2%
55-64	3	13%	1	3%	-	-	4	13%	3	7%
65+	-	-	-	-	1	2%	1	3%	1	2%
Race/Ethnicity										
White, not Hispanic	11	46%	13	39%	26	57%	12	39%	16	36%
Black/African American, not Hispanic	10	42%	17	52%	17	37%	18	58%	22	50%
Hispanic/Latino	-	-	1	3%	2	4%	1	3%	4	9%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	3	13%	2	6%	1	2%	-	-	2	5%
Race/Sex										
White, not Hispanic Males	9	38%	12	36%	23	50%	10	32%	14	32%
White, not Hispanic Females	2	8%	1	3%	3	7%	2	6%	2	5%
Black/African American, not Hispanic Males	8	33%	12	36%	14	30%	17	55%	20	45%
Black/African American, not Hispanic Females	2	8%	5	15%	3	7%	1	3%	2	5%
Hispanic/Latino Males	-	-	1	3%	2	4%	1	3%	3	7%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	1	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	13%	2	6%	1	2%	-	-	2	5%
Total	24		33		46		31		44	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	15	75%	21	78%	29	73%	23	85%	30	77%
Injection drug use (IDU)	-	-	-	-	1	3%	1	4%	1	3%
Male-to-male sexual contact and IDU	1	5%	2	7%	4	10%	-	-	1	3%
Heterosexual contact	3	15%	1	4%	-	-	1	4%	5	13%
Other/unknown	1	5%	3	11%	6	15%	2	7%	2	5%
Subtotal	20	100%	27	100%	40	100%	27	100%	39	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	1	20%	-	-	-	-
Heterosexual contact	4	100%	6	100%	4	80%	3	100%	4	80%
Other/unknown	-	-	-	-	-	-	-	-	1	20%
Subtotal	4	100%	6	100%	5	100%	3	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	-	-
Total	24		33		46		31		44	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	58%	21	64%	28	61%	21	68%	29	66%
Injection drug use (IDU) only	-	-	-	-	-	-	1	3%	1	2%
Heterosexual contact only	7	29%	7	21%	4	9%	4	13%	9	20%
MSM & IDU	1	4%	1	3%	4	9%	-	-	1	2%
IDU & Heterosexual contact	-	-	-	-	2	4%	-	-	-	-
MSM & Heterosexual contact	1	4%	-	-	1	2%	2	6%	1	2%
MSM & IDU & Heterosexual contact	-	-	1	3%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	1	2%	1	3%	-	-
Other/unknown	1	4%	3	9%	6	13%	2	6%	3	7%
Total	24		33		46		31		44	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 1

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	24	77%	29	83%	15	79%	15	88%	9	100%	897	80%
Females	7	23%	6	17%	4	21%	2	12%	-	-	223	20%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	19	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	3	10%	1	3%	-	-	-	-	-	-	33	3%
20-24	1	3%	3	9%	2	11%	6	35%	-	-	108	10%
25-29	5	16%	2	6%	1	5%	4	24%	1	11%	180	16%
30-34	6	19%	5	14%	3	16%	2	12%	2	22%	219	20%
35-39	4	13%	8	23%	3	16%	2	12%	-	-	211	19%
40-44	5	16%	6	17%	4	21%	1	6%	3	33%	138	12%
45-49	4	13%	3	9%	3	16%	-	-	1	11%	92	8%
50-54	2	6%	4	11%	1	5%	-	-	1	11%	64	6%
55-64	1	3%	3	9%	2	11%	1	6%	1	11%	46	4%
65+	-	-	-	-	-	-	1	6%	-	-	9	1%
Race/Ethnicity												
White, not Hispanic	18	58%	16	46%	11	58%	10	59%	6	67%	664	59%
Black/African American, not Hispanic	9	29%	16	46%	4	21%	3	18%	1	11%	378	34%
Hispanic/Latino	4	13%	3	9%	2	11%	4	24%	2	22%	62	6%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	2	11%	-	-	-	-	11	1%
Race/Sex												
White, not Hispanic Males	14	45%	15	43%	8	42%	9	53%	6	67%	580	52%
White, not Hispanic Females	4	13%	1	3%	3	16%	1	6%	-	-	84	8%
Black/African American, not Hispanic Males	6	19%	11	31%	4	21%	3	18%	1	11%	259	23%
Black/African American, not Hispanic Females	3	10%	5	14%	-	-	-	-	-	-	119	11%
Hispanic/Latino Males	4	13%	3	9%	1	5%	3	18%	2	22%	45	4%
Hispanic/Latina Females	-	-	-	-	1	5%	1	6%	-	-	17	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	2	11%	-	-	-	-	11	1%
Total	31		35		19		17		9		1,120	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	20	83%	19	66%	10	67%	13	87%	7	78%	647	73%
Injection drug use (IDU)	-	-	1	3%	-	-	-	-	-	-	47	5%
Male-to-male sexual contact and IDU	1	4%	1	3%	1	7%	1	7%	1	11%	68	8%
Heterosexual contact	3	13%	3	10%	-	-	-	-	-	-	58	7%
Other/unknown	-	-	5	17%	4	27%	1	7%	1	11%	67	8%
Subtotal	24	100%	29	100%	15	100%	15	100%	9	100%	887	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	31	14%
Heterosexual contact	7	100%	6	100%	3	75%	2	100%	-	-	170	79%
Other/unknown	-	-	-	-	1	25%	-	-	-	-	13	6%
Subtotal	7	100%	6	100%	4	100%	2	100%	-	-	214	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	16	84%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	16%
Subtotal	-	-	-	-	-	-	-	-	-	-	19	100%
Total	31		35		19		17		9		1,120	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	20	65%	18	51%	9	47%	13	76%	7	78%	609	54%
Injection drug use (IDU) only	-	-	1	3%	-	-	-	-	-	-	39	3%
Heterosexual contact only	10	32%	9	26%	3	16%	2	12%	-	-	228	20%
MSM & IDU	1	3%	1	3%	1	5%	1	6%	1	11%	52	5%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	39	3%
MSM & Heterosexual contact	-	-	1	3%	1	5%	-	-	-	-	38	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	16	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	16	1%
Other/unknown	-	-	5	14%	5	26%	1	6%	1	11%	83	7%
Total	31		35		19		17		9		1,120	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 1

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 1

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	221.8	881	75%	442	74%	439	76%
Females	70.8	294	25%	157	26%	137	24%
Age at end of year							
<13	5.3	7	1%	5	1%	2	<1%
13-14	*	2	<1%	-	-	2	<1%
15-19	21.4	12	1%	10	2%	2	<1%
20-24	76.2	47	4%	40	7%	7	1%
25-29	139.8	74	6%	61	10%	13	2%
30-34	145.6	71	6%	50	8%	21	4%
35-39	237.6	108	9%	62	10%	46	8%
40-44	365.0	182	15%	84	14%	98	17%
45-49	388.8	193	16%	90	15%	103	18%
50-54	361.0	209	18%	84	14%	125	22%
55-64	193.8	217	18%	89	15%	128	22%
65+	42.2	53	5%	24	4%	29	5%
Race/Ethnicity							
White, not Hispanic	92.2	590	50%	295	49%	295	51%
Black/African American, not Hispanic	528.2	474	40%	248	41%	226	39%
Hispanic/Latino	137.3	75	6%	32	5%	43	7%
Asian/Pacific Islander	64.0	7	1%	5	1%	2	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	28	2%	19	3%	9	2%
Race/Sex							
White, not Hispanic Males	153.9	482	41%	229	38%	253	44%
White, not Hispanic Females	33.0	108	9%	66	11%	42	7%
Black/African American, not Hispanic Males	732.9	314	27%	168	28%	146	25%
Black/African American, not Hispanic Females	341.2	160	14%	80	13%	80	14%
Hispanic/Latino Males	197.0	54	5%	24	4%	30	5%
Hispanic/Latina Females	77.1	21	2%	8	1%	13	2%
Asian/Pacific Islander Males	113.0	6	1%	4	1%	2	<1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	28	2%	19	3%	9	2%
Total	144.6	1,175		599		576	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	624	71%	309	71%	315	72%
Injection drug use (IDU)	41	5%	16	4%	25	6%
Male-to-male sexual contact and IDU	56	6%	30	7%	26	6%
Heterosexual contact	80	9%	38	9%	42	10%
Other/unknown	72	8%	44	10%	28	6%
Subtotal	873	100%	437	100%	436	100%
Female adult or adolescent						
Injection drug use	26	9%	10	7%	16	12%
Heterosexual contact	233	82%	125	82%	108	83%
Other/unknown	24	8%	18	12%	6	5%
Subtotal	283	100%	153	100%	130	100%
Child (<13 yrs at diagnosis)						
Perinatal	19	100%	9	100%	10	100%
Other/unknown	-	-	-	-	-	-
Subtotal	19	100%	9	100%	10	100%
Total	1,175		599		576	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	575	49%	286	48%	289	50%
Injection drug use (IDU) only	29	2%	13	2%	16	3%
Heterosexual contact only	313	27%	163	27%	150	26%
MSM & IDU	41	3%	26	4%	15	3%
IDU & Heterosexual contact	38	3%	13	2%	25	4%
MSM & Heterosexual contact	49	4%	23	4%	26	5%
MSM & IDU & Heterosexual contact	15	1%	4	1%	11	2%
Perinatal exposure	19	2%	9	2%	10	2%
Other/unknown	96	8%	62	10%	34	6%
Total	1,175		599		576	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 1

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	373	78%	188	61%	44	81%	3	50%	-	-	16	67%
Injection drug use (IDU)	19	4%	18	6%	3	6%	-	-	-	-	1	4%
Male-to-male sexual contact and IDU	35	7%	16	5%	3	6%	-	-	-	-	2	8%
Heterosexual contact	25	5%	51	16%	2	4%	-	-	-	-	2	8%
Other/unknown	27	6%	37	12%	2	4%	3	50%	-	-	3	13%
Subtotal	479	100%	310	100%	54	100%	6	100%	-	-	24	100%
Female adult or adolescent												
Injection drug use	8	8%	18	12%	-	-	-	-	-	-	-	-
Heterosexual contact	94	89%	121	79%	16	80%	1	100%	1	100%	-	-
Other/unknown	4	4%	14	9%	4	20%	-	-	-	-	2	100%
Subtotal	106	100%	153	100%	20	100%	1	100%	1	100%	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	5	100%	11	100%	1	100%	-	-	-	-	2	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	5	100%	11	100%	1	100%	-	-	-	-	2	100%
Total	590		474		75		7		1		28	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 1

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	346	59%	169	36%	41	55%	3	43%	-	-	16	57%
Injection drug use (IDU) only	13	2%	16	3%	-	-	-	-	-	-	-	-
Heterosexual contact only	119	20%	172	36%	18	24%	1	14%	1	100%	2	7%
MSM & IDU	23	4%	15	3%	2	3%	-	-	-	-	1	4%
IDU & Heterosexual contact	14	2%	20	4%	3	4%	-	-	-	-	1	4%
MSM & Heterosexual contact	27	5%	19	4%	3	4%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	12	2%	1	0%	1	1%	-	-	-	-	1	4%
Perinatal exposure	5	1%	11	2%	1	1%	-	-	-	-	2	7%
Other/unknown	31	5%	51	11%	6	8%	3	43%	-	-	5	18%
Total	590		474		75		7		1		28	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 1

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	711	73%	752	73%	798	74%	834	74%	881	75%
Females	266	27%	276	27%	286	26%	290	26%	294	25%
Age at end of year										
<13	7	1%	7	1%	8	1%	8	1%	7	1%
13-14	4	<1%	2	<1%	1	<1%	2	<1%	2	<1%
15-19	6	1%	9	1%	10	1%	12	1%	12	1%
20-24	31	3%	37	4%	48	4%	47	4%	47	4%
25-29	46	5%	50	5%	58	5%	61	5%	74	6%
30-34	81	8%	75	7%	68	6%	73	6%	71	6%
35-39	142	15%	131	13%	127	12%	116	10%	108	9%
40-44	179	18%	181	18%	189	17%	184	16%	182	15%
45-49	193	20%	202	20%	199	18%	197	18%	193	16%
50-54	138	14%	163	16%	178	16%	189	17%	209	18%
55-64	129	13%	144	14%	163	15%	195	17%	217	18%
65+	21	2%	27	3%	35	3%	40	4%	53	5%
Race/Ethnicity										
White, not Hispanic	500	51%	522	51%	550	51%	569	51%	590	50%
Black/African American, not Hispanic	387	40%	412	40%	433	40%	452	40%	474	40%
Hispanic/Latino	61	6%	63	6%	67	6%	69	6%	75	6%
Asian/Pacific Islander	7	1%	7	1%	7	1%	7	1%	7	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	21	2%	23	2%	26	2%	26	2%	28	2%
Race/Sex										
White, not Hispanic Males	404	41%	424	41%	447	41%	463	41%	482	41%
White, not Hispanic Females	96	10%	98	10%	103	10%	106	9%	108	9%
Black/African American, not Hispanic Males	241	25%	258	25%	275	25%	293	26%	314	27%
Black/African American, not Hispanic Females	146	15%	154	15%	158	15%	159	14%	160	14%
Hispanic/Latino Males	42	4%	44	4%	47	4%	49	4%	54	5%
Hispanic/Latina Females	19	2%	19	2%	20	2%	20	2%	21	2%
Asian/Pacific Islander Males	6	1%	6	1%	6	1%	6	1%	6	1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	21	2%	23	2%	26	2%	26	2%	28	2%
Total	977		1,028		1,084		1,124		1,175	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 1

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	495	70%	525	70%	558	71%	588	71%	624	71%
Injection drug use (IDU)	37	5%	38	5%	39	5%	40	5%	41	5%
Male-to-male sexual contact and IDU	52	7%	52	7%	55	7%	55	7%	56	6%
Heterosexual contact	70	10%	74	10%	74	9%	75	9%	80	9%
Other/unknown	51	7%	57	8%	65	8%	68	8%	72	8%
Subtotal	705	100%	746	100%	791	100%	826	100%	873	100%
Female adult or adolescent										
Injection drug use	25	10%	25	9%	26	9%	26	9%	26	9%
Heterosexual contact	208	81%	218	82%	226	82%	230	82%	233	82%
Other/unknown	23	9%	23	9%	23	8%	23	8%	24	8%
Subtotal	256	100%	266	100%	275	100%	279	100%	283	100%
Child (<13 yrs at diagnosis)										
Perinatal	16	100%	16	100%	18	100%	19	100%	19	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	16	100%	16	100%	18	100%	19	100%	19	100%
Total	977		1,028		1,084		1,124		1,175	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 1

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	452	46%	481	47%	512	47%	540	48%	575	49%
Injection drug use (IDU) only	26	3%	27	3%	27	2%	28	2%	29	2%
Heterosexual contact only	278	28%	292	28%	300	28%	305	27%	313	27%
MSM & IDU	37	4%	37	4%	40	4%	40	4%	41	3%
IDU & Heterosexual contact	36	4%	36	4%	38	4%	38	3%	38	3%
MSM & Heterosexual contact	43	4%	44	4%	46	4%	48	4%	49	4%
MSM & IDU & Heterosexual contact	15	2%	15	1%	15	1%	15	1%	15	1%
Perinatal exposure	16	2%	16	2%	18	2%	19	2%	19	2%
Other/unknown	74	8%	80	8%	88	8%	91	8%	96	8%
Total	977		1,028		1,084		1,124		1,175	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 2

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 2

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	31.0	186	89%	154	90%	21	81%	11	85%	
Females	3.6	24	11%	17	10%	5	19%	2	15%	
Age at diagnosis (yr)										
<13	*	1	<1%	1	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	18.7	15	7%	13	8%	1	4%	1	8%	
20-24	70.8	60	29%	56	33%	3	12%	1	8%	
25-29	40.2	34	16%	30	18%	2	8%	2	15%	
30-34	34.6	28	13%	22	13%	5	19%	1	8%	
35-39	32.2	23	11%	19	11%	3	12%	1	8%	
40-44	22.6	17	8%	11	6%	5	19%	1	8%	
45-49	16.0	13	6%	10	6%	1	4%	2	15%	
50-54	7.5	7	3%	4	2%	3	12%	-	-	
55-64	5.6	10	5%	5	3%	3	12%	2	15%	
65+	*	2	1%	-	-	-	-	2	15%	
Race/Ethnicity										
White, not Hispanic	4.5	34	16%	29	17%	4	15%	1	8%	
Black/African American, not Hispanic	41.7	155	74%	125	73%	19	73%	11	85%	
Hispanic/Latino	19.2	13	6%	9	5%	3	12%	1	8%	
Asian/Pacific Islander	*	2	1%	2	1%	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	6	3%	6	4%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	7.6	28	13%	24	14%	3	12%	1	8%	
White, not Hispanic Females	1.5	6	3%	5	3%	1	4%	-	-	
Black/African American, not Hispanic Males	81.8	138	66%	114	67%	15	58%	9	69%	
Black/African American, not Hispanic Females	8.4	17	8%	11	6%	4	15%	2	15%	
Hispanic/Latino Males	35.8	12	6%	8	5%	3	12%	1	8%	
Hispanic/Latina Females	*	1	<1%	1	1%	-	-	-	-	
Asian/Pacific Islander Males	*	2	1%	2	1%	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	6	3%	6	4%	-	-	-	-	
Total	16.7	210		171		26		13		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	122	66%	104	68%	12	57%	6	55%
Injection drug use (IDU)	1	1%	-	-	1	5%	-	-
Male-to-male sexual contact and IDU	2	1%	2	1%	-	-	-	-
Heterosexual contact	11	6%	9	6%	1	5%	1	9%
Other/unknown	49	26%	38	25%	7	33%	4	36%
Subtotal	185	100%	153	100%	21	100%	11	100%
Female adult or adolescent								
Injection drug use	3	13%	2	12%	1	20%	-	-
Heterosexual contact	16	67%	13	76%	3	60%	-	-
Other/unknown	5	21%	2	12%	1	20%	2	100%
Subtotal	24	100%	17	100%	5	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	210		171		26		13	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	121	58%	103	60%	12	46%	6	46%
Injection drug use (IDU) only	2	1%	1	1%	1	4%	-	-
Heterosexual contact only	27	13%	22	13%	4	15%	1	8%
MSM & IDU	2	1%	2	1%	-	-	-	-
IDU & Heterosexual contact	2	1%	1	1%	1	4%	-	-
MSM & Heterosexual contact	1	<1%	1	1%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	55	26%	41	24%	8	31%	6	46%
Total	210		171		26		13	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	16	57%	94	69%	9	75%	1	50%	-	-	2	33%
Injection drug use (IDU)	-	-	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	4%	-	-	1	8%	-	-	-	-	-	-
Heterosexual contact	2	7%	7	5%	-	-	-	-	-	-	2	33%
Other/unknown	9	32%	35	26%	2	17%	1	50%	-	-	2	33%
Subtotal	28	100%	137	100%	12	100%	2	100%	-	-	6	100%
Female adult or adolescent												
Injection drug use	3	50%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	33%	13	76%	1	100%	-	-	-	-	-	-
Other/unknown	1	17%	4	24%	-	-	-	-	-	-	-	-
Subtotal	6	100%	17	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	34		155		13		2		0		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	47%	93	60%	9	69%	1	50%	-	-	2	33%
Injection drug use (IDU) only	1	3%	1	1%	-	-	-	-	-	-	-	-
Heterosexual contact only	4	12%	20	13%	1	8%	-	-	-	-	2	33%
MSM & IDU	1	3%	-	-	1	8%	-	-	-	-	-	-
IDU & Heterosexual contact	2	6%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	1	1%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	10	29%	40	26%	2	15%	1	50%	-	-	2	33%
Total	34		155		13		2		0		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 2

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	23.2	139	82%	26.2	157	79%	28.3	170	82%	31.5	189	83%
Females	4.5	30	18%	6.2	41	21%	5.8	38	18%	5.9	39	17%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	1	<1%
13-14	*	1	1%	*	0	<1%	*	-	-	*	1	<1%
15-19	23.7	19	11%	15.0	12	6%	16.2	13	6%	21.2	17	7%
20-24	38.9	33	20%	44.8	38	19%	48.3	41	20%	57.8	49	21%
25-29	24.8	21	12%	31.9	27	14%	36.7	31	15%	41.4	35	15%
30-34	28.4	23	14%	33.4	27	14%	39.5	32	15%	39.5	32	14%
35-39	23.8	17	10%	14.0	10	5%	29.4	21	10%	29.4	21	9%
40-44	18.6	14	8%	34.6	26	13%	22.6	17	8%	25.3	19	8%
45-49	19.7	16	9%	32.0	26	13%	22.2	18	9%	28.3	23	10%
50-54	12.8	12	7%	16.0	15	8%	21.3	20	10%	17.0	16	7%
55-64	7.3	13	8%	8.4	15	8%	6.7	12	6%	6.7	12	5%
65+	*	-	-	*	2	1%	*	3	1%	*	2	1%
Race/Ethnicity												
White, not Hispanic	5.4	41	24%	6.3	48	24%	6.1	46	22%	7.5	57	25%
Black/African American, not Hispanic	28.8	107	63%	35.0	130	66%	33.4	124	60%	37.1	138	61%
Hispanic/Latino	19.2	13	8%	19.2	13	7%	35.5	24	12%	31.0	21	9%
Asian/Pacific Islander	*	1	1%	*	1	1%	*	1	<1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	6	3%	*	13	6%	*	12	5%
Race/Sex												
White, not Hispanic Males	10.3	38	22%	11.2	41	21%	11.2	41	20%	13.6	50	22%
White, not Hispanic Females	*	3	2%	1.8	7	4%	1.3	5	2%	1.8	7	3%
Black/African American, not Hispanic Males	49.2	83	49%	59.3	100	51%	57.5	97	47%	67.0	113	50%
Black/African American, not Hispanic Females	11.8	24	14%	14.8	30	15%	13.3	27	13%	12.3	25	11%
Hispanic/Latino Males	29.8	10	6%	29.8	10	5%	62.7	21	10%	50.7	17	7%
Hispanic/Latina Females	*	3	2%	*	3	2%	*	3	1%	*	4	2%
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	<1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	6	3%	*	13	6%	*	12	5%
Total	13.4	169		15.7	198		16.5	208		18.1	228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 2

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	96	70%	118	75%	116	68%	120	63%
Injection drug use (IDU)	1	1%	3	2%	4	2%	3	2%
Male-to-male sexual contact and IDU	1	1%	1	1%	10	6%	2	1%
Heterosexual contact	3	2%	5	3%	5	3%	9	5%
Other/unknown	37	27%	30	19%	35	21%	55	29%
Subtotal	138	100%	157	100%	170	100%	189	100%
Female adult or adolescent								
Injection drug use	4	13%	2	5%	2	5%	3	8%
Heterosexual contact	14	47%	23	56%	17	46%	20	54%
Other/unknown	12	40%	16	39%	18	49%	14	38%
Subtotal	30	100%	41	100%	37	100%	37	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	2	100%
Subtotal	1	100%	-	-	1	100%	2	100%
Total	169		198		208		228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 2

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	96	57%	118	60%	113	54%	117	51%
Injection drug use (IDU) only	1	1%	3	2%	5	2%	5	2%
Heterosexual contact only	17	10%	28	14%	22	11%	29	13%
MSM & IDU	1	1%	1	1%	6	3%	2	1%
IDU & Heterosexual contact	4	2%	2	1%	1	<1%	1	<1%
MSM & Heterosexual contact	-	-	-	-	3	1%	3	1%
MSM & IDU & Heterosexual contact	-	-	-	-	4	2%	-	-
Perinatal exposure	1	1%	-	-	1	<1%	1	<1%
Other/unknown	49	29%	46	23%	53	25%	70	31%
Total	169		198		208		228	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 2

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	94	80%	114	83%	131	83%	137	83%	154	90%
Females	23	20%	24	17%	27	17%	29	17%	17	10%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	1	1%	1	1%
13-14	1	1%	-	-	-	-	1	1%	-	-
15-19	15	13%	11	8%	13	8%	17	10%	13	8%
20-24	24	21%	34	25%	34	22%	38	23%	56	33%
25-29	16	14%	20	14%	29	18%	25	15%	30	18%
30-34	15	13%	19	14%	25	16%	25	15%	22	13%
35-39	13	11%	8	6%	14	9%	15	9%	19	11%
40-44	8	7%	17	12%	11	7%	12	7%	11	6%
45-49	10	9%	8	6%	11	7%	15	9%	10	6%
50-54	6	5%	12	9%	11	7%	10	6%	4	2%
55-64	9	8%	8	6%	9	6%	6	4%	5	3%
65+	-	-	1	1%	1	1%	1	1%	-	-
Race/Ethnicity										
White, not Hispanic	26	22%	30	22%	37	23%	41	25%	29	17%
Black/African American, not Hispanic	78	67%	91	66%	94	59%	101	61%	125	73%
Hispanic/Latino	7	6%	12	9%	14	9%	13	8%	9	5%
Asian/Pacific Islander	1	1%	1	1%	1	1%	-	-	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	5	4%	4	3%	12	8%	11	7%	6	4%
Race/Sex										
White, not Hispanic Males	24	21%	26	19%	33	21%	38	23%	24	14%
White, not Hispanic Females	2	2%	4	3%	4	3%	3	2%	5	3%
Black/African American, not Hispanic Males	58	50%	75	54%	76	48%	81	49%	114	67%
Black/African American, not Hispanic Females	20	17%	16	12%	18	11%	20	12%	11	6%
Hispanic/Latino Males	6	5%	9	7%	11	7%	10	6%	8	5%
Hispanic/Latina Females	1	1%	3	2%	3	2%	3	2%	1	1%
Asian/Pacific Islander Males	1	1%	1	1%	1	1%	-	-	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	5	4%	4	3%	12	8%	11	7%	6	4%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	68	73%	87	76%	94	72%	87	64%	104	68%
Injection drug use (IDU)	-	-	3	3%	1	1%	2	1%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	7	5%	2	1%	2	1%
Heterosexual contact	3	3%	5	4%	3	2%	4	3%	9	6%
Other/unknown	22	24%	19	17%	26	20%	42	31%	38	25%
Subtotal	93	100%	114	100%	131	100%	137	100%	153	100%
Female adult or adolescent										
Injection drug use	2	9%	1	4%	1	4%	2	7%	2	12%
Heterosexual contact	13	57%	14	58%	13	50%	15	56%	13	76%
Other/unknown	8	35%	9	38%	12	46%	10	37%	2	12%
Subtotal	23	100%	24	100%	26	100%	27	100%	17	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	2	100%	1	100%
Subtotal	1	100%	-	-	1	100%	2	100%	1	100%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	68	58%	87	63%	91	58%	84	51%	103	60%
Injection drug use (IDU) only	-	-	3	2%	2	1%	4	2%	1	1%
Heterosexual contact only	16	14%	19	14%	16	10%	19	11%	22	13%
MSM & IDU	-	-	-	-	5	3%	2	1%	2	1%
IDU & Heterosexual contact	2	2%	1	1%	-	-	-	-	1	1%
MSM & Heterosexual contact	-	-	-	-	3	2%	3	2%	1	1%
MSM & IDU & Heterosexual contact	-	-	-	-	2	1%	-	-	-	-
Perinatal exposure	1	1%	-	-	1	1%	1	1%	-	-
Other/unknown	30	26%	28	20%	38	24%	53	32%	41	24%
Total	117		138		158		166		171	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 2

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	67	74%	65	78%	54	72%	76	84%	43	83%	3,531	82%
Females	24	26%	18	22%	21	28%	15	16%	9	17%	771	18%
Age at AIDS diagnosis (yr)												
<13	2	2%	1	1%	1	1%	-	-	-	-	51	1%
13-14	1	1%	-	-	-	-	-	-	-	-	6	<1%
15-19	5	5%	4	5%	3	4%	2	2%	2	4%	91	2%
20-24	14	15%	10	12%	9	12%	13	14%	8	15%	363	8%
25-29	12	13%	10	12%	4	5%	18	20%	8	15%	719	17%
30-34	18	20%	10	12%	8	11%	12	13%	7	13%	878	20%
35-39	10	11%	5	6%	11	15%	10	11%	5	10%	827	19%
40-44	8	9%	11	13%	11	15%	9	10%	7	13%	557	13%
45-49	10	11%	21	25%	10	13%	10	11%	3	6%	373	9%
50-54	6	7%	4	5%	11	15%	7	8%	5	10%	203	5%
55-64	5	5%	6	7%	5	7%	8	9%	5	10%	183	4%
65+	-	-	1	1%	2	3%	2	2%	2	4%	51	1%
Race/Ethnicity												
White, not Hispanic	24	26%	20	24%	12	16%	24	26%	12	23%	1,545	36%
Black/African American, not Hispanic	59	65%	59	71%	52	69%	48	53%	34	65%	2,362	55%
Hispanic/Latino	7	8%	3	4%	10	13%	12	13%	4	8%	329	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	5	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	5	<1%
Unknown	1	1%	1	1%	1	1%	7	8%	2	4%	56	1%
Race/Sex												
White, not Hispanic Males	20	22%	18	22%	10	13%	18	20%	11	21%	1,425	33%
White, not Hispanic Females	4	4%	2	2%	2	3%	6	7%	1	2%	120	3%
Black/African American, not Hispanic Males	42	46%	43	52%	34	45%	42	46%	26	50%	1,816	42%
Black/African American, not Hispanic Females	17	19%	16	19%	18	24%	6	7%	8	15%	546	13%
Hispanic/Latino Males	4	4%	3	4%	9	12%	11	12%	4	8%	240	6%
Hispanic/Latina Females	3	3%	-	-	1	1%	1	1%	-	-	89	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	5	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	1	1%	1	1%	1	1%	7	8%	2	4%	56	1%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	44	66%	41	64%	32	59%	52	68%	26	60%	2,391	68%
Injection drug use (IDU)	2	3%	1	2%	3	6%	4	5%	1	2%	346	10%
Male-to-male sexual contact and IDU	3	4%	2	3%	4	7%	-	-	2	5%	268	8%
Heterosexual contact	2	3%	1	2%	2	4%	6	8%	4	9%	190	5%
Other/unknown	16	24%	19	30%	13	24%	14	18%	10	23%	316	9%
Subtotal	67	100%	64	100%	54	100%	76	100%	43	100%	3,511	100%
Female adult or adolescent												
Injection drug use	5	23%	1	6%	1	5%	2	13%	1	11%	182	25%
Heterosexual contact	9	41%	12	67%	10	50%	9	60%	3	33%	472	64%
Other/unknown	8	36%	5	28%	9	45%	4	27%	5	56%	86	12%
Subtotal	22	100%	18	100%	20	100%	15	100%	9	100%	740	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	1	100%	1	100%	-	-	-	-	46	90%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	10%
Subtotal	2	100%	1	100%	1	100%	-	-	-	-	51	100%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	41	45%	39	47%	32	43%	52	57%	26	50%	2,251	52%
Injection drug use (IDU) only	5	5%	1	1%	3	4%	3	3%	1	2%	318	7%
Heterosexual contact only	11	12%	13	16%	12	16%	15	16%	7	13%	662	15%
MSM & IDU	3	3%	2	2%	2	3%	-	-	2	4%	220	5%
IDU & Heterosexual contact	2	2%	1	1%	1	1%	3	3%	1	2%	210	5%
MSM & Heterosexual contact	3	3%	2	2%	-	-	-	-	-	-	138	3%
MSM & IDU & Heterosexual contact	-	-	-	-	2	3%	-	-	-	-	48	1%
Perinatal exposure	2	2%	1	1%	1	1%	-	-	-	-	46	1%
Other/unknown	24	26%	24	29%	22	29%	18	20%	15	29%	409	10%
Total	91		83		75		91		52		4,302	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 2

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 2

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	572.1	3,432	78%	1,766	78%	1,666	78%
Females	143.8	949	22%	489	22%	460	22%
Age at end of year							
<13	6.3	12	<1%	9	<1%	3	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	46.1	37	1%	32	1%	5	<1%
20-24	260.6	221	5%	178	8%	43	2%
25-29	350.1	296	7%	223	10%	73	3%
30-34	436.1	353	8%	241	11%	112	5%
35-39	478.6	342	8%	209	9%	133	6%
40-44	648.5	488	11%	259	11%	229	11%
45-49	851.5	691	16%	312	14%	379	18%
50-54	851.8	800	18%	334	15%	466	22%
55-64	511.4	911	21%	367	16%	544	26%
65+	110.6	229	5%	91	4%	138	6%
Race/Ethnicity							
White, not Hispanic	172.2	1,306	30%	655	29%	651	31%
Black/African American, not Hispanic	674.1	2,506	57%	1,304	58%	1,202	57%
Hispanic/Latino	572.0	387	9%	174	8%	213	10%
Asian/Pacific Islander	39.8	15	<1%	11	<1%	4	<1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	163	4%	109	5%	54	3%
Race/Sex							
White, not Hispanic Males	316.9	1,164	27%	579	26%	585	28%
White, not Hispanic Females	36.3	142	3%	76	3%	66	3%
Black/African American, not Hispanic Males	1,102.0	1,859	42%	968	43%	891	42%
Black/African American, not Hispanic Females	318.6	647	15%	336	15%	311	15%
Hispanic/Latino Males	790.6	265	6%	122	5%	143	7%
Hispanic/Latina Females	357.3	122	3%	52	2%	70	3%
Asian/Pacific Islander Males	64.9	12	<1%	9	<1%	3	<1%
Asian/Pacific Islander Females	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Males	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	163	4%	109	5%	54	3%
Total	347.7	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,317	68%	1,170	67%	1,147	69%
Injection drug use (IDU)	159	5%	54	3%	105	6%
Male-to-male sexual contact and IDU	163	5%	53	3%	110	7%
Heterosexual contact	231	7%	111	6%	120	7%
Other/unknown	537	16%	361	21%	176	11%
Subtotal	3,407	100%	1,749	100%	1,658	100%
Female adult or adolescent						
Injection drug use	124	13%	40	8%	84	19%
Heterosexual contact	593	64%	289	61%	304	68%
Other/unknown	203	22%	146	31%	57	13%
Subtotal	920	100%	475	100%	445	100%
Child (<13 yrs at diagnosis)						
Perinatal	44	81%	22	71%	22	96%
Other/unknown	10	19%	9	29%	1	4%
Subtotal	54	100%	31	100%	23	100%
Total	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,197	50%	1,128	50%	1,069	50%
Injection drug use (IDU) only	130	3%	50	2%	80	4%
Heterosexual contact only	824	19%	400	18%	424	20%
MSM & IDU	125	3%	48	2%	77	4%
IDU & Heterosexual contact	153	3%	44	2%	109	5%
MSM & Heterosexual contact	120	3%	42	2%	78	4%
MSM & IDU & Heterosexual contact	38	1%	5	<1%	33	2%
Perinatal exposure	45	1%	23	1%	22	1%
Other/unknown	749	17%	515	23%	234	11%
Total	4,381		2,255		2,126	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	914	79%	1,194	65%	125	48%	6	50%	2	67%	76	59%
Injection drug use (IDU)	30	3%	71	4%	54	21%	-	-	-	-	4	3%
Male-to-male sexual contact and IDU	67	6%	71	4%	21	8%	-	-	-	-	4	3%
Heterosexual contact	33	3%	163	9%	23	9%	-	-	-	-	12	9%
Other/unknown	116	10%	344	19%	37	14%	6	50%	1	33%	33	26%
Subtotal	1,160	100%	1,843	100%	260	100%	12	100%	3	100%	129	100%
Female adult or adolescent												
Injection drug use	26	18%	68	11%	28	24%	-	-	-	-	2	6%
Heterosexual contact	86	61%	414	66%	72	61%	2	67%	1	100%	18	53%
Other/unknown	29	21%	141	23%	18	15%	1	33%	-	-	14	41%
Subtotal	141	100%	623	100%	118	100%	3	100%	1	100%	34	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	40%	33	83%	9	100%	-	-	-	-	-	-
Other/unknown	3	60%	7	18%	-	-	-	-	-	-	-	-
Subtotal	5	100%	40	100%	9	100%	-	-	-	-	-	-
Total	1,306		2,506		387		15		4		163	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 2

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	895	69%	1106	44%	119	31%	5	33%	2	50%	70	43%
Injection drug use (IDU) only	29	2%	57	2%	41	11%	-	-	-	-	3	2%
Heterosexual contact only	119	9%	577	23%	95	25%	2	13%	1	25%	30	18%
MSM & IDU	60	5%	48	2%	14	4%	-	-	-	-	3	2%
IDU & Heterosexual contact	27	2%	82	3%	41	11%	-	-	-	-	3	2%
MSM & Heterosexual contact	19	1%	88	4%	6	2%	1	7%	-	-	6	4%
MSM & IDU & Heterosexual contact	7	1%	23	1%	7	2%	-	-	-	-	1	1%
Perinatal exposure	2	0%	34	1%	9	2%	-	-	-	-	-	-
Other/unknown	148	11%	491	20%	55	14%	7	47%	1	25%	47	29%
Total	1,306		2,506		387		15		4		163	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 2

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	2,753	77%	2,901	77%	3,067	78%	3,252	78%	3,432	78%
Females	811	23%	848	23%	887	22%	927	22%	949	22%
Age at end of year										
<13	15	<1%	12	<1%	11	<1%	11	<1%	12	<1%
13-14	7	<1%	7	<1%	4	<1%	2	<1%	1	<1%
15-19	41	1%	34	1%	33	1%	36	1%	37	1%
20-24	146	4%	159	4%	182	5%	208	5%	221	5%
25-29	217	6%	230	6%	245	6%	272	7%	296	7%
30-34	254	7%	276	7%	314	8%	332	8%	353	8%
35-39	352	10%	321	9%	313	8%	328	8%	342	8%
40-44	596	17%	587	16%	552	14%	496	12%	488	11%
45-49	723	20%	732	20%	705	18%	726	17%	691	16%
50-54	579	16%	631	17%	716	18%	763	18%	800	18%
55-64	529	15%	632	17%	725	18%	815	20%	911	21%
65+	105	3%	128	3%	154	4%	190	5%	229	5%
Race/Ethnicity										
White, not Hispanic	1,138	32%	1,182	32%	1,220	31%	1,273	30%	1,306	30%
Black/African American, not Hispanic	1,974	55%	2,094	56%	2,219	56%	2,357	56%	2,506	57%
Hispanic/Latino	315	9%	328	9%	355	9%	376	9%	387	9%
Asian/Pacific Islander	12	<1%	12	<1%	13	<1%	13	<1%	15	<1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	121	3%	129	3%	143	4%	156	4%	163	4%
Race/Sex										
White, not Hispanic Males	1,020	29%	1,057	28%	1,090	28%	1,136	27%	1,164	27%
White, not Hispanic Females	118	3%	125	3%	130	3%	137	3%	142	3%
Black/African American, not Hispanic Males	1,423	40%	1,517	40%	1,615	41%	1,726	41%	1,859	42%
Black/African American, not Hispanic Females	551	15%	577	15%	604	15%	631	15%	647	15%
Hispanic/Latino Males	204	6%	214	6%	237	6%	255	6%	265	6%
Hispanic/Latina Females	111	3%	114	3%	118	3%	121	3%	122	3%
Asian/Pacific Islander Males	9	<1%	9	<1%	10	<1%	10	<1%	12	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	121	3%	129	3%	143	4%	156	4%	163	4%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 2

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,848	68%	1,958	68%	2,076	68%	2,198	68%	2,317	68%
Injection drug use (IDU)	150	5%	152	5%	155	5%	158	5%	159	5%
Male-to-male sexual contact and IDU	148	5%	149	5%	158	5%	161	5%	163	5%
Heterosexual contact	201	7%	206	7%	211	7%	220	7%	231	7%
Other/unknown	382	14%	412	14%	443	15%	491	15%	537	16%
Subtotal	2,729	100%	2,877	100%	3,043	100%	3,228	100%	3,407	100%
Female adult or adolescent										
Injection drug use	114	15%	116	14%	118	14%	121	13%	124	13%
Heterosexual contact	521	66%	542	66%	558	65%	579	64%	593	64%
Other/unknown	149	19%	163	20%	184	21%	198	22%	203	22%
Subtotal	784	100%	821	100%	860	100%	898	100%	920	100%
Child (<13 yrs at diagnosis)										
Perinatal	44	86%	44	86%	44	86%	44	83%	44	81%
Other/unknown	7	14%	7	14%	7	14%	9	17%	10	19%
Subtotal	51	100%	51	100%	51	100%	53	100%	54	100%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 2

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	1,737	49%	1,846	49%	1,961	50%	2,079	50%	2,197	50%
Injection drug use (IDU) only	117	3%	119	3%	123	3%	128	3%	130	3%
Heterosexual contact only	722	20%	748	20%	769	19%	799	19%	824	19%
MSM & IDU	113	3%	114	3%	120	3%	123	3%	125	3%
IDU & Heterosexual contact	147	4%	149	4%	150	4%	151	4%	153	3%
MSM & Heterosexual contact	111	3%	112	3%	115	3%	119	3%	120	3%
MSM & IDU & Heterosexual contact	35	1%	35	1%	38	1%	38	1%	38	1%
Perinatal exposure	44	1%	44	1%	44	1%	45	1%	45	1%
Other/unknown	538	15%	582	16%	634	16%	697	17%	749	17%
Total	3,564		3,749		3,954		4,179		4,381	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 3

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 3

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	8.8	57	88%	48	87%	8	89%	1	100%
Females	1.2	8	12%	7	13%	1	11%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	5.7	5	8%	5	9%	-	-	-	-
20-24	15.5	14	22%	14	25%	-	-	-	-
25-29	10.2	8	12%	8	15%	-	-	-	-
30-34	7.9	6	9%	4	7%	2	22%	-	-
35-39	6.8	5	8%	4	7%	1	11%	-	-
40-44	6.1	5	8%	4	7%	1	11%	-	-
45-49	11.1	10	15%	8	15%	1	11%	1	100%
50-54	6.9	7	11%	5	9%	2	22%	-	-
55-64	2.6	5	8%	3	5%	2	22%	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	3.3	37	57%	28	51%	8	89%	1	100%
Black/African American, not Hispanic	20.6	24	37%	23	42%	1	11%	-	-
Hispanic/Latino	*	3	5%	3	5%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	2%	1	2%	-	-	-	-
Race/Sex									
White, not Hispanic Males	6.1	34	52%	26	47%	7	78%	1	100%
White, not Hispanic Females	*	3	5%	2	4%	1	11%	-	-
Black/African American, not Hispanic Males	37.9	21	32%	20	36%	1	11%	-	-
Black/African American, not Hispanic Females	*	3	5%	3	5%	-	-	-	-
Hispanic/Latino Males	*	1	2%	1	2%	-	-	-	-
Hispanic/Latina Females	*	2	3%	2	4%	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	2%	1	2%	-	-	-	-
Total	4.9	65		55		9		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 3

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	44	77%	37	77%	7	88%	-	-
Injection drug use (IDU)	2	4%	2	4%	-	-	-	-
Male-to-male sexual contact and IDU	1	2%	1	2%	-	-	-	-
Heterosexual contact	2	4%	2	4%	-	-	-	-
Other/unknown	8	14%	6	13%	1	13%	1	100%
Subtotal	57	100%	48	100%	8	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	5	71%	5	83%	-	-	-	-
Other/unknown	2	29%	1	17%	1	100%	-	-
Subtotal	7	100%	6	100%	1	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	65		55		9		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 3

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	44	68%	37	67%	7	78%	-	-
Injection drug use (IDU) only	1	2%	1	2%	-	-	-	-
Heterosexual contact only	7	11%	7	13%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	2%	1	2%	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	2%	1	2%	-	-	-	-
Perinatal exposure	1	2%	1	2%	-	-	-	-
Other/unknown	10	15%	7	13%	2	22%	1	100%
Total	65		55		9		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 3

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	26	76%	16	76%	1	100%	-	-	-	-	1	100%
Injection drug use (IDU)	1	3%	1	5%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	3%	1	5%	-	-	-	-	-	-	-	-
Other/unknown	5	15%	3	14%	-	-	-	-	-	-	-	-
Subtotal	34	100%	21	100%	1	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	67%	2	67%	1	100%	-	-	-	-	-	-
Other/unknown	1	33%	1	33%	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-
Total	37		24		3		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 3

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	26	70%	16	67%	1	33%	-	-	-	-	1	100%
Injection drug use (IDU) only	1	3%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	8%	3	13%	1	33%	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	4%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	1	33%	-	-	-	-	-	-
Other/unknown	6	16%	4	17%	-	-	-	-	-	-	-	-
Total	37		24		3		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 3

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	4.9	32	71%	7.7	50	81%	6.1	40	78%	8.9	58	84%
Females	1.9	13	29%	1.8	12	19%	1.6	11	22%	1.6	11	16%
Age at diagnosis (yr)												
<13	*	-	-	*	2	3%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	2%	*	4	6%	6.8	6	12%	*	3	4%
20-24	*	4	9%	13.2	12	19%	7.7	7	14%	14.4	13	19%
25-29	6.4	5	11%	15.3	12	19%	10.2	8	16%	16.6	13	19%
30-34	11.9	9	20%	*	4	6%	13.2	10	20%	6.6	5	7%
35-39	*	1	2%	12.3	9	15%	*	1	2%	9.6	7	10%
40-44	*	2	4%	6.1	5	8%	*	4	8%	9.8	8	12%
45-49	10.0	9	20%	*	2	3%	*	4	8%	*	4	6%
50-54	5.9	6	13%	8.9	9	15%	4.9	5	10%	*	4	6%
55-64	2.6	5	11%	*	3	5%	*	4	8%	4.1	8	12%
65+	*	3	7%	*	-	-	*	2	4%	*	4	6%
Race/Ethnicity												
White, not Hispanic	2.1	24	53%	2.9	33	53%	2.5	28	55%	3.0	34	49%
Black/African American, not Hispanic	12.9	15	33%	21.5	25	40%	17.2	20	39%	24.1	28	41%
Hispanic/Latino	*	2	4%	*	1	2%	*	1	2%	*	3	4%
Asian/Pacific Islander	*	2	4%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	4%	*	3	5%	*	2	4%	*	4	6%
Race/Sex												
White, not Hispanic Males	3.1	17	38%	4.9	27	44%	3.8	21	41%	5.2	29	42%
White, not Hispanic Females	1.2	7	16%	1.0	6	10%	1.2	7	14%	0.9	5	7%
Black/African American, not Hispanic Males	19.9	11	24%	34.3	19	31%	28.9	16	31%	39.7	22	32%
Black/African American, not Hispanic Females	*	4	9%	9.9	6	10%	*	4	8%	9.9	6	9%
Hispanic/Latino Males	*	1	2%	*	1	2%	*	1	2%	*	3	4%
Hispanic/Latina Females	*	1	2%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	2%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	1	2%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	4%	*	3	5%	*	2	4%	*	4	6%
Total	3.4	45		4.7	62		3.8	51		5.2	69	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AID infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 3

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	20	63%	34	71%	29	73%	36	62%
Injection drug use (IDU)	2	6%	1	2%	-	-	3	5%
Male-to-male sexual contact and IDU	2	6%	-	-	-	-	1	2%
Heterosexual contact	1	3%	1	2%	1	3%	3	5%
Other/unknown	7	22%	12	25%	10	25%	15	26%
Subtotal	32	100%	48	100%	40	100%	58	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	9	69%	7	58%	8	73%	9	82%
Other/unknown	4	31%	5	42%	3	27%	2	18%
Subtotal	13	100%	12	100%	11	100%	11	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	2	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	-	-	-	-
Total	45		62		51		69	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 3

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	19	42%	31	50%	28	55%	36	52%
Injection drug use (IDU) only	1	2%	-	-	-	-	3	4%
Heterosexual contact only	10	22%	8	13%	9	18%	12	17%
MSM & IDU	1	2%	-	-	-	-	1	1%
IDU & Heterosexual contact	1	2%	1	2%	-	-	-	-
MSM & Heterosexual contact	1	2%	3	5%	1	2%	-	-
MSM & IDU & Heterosexual contact	1	2%	-	-	-	-	-	-
Perinatal exposure	-	-	2	3%	-	-	-	-
Other/unknown	11	24%	17	27%	13	25%	17	25%
Total	45		62		51		69	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 3

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	16	67%	44	88%	35	83%	46	82%	48	87%
Females	8	33%	6	12%	7	17%	10	18%	7	13%
Age at HIV diagnosis (yr)										
<13	-	-	2	4%	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	4	8%	6	14%	3	5%	5	9%
20-24	3	13%	10	20%	7	17%	11	20%	14	25%
25-29	3	13%	10	20%	6	14%	13	23%	8	15%
30-34	5	21%	4	8%	9	21%	5	9%	4	7%
35-39	1	4%	9	18%	1	2%	6	11%	4	7%
40-44	1	4%	4	8%	3	7%	4	7%	4	7%
45-49	5	21%	1	2%	4	10%	2	4%	8	15%
50-54	2	8%	5	10%	2	5%	3	5%	5	9%
55-64	2	8%	1	2%	2	5%	7	13%	3	5%
65+	2	8%	-	-	2	5%	2	4%	-	-
Race/Ethnicity										
White, not Hispanic	12	50%	26	52%	25	60%	25	45%	28	51%
Black/African American, not Hispanic	10	42%	21	42%	14	33%	25	45%	23	42%
Hispanic/Latino	-	-	1	2%	1	2%	2	4%	3	5%
Asian/Pacific Islander	1	4%	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	2	4%	2	5%	4	7%	1	2%
Race/Sex										
White, not Hispanic Males	9	38%	22	44%	19	45%	20	36%	26	47%
White, not Hispanic Females	3	13%	4	8%	6	14%	5	9%	2	4%
Black/African American, not Hispanic Males	6	25%	19	38%	13	31%	20	36%	20	36%
Black/African American, not Hispanic Females	4	17%	2	4%	1	2%	5	9%	3	5%
Hispanic/Latino Males	-	-	1	2%	1	2%	2	4%	1	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	2	4%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	1	4%	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	2	4%	2	5%	4	7%	1	2%
Total	24		50		42		56		55	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 3

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	11	69%	29	69%	26	74%	33	72%	37	77%
Injection drug use (IDU)	-	-	1	2%	-	-	1	2%	2	4%
Male-to-male sexual contact and IDU	1	6%	-	-	-	-	1	2%	1	2%
Heterosexual contact	-	-	1	2%	1	3%	3	7%	2	4%
Other/unknown	4	25%	11	26%	8	23%	8	17%	6	13%
Subtotal	16	100%	42	100%	35	100%	46	100%	48	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	7	88%	3	50%	6	86%	8	80%	5	83%
Other/unknown	1	13%	3	50%	1	14%	2	20%	1	17%
Subtotal	8	100%	6	100%	7	100%	10	100%	6	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	100%
Subtotal	-	-	2	100%	-	-	-	-	1	100%
Total	24		50		42		56		55	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 3

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	42%	27	54%	25	60%	33	59%	37	67%
Injection drug use (IDU) only	-	-	-	-	-	-	1	2%	1	2%
Heterosexual contact only	7	29%	4	8%	7	17%	11	20%	7	13%
MSM & IDU	-	-	-	-	-	-	1	2%	-	-
IDU & Heterosexual contact	-	-	1	2%	-	-	-	-	1	2%
MSM & Heterosexual contact	1	4%	2	4%	1	2%	-	-	-	-
MSM & IDU & Heterosexual contact	1	4%	-	-	-	-	-	-	1	2%
Perinatal exposure	-	-	2	4%	-	-	-	-	1	2%
Other/unknown	5	21%	14	28%	9	21%	10	18%	7	13%
Total	24		50		42		56		55	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 3

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	15	75%	20	77%	9	69%	17	81%	14	82%	1,082	86%
Females	5	25%	6	23%	4	31%	4	19%	3	18%	175	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	<1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	2	10%	1	4%	-	-	2	10%	-	-	25	2%
20-24	1	5%	5	19%	1	8%	2	10%	1	6%	102	8%
25-29	3	15%	5	19%	2	15%	2	10%	2	12%	190	15%
30-34	2	10%	3	12%	2	15%	1	5%	3	18%	256	20%
35-39	1	5%	-	-	-	-	3	14%	-	-	221	18%
40-44	2	10%	5	19%	1	8%	4	19%	2	12%	191	15%
45-49	3	15%	2	8%	-	-	3	14%	3	18%	122	10%
50-54	3	15%	3	12%	4	31%	-	-	3	18%	74	6%
55-64	2	10%	2	8%	3	23%	3	14%	2	12%	56	4%
65+	1	5%	-	-	-	-	1	5%	1	6%	13	1%
Race/Ethnicity												
White, not Hispanic	13	65%	15	58%	3	23%	12	57%	12	71%	822	65%
Black/African American, not Hispanic	4	20%	9	35%	10	77%	8	38%	5	29%	385	31%
Hispanic/Latino	1	5%	-	-	-	-	1	5%	-	-	32	3%
Asian/Pacific Islander	1	5%	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	1	5%	2	8%	-	-	-	-	-	-	13	1%
Race/Sex												
White, not Hispanic Males	10	50%	12	46%	2	15%	12	57%	10	59%	740	59%
White, not Hispanic Females	3	15%	3	12%	1	8%	-	-	2	12%	82	7%
Black/African American, not Hispanic Males	3	15%	6	23%	7	54%	4	19%	4	24%	305	24%
Black/African American, not Hispanic Females	1	5%	3	12%	3	23%	4	19%	1	6%	80	6%
Hispanic/Latino Males	-	-	-	-	-	-	1	5%	-	-	25	2%
Hispanic/Latina Females	1	5%	-	-	-	-	-	-	-	-	7	1%
Asian/Pacific Islander Males	1	5%	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	1	5%	2	8%	-	-	-	-	-	-	13	1%
Total	20		26		13		21		17		1,257	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 3

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	10	67%	16	80%	4	44%	8	47%	8	57%	759	70%
Injection drug use (IDU)	1	7%	1	5%	1	11%	1	6%	-	-	77	7%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	6%	-	-	66	6%
Heterosexual contact	1	7%	2	10%	1	11%	-	-	1	7%	55	5%
Other/unknown	3	20%	1	5%	3	33%	7	41%	5	36%	122	11%
Subtotal	15	100%	20	100%	9	100%	17	100%	14	100%	1,079	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	23	13%
Heterosexual contact	4	80%	5	83%	2	50%	3	75%	2	67%	125	73%
Other/unknown	1	20%	1	17%	2	50%	1	25%	1	33%	24	14%
Subtotal	5	100%	6	100%	4	100%	4	100%	3	100%	172	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	20		26		13		21		17		1,257	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 3

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	10	50%	12	46%	4	31%	8	38%	8	47%	731	58%
Injection drug use (IDU) only	1	5%	-	-	-	-	1	5%	-	-	64	5%
Heterosexual contact only	5	25%	7	27%	3	23%	3	14%	3	18%	180	14%
MSM & IDU	-	-	-	-	-	-	1	5%	-	-	61	5%
IDU & Heterosexual contact	-	-	1	4%	1	8%	-	-	-	-	36	3%
MSM & Heterosexual contact	-	-	4	15%	-	-	-	-	-	-	27	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	5	<1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	<1%
Other/unknown	4	20%	2	8%	5	38%	8	38%	6	35%	148	12%
Total	20		26		13		21		17		1,257	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 3

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 3

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	172.8	1,124	80%	581	79%	543	82%
Females	40.5	276	20%	159	21%	117	18%
Age at end of year							
<13	*	4	<1%	4	1%	-	-
13-14	*	2	<1%	2	<1%	-	-
15-19	13.6	12	1%	12	2%	-	-
20-24	54.1	49	4%	43	6%	6	1%
25-29	128.8	101	7%	77	10%	24	4%
30-34	117.6	89	6%	70	9%	19	3%
35-39	162.9	119	9%	80	11%	39	6%
40-44	169.0	138	10%	66	9%	72	11%
45-49	255.9	230	16%	104	14%	126	19%
50-54	256.6	260	19%	112	15%	148	22%
55-64	157.4	306	22%	132	18%	174	26%
65+	39.7	90	6%	38	5%	52	8%
Race/Ethnicity							
White, not Hispanic	68.8	781	56%	398	54%	383	58%
Black/African American, not Hispanic	437.7	509	36%	282	38%	227	34%
Hispanic/Latino	218.8	67	5%	30	4%	37	6%
Asian/Pacific Islander	38.4	9	1%	6	1%	3	<1%
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	32	2%	23	3%	9	1%
Race/Sex							
White, not Hispanic Males	121.4	674	48%	335	45%	339	51%
White, not Hispanic Females	18.4	107	8%	63	9%	44	7%
Black/African American, not Hispanic Males	660.5	366	26%	200	27%	166	25%
Black/African American, not Hispanic Females	234.9	143	10%	82	11%	61	9%
Hispanic/Latino Males	333.9	53	4%	23	3%	30	5%
Hispanic/Latina Females	94.9	14	1%	7	1%	7	1%
Asian/Pacific Islander Males	53.4	6	<1%	5	1%	1	<1%
Asian/Pacific Islander Females	*	3	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	32	2%	23	3%	9	1%
Total	105.1	1,400		740		660	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 3

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	755	68%	375	66%	380	70%
Injection drug use (IDU)	54	5%	15	3%	39	7%
Male-to-male sexual contact and IDU	53	5%	19	3%	34	6%
Heterosexual contact	63	6%	31	5%	32	6%
Other/unknown	186	17%	131	23%	55	10%
Subtotal	1,111	100%	571	100%	540	100%
Female adult or adolescent						
Injection drug use	23	9%	12	8%	11	9%
Heterosexual contact	178	66%	90	59%	88	76%
Other/unknown	67	25%	50	33%	17	15%
Subtotal	268	100%	152	100%	116	100%
Child (<13 yrs at diagnosis)						
Perinatal	18	86%	14	82%	4	100%
Other/unknown	3	14%	3	18%	-	-
Subtotal	21	100%	17	100%	4	100%
Total	1,400		740		660	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 3

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	723	52%	364	49%	359	54%
Injection drug use (IDU) only	48	3%	16	2%	32	5%
Heterosexual contact only	241	17%	121	16%	120	18%
MSM & IDU	41	3%	12	2%	29	4%
IDU & Heterosexual contact	29	2%	11	1%	18	3%
MSM & Heterosexual contact	32	2%	11	1%	21	3%
MSM & IDU & Heterosexual contact	12	1%	7	1%	5	1%
Perinatal exposure	19	1%	15	2%	4	1%
Other/unknown	255	18%	183	25%	72	11%
Total	1,400		740		660	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 3

Living with a Diagnosis of HIV Infection in 2014												
Transmission Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	488	73%	222	62%	28	57%	4	67%	-	-	13	54%
Injection drug use (IDU)	25	4%	22	6%	5	10%	-	-	1	100%	1	4%
Male-to-male sexual contact and IDU	34	5%	17	5%	2	4%	-	-	-	-	-	-
Heterosexual contact	31	5%	20	6%	8	16%	1	17%	-	-	3	13%
Other/unknown	95	14%	77	22%	6	12%	1	17%	-	-	7	29%
Subtotal	673	100%	358	100%	49	100%	6	100%	1	100%	24	100%
Female adult or adolescent												
Injection drug use	5	5%	15	11%	2	15%	-	-	-	-	1	14%
Heterosexual contact	77	73%	87	63%	10	77%	2	67%	-	-	2	29%
Other/unknown	23	22%	37	27%	1	8%	1	33%	1	100%	4	57%
Subtotal	105	100%	139	100%	13	100%	3	100%	1	100%	7	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	67%	11	92%	4	80%	-	-	-	-	1	100%
Other/unknown	1	33%	1	8%	1	20%	-	-	-	-	-	-
Subtotal	3	100%	12	100%	5	100%	-	-	-	-	1	100%
Total	781		509		67		9		2		32	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 3

Living with a Diagnosis of HIV Infection in 2014												
Exposure Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	474	61%	209	41%	26	39%	3	33%	-	-	11	34%
Injection drug use (IDU) only	22	3%	21	4%	4	6%	-	-	1	50%	-	-
Heterosexual contact only	108	14%	107	21%	18	27%	3	33%	-	-	5	16%
MSM & IDU	26	3%	13	3%	2	3%	-	-	-	-	-	-
IDU & Heterosexual contact	8	1%	16	3%	3	4%	-	-	-	-	2	6%
MSM & Heterosexual contact	14	2%	13	3%	2	3%	1	11%	-	-	2	6%
MSM & IDU & Heterosexual contact	8	1%	4	1%	-	-	-	-	-	-	-	-
Perinatal exposure	3	0%	11	2%	4	6%	-	-	-	-	1	3%
Other/unknown	118	15%	115	23%	8	12%	2	22%	1	50%	11	34%
Total	781		509		67		9		2		32	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 3

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	925	80%	972	79%	1,011	79%	1,066	80%	1,124	80%
Females	238	20%	252	21%	262	21%	271	20%	276	20%
Age at end of year										
<13	3	<1%	5	<1%	6	<1%	6	<1%	4	<1%
13-14	3	<1%	2	<1%	-	-	-	-	2	<1%
15-19	10	1%	9	1%	12	1%	9	1%	12	1%
20-24	46	4%	48	4%	40	3%	45	3%	49	4%
25-29	65	6%	79	6%	89	7%	100	7%	101	7%
30-34	90	8%	85	7%	87	7%	89	7%	89	6%
35-39	109	9%	117	10%	118	9%	114	9%	119	9%
40-44	188	16%	172	14%	156	12%	148	11%	138	10%
45-49	241	21%	242	20%	243	19%	237	18%	230	16%
50-54	181	16%	201	16%	229	18%	246	18%	260	19%
55-64	192	17%	220	18%	238	19%	268	20%	306	22%
65+	35	3%	44	4%	55	4%	75	6%	90	6%
Race/Ethnicity										
White, not Hispanic	658	57%	690	56%	717	56%	748	56%	781	56%
Black/African American, not Hispanic	415	36%	439	36%	457	36%	483	36%	509	36%
Hispanic/Latino	57	5%	58	5%	61	5%	64	5%	67	5%
Asian/Pacific Islander	8	1%	9	1%	9	1%	9	1%	9	1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	23	2%	26	2%	27	2%	31	2%	32	2%
Race/Sex										
White, not Hispanic Males	572	49%	597	49%	617	48%	643	48%	674	48%
White, not Hispanic Females	86	7%	93	8%	100	8%	105	8%	107	8%
Black/African American, not Hispanic Males	287	25%	305	25%	320	25%	342	26%	366	26%
Black/African American, not Hispanic Females	128	11%	134	11%	137	11%	141	11%	143	10%
Hispanic/Latino Males	44	4%	45	4%	48	4%	51	4%	53	4%
Hispanic/Latina Females	13	1%	13	1%	13	1%	13	1%	14	1%
Asian/Pacific Islander Males	6	1%	6	<1%	6	<1%	6	<1%	6	<1%
Asian/Pacific Islander Females	2	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	23	2%	26	2%	27	2%	31	2%	32	2%
Total	1,163		1,224		1,273		1,337		1,400	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 3

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	617	67%	650	68%	678	68%	710	67%	755	68%
Injection drug use (IDU)	48	5%	49	5%	49	5%	52	5%	54	5%
Male-to-male sexual contact and IDU	51	6%	52	5%	52	5%	52	5%	53	5%
Heterosexual contact	54	6%	55	6%	57	6%	60	6%	63	6%
Other/unknown	145	16%	154	16%	162	16%	179	17%	186	17%
Subtotal	915	100%	960	100%	998	100%	1,053	100%	1,111	100%
Female adult or adolescent										
Injection drug use	22	10%	23	9%	23	9%	23	9%	23	9%
Heterosexual contact	154	67%	161	66%	168	66%	175	67%	178	66%
Other/unknown	54	23%	60	25%	63	25%	65	25%	67	25%
Subtotal	230	100%	244	100%	254	100%	263	100%	268	100%
Child (<13 yrs at diagnosis)										
Perinatal	15	83%	17	85%	18	86%	18	86%	18	86%
Other/unknown	3	17%	3	15%	3	14%	3	14%	3	14%
Subtotal	18	100%	20	100%	21	100%	21	100%	21	100%
Total	1,163		1,224		1,273		1,337		1,400	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 3

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	587	50%	618	50%	646	51%	678	51%	723	52%
Injection drug use (IDU) only	44	4%	45	4%	45	4%	47	4%	48	3%
Heterosexual contact only	208	18%	216	18%	225	18%	235	18%	241	17%
MSM & IDU	40	3%	41	3%	41	3%	41	3%	41	3%
IDU & Heterosexual contact	26	2%	27	2%	27	2%	28	2%	29	2%
MSM & Heterosexual contact	30	3%	32	3%	32	3%	32	2%	32	2%
MSM & IDU & Heterosexual contact	11	1%	11	1%	11	1%	11	1%	12	1%
Perinatal exposure	16	1%	18	1%	19	1%	19	1%	19	1%
Other/unknown	201	17%	216	18%	227	18%	246	18%	255	18%
Total	1,163		1,224		1,273		1,337		1,400	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 4

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 4

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	9.1	41	85%	23	82%	11	85%	7	100%
Females	1.5	7	15%	5	18%	2	15%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	3	6%	2	7%	-	-	1	14%
20-24	12.2	7	15%	4	14%	2	15%	1	14%
25-29	25.0	13	27%	9	32%	3	23%	1	14%
30-34	*	3	6%	2	7%	1	8%	-	-
35-39	*	1	2%	1	4%	-	-	-	-
40-44	8.9	5	10%	3	11%	2	15%	-	-
45-49	17.1	10	21%	5	18%	3	23%	2	29%
50-54	*	-	-	-	-	-	-	-	-
55-64	4.4	6	13%	2	7%	2	15%	2	29%
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	2.6	21	44%	14	50%	3	23%	4	57%
Black/African American, not Hispanic	26.6	19	40%	9	32%	7	54%	3	43%
Hispanic/Latino	20.1	5	10%	2	7%	3	23%	-	-
Asian/Pacific Islander	*	1	2%	1	4%	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	2	4%	2	7%	-	-	-	-
Race/Sex									
White, not Hispanic Males	4.9	19	40%	12	43%	3	23%	4	57%
White, not Hispanic Females	*	2	4%	2	7%	-	-	-	-
Black/African American, not Hispanic Males	45.5	16	33%	8	29%	5	38%	3	43%
Black/African American, not Hispanic Females	*	3	6%	1	4%	2	15%	-	-
Hispanic/Latino Males	36.4	5	10%	2	7%	3	23%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	2%	1	4%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	2	4%	2	7%	-	-	-	-
Total	5.2	48		28		13		7	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	28	68%	15	65%	8	73%	5	71%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	3	7%	2	9%	-	-	1	14%
Other/unknown	10	24%	6	26%	3	27%	1	14%
Subtotal	41	100%	23	100%	11	100%	7	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	7	100%	5	100%	2	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	7	100%	5	100%	2	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	48		28		13		7	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	28	58%	15	54%	8	62%	5	71%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	10	21%	7	25%	2	15%	1	14%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	10	21%	6	21%	3	23%	1	14%
Total	48		28		13		7	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	14	74%	12	75%	2	40%	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	11%	1	6%	-	-	-	-	-	-	-	-
Other/unknown	3	16%	3	19%	3	60%	-	-	-	-	1	100%
Subtotal	19	100%	16	100%	5	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	3	100%	-	-	1	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	3	100%	-	-	1	100%	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	21		19		5		1		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	67%	12	63%	2	40%	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	4	19%	4	21%	-	-	1	100%	-	-	1	50%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	14%	3	16%	3	60%	-	-	-	-	1	50%
Total	21		19		5		1		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 4

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	10.9	49	89%	10.7	48	79%	9.6	43	88%	6.4	29	74%
Females	1.3	6	11%	2.8	13	21%	1.3	6	12%	2.1	10	26%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	1	2%	*	1	3%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	12.0	7	13%	12.0	7	11%	*	3	6%	*	-	-
20-24	12.2	7	13%	24.4	14	23%	10.5	6	12%	10.5	6	15%
25-29	*	4	7%	23.0	12	20%	13.4	7	14%	11.5	6	15%
30-34	*	4	7%	*	4	7%	9.7	5	10%	*	4	10%
35-39	15.8	8	15%	*	4	7%	*	4	8%	*	4	10%
40-44	19.6	11	20%	*	3	5%	8.9	5	10%	*	2	5%
45-49	*	4	7%	*	3	5%	13.7	8	16%	11.9	7	18%
50-54	10.2	7	13%	8.8	6	10%	8.8	6	12%	*	2	5%
55-64	*	3	5%	4.4	6	10%	*	3	6%	4.4	6	15%
65+	*	-	-	*	2	3%	*	1	2%	*	1	3%
Race/Ethnicity												
White, not Hispanic	3.8	30	55%	3.6	29	48%	3.6	29	59%	3.4	27	69%
Black/African American, not Hispanic	26.6	19	35%	36.4	26	43%	19.6	14	29%	11.2	8	21%
Hispanic/Latino	*	2	4%	*	2	3%	*	3	6%	*	2	5%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	1	2%	*	1	2%	*	-	-	*	-	-
Unknown	*	3	5%	*	3	5%	*	3	6%	*	2	5%
Race/Sex												
White, not Hispanic Males	6.7	26	47%	5.7	22	36%	6.7	26	53%	5.4	21	54%
White, not Hispanic Females	*	4	7%	1.7	7	11%	*	3	6%	1.5	6	15%
Black/African American, not Hispanic Males	48.4	17	31%	62.6	22	36%	34.2	12	24%	14.2	5	13%
Black/African American, not Hispanic Females	*	2	4%	*	4	7%	*	2	4%	*	3	8%
Hispanic/Latino Males	*	2	4%	*	1	2%	*	3	6%	*	2	5%
Hispanic/Latina Females	*	-	-	*	1	2%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	1	2%	*	1	2%	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	5%	*	3	5%	*	3	6%	*	2	5%
Total	6.0	55		6.6	61		5.3	49		4.2	39	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AID infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 4

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	32	65%	33	69%	27	64%	19	66%
Injection drug use (IDU)	2	4%	1	2%	1	2%	1	3%
Male-to-male sexual contact and IDU	2	4%	-	-	1	2%	1	3%
Heterosexual contact	7	14%	3	6%	1	2%	3	10%
Other/unknown	6	12%	11	23%	12	29%	5	17%
Subtotal	49	100%	48	100%	42	100%	29	100%
Female adult or adolescent								
Injection drug use	1	17%	3	23%	1	17%	1	11%
Heterosexual contact	5	83%	7	54%	3	50%	7	78%
Other/unknown	-	-	3	23%	2	33%	1	11%
Subtotal	6	100%	13	100%	6	100%	9	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	1	100%
Subtotal	-	-	-	-	1	100%	1	100%
Total	55		61		49		39	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 4

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	29	53%	31	51%	25	51%	19	49%
Injection drug use (IDU) only	3	5%	2	3%	2	4%	1	3%
Heterosexual contact only	12	22%	10	16%	4	8%	10	26%
MSM & IDU	1	2%	-	-	-	-	1	3%
IDU & Heterosexual contact	-	-	2	3%	-	-	1	3%
MSM & Heterosexual contact	3	5%	2	3%	2	4%	-	-
MSM & IDU & Heterosexual contact	1	2%	-	-	1	2%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	6	11%	14	23%	15	31%	7	18%
Total	55		61		49		39	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 4

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	30	91%	31	79%	28	90%	17	71%	23	82%
Females	3	9%	8	21%	3	10%	7	29%	5	18%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	1	3%	1	4%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	5	15%	7	18%	3	10%	-	-	2	7%
20-24	5	15%	11	28%	3	10%	5	21%	4	14%
25-29	3	9%	8	21%	5	16%	6	25%	9	32%
30-34	3	9%	3	8%	2	6%	3	13%	2	7%
35-39	6	18%	3	8%	2	6%	1	4%	1	4%
40-44	4	12%	1	3%	5	16%	-	-	3	11%
45-49	1	3%	-	-	5	16%	4	17%	5	18%
50-54	5	15%	3	8%	3	10%	1	4%	-	-
55-64	1	3%	2	5%	1	3%	3	13%	2	7%
65+	-	-	1	3%	1	3%	-	-	-	-
Race/Ethnicity										
White, not Hispanic	19	58%	21	54%	19	61%	12	50%	14	50%
Black/African American, not Hispanic	10	30%	14	36%	8	26%	8	33%	9	32%
Hispanic/Latino	2	6%	2	5%	2	6%	2	8%	2	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	4%
American Indian/Alaska Native	-	-	1	3%	-	-	-	-	-	-
Unknown	2	6%	1	3%	2	6%	2	8%	2	7%
Race/Sex										
White, not Hispanic Males	16	48%	16	41%	17	55%	9	38%	12	43%
White, not Hispanic Females	3	9%	5	13%	2	6%	3	13%	2	7%
Black/African American, not Hispanic Males	10	30%	12	31%	8	26%	5	21%	8	29%
Black/African American, not Hispanic Females	-	-	2	5%	-	-	3	13%	1	4%
Hispanic/Latino Males	2	6%	1	3%	2	6%	2	8%	2	7%
Hispanic/Latina Females	-	-	1	3%	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	4%
American Indian/Alaska Native Males	-	-	1	3%	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	6%	1	3%	2	6%	2	8%	2	7%
Total	33		39		31		24		28	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	20	67%	24	77%	19	70%	11	65%	15	65%
Injection drug use (IDU)	1	3%	-	-	1	4%	1	6%	-	-
Male-to-male sexual contact and IDU	2	7%	-	-	-	-	1	6%	-	-
Heterosexual contact	4	13%	1	3%	1	4%	2	12%	2	9%
Other/unknown	3	10%	6	19%	6	22%	2	12%	6	26%
Subtotal	30	100%	31	100%	27	100%	17	100%	23	100%
Female adult or adolescent										
Injection drug use	1	33%	-	-	-	-	1	17%	-	-
Heterosexual contact	2	67%	6	75%	1	33%	4	67%	5	100%
Other/unknown	-	-	2	25%	2	67%	1	17%	-	-
Subtotal	3	100%	8	100%	3	100%	6	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	1	100%	-	-
Subtotal	-	-	-	-	1	100%	1	100%	-	-
Total	33		39		31		24		28	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	18	55%	23	59%	17	55%	11	46%	15	54%
Injection drug use (IDU) only	2	6%	-	-	1	3%	1	4%	-	-
Heterosexual contact only	6	18%	7	18%	2	6%	6	25%	7	25%
MSM & IDU	1	3%	-	-	-	-	1	4%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	1	4%	-	-
MSM & Heterosexual contact	2	6%	1	3%	2	6%	-	-	-	-
MSM & IDU & Heterosexual contact	1	3%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	9%	8	21%	9	29%	4	17%	6	21%
Total	33		39		31		24		28	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 4

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	18	67%	27	77%	16	76%	17	81%	22	92%	746	79%
Females	9	33%	8	23%	5	24%	4	19%	2	8%	201	21%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	10	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	3	11%	1	3%	-	-	-	-	1	4%	22	2%
20-24	1	4%	5	14%	3	14%	3	14%	4	17%	78	8%
25-29	3	11%	6	17%	2	10%	1	5%	3	13%	148	16%
30-34	3	11%	4	11%	4	19%	3	14%	1	4%	178	19%
35-39	3	11%	1	3%	4	19%	2	10%	1	4%	177	19%
40-44	7	26%	5	14%	-	-	2	10%	4	17%	128	14%
45-49	4	15%	5	14%	3	14%	4	19%	5	21%	94	10%
50-54	1	4%	3	9%	3	14%	2	10%	1	4%	45	5%
55-64	2	7%	4	11%	2	10%	3	14%	4	17%	54	6%
65+	-	-	1	3%	-	-	1	5%	-	-	11	1%
Race/Ethnicity												
White, not Hispanic	10	37%	14	40%	11	52%	17	81%	8	33%	581	61%
Black/African American, not Hispanic	14	52%	18	51%	8	38%	4	19%	11	46%	317	33%
Hispanic/Latino	1	4%	1	3%	1	5%	-	-	3	13%	36	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	4%	1	<1%
Unknown	2	7%	2	6%	1	5%	-	-	1	4%	12	1%
Race/Sex												
White, not Hispanic Males	9	33%	11	31%	10	48%	13	62%	8	33%	501	53%
White, not Hispanic Females	1	4%	3	9%	1	5%	4	19%	-	-	80	8%
Black/African American, not Hispanic Males	6	22%	15	43%	4	19%	4	19%	9	38%	208	22%
Black/African American, not Hispanic Females	8	30%	3	9%	4	19%	-	-	2	8%	109	12%
Hispanic/Latino Males	1	4%	-	-	1	5%	-	-	3	13%	26	3%
Hispanic/Latina Females	-	-	1	3%	-	-	-	-	-	-	10	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	1	4%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	7%	2	6%	1	5%	-	-	1	4%	12	1%
Total	27		35		21		21		24		947	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	11	61%	14	52%	9	56%	13	76%	14	64%	477	64%
Injection drug use (IDU)	2	11%	1	4%	-	-	-	-	1	5%	43	6%
Male-to-male sexual contact and IDU	-	-	-	-	1	6%	-	-	-	-	36	5%
Heterosexual contact	1	6%	5	19%	-	-	2	12%	2	9%	58	8%
Other/unknown	4	22%	7	26%	6	38%	2	12%	5	23%	127	17%
Subtotal	18	100%	27	100%	16	100%	17	100%	22	100%	741	100%
Female adult or adolescent												
Injection drug use	1	11%	4	50%	1	20%	-	-	-	-	44	22%
Heterosexual contact	8	89%	3	38%	3	60%	4	100%	2	100%	121	62%
Other/unknown	-	-	1	13%	1	20%	-	-	-	-	31	16%
Subtotal	9	100%	8	100%	5	100%	4	100%	2	100%	196	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	8	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	20%
Subtotal	-	-	-	-	-	-	-	-	-	-	10	100%
Total	27		35		21		21		24		947	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	11	41%	10	29%	9	43%	13	62%	14	58%	455	48%
Injection drug use (IDU) only	2	7%	3	9%	1	5%	-	-	1	4%	46	5%
Heterosexual contact only	9	33%	8	23%	3	14%	6	29%	4	17%	179	19%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	24	3%
IDU & Heterosexual contact	1	4%	2	6%	-	-	-	-	-	-	40	4%
MSM & Heterosexual contact	-	-	4	11%	-	-	-	-	-	-	22	2%
MSM & IDU & Heterosexual contact	-	-	-	-	1	5%	-	-	-	-	12	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	8	1%
Other/unknown	4	15%	8	23%	7	33%	2	10%	5	21%	161	17%
Total	27		35		21		21		24		947	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 4

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 4

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	174.6	785	75%	406	75%	379	74%
Females	56.7	266	25%	135	25%	131	26%
Age at end of year							
<13	*	4	<1%	4	1%	-	-
13-14	*	1	<1%	1	<1%	-	-
15-19	12.0	7	1%	6	1%	1	<1%
20-24	80.2	46	4%	37	7%	9	2%
25-29	153.6	80	8%	59	11%	21	4%
30-34	122.2	63	6%	41	8%	22	4%
35-39	128.2	65	6%	39	7%	26	5%
40-44	228.2	128	12%	59	11%	69	14%
45-49	290.1	170	16%	84	16%	86	17%
50-54	294.0	201	19%	73	13%	128	25%
55-64	156.5	215	20%	104	19%	111	22%
65+	42.1	71	7%	34	6%	37	7%
Race/Ethnicity							
White, not Hispanic	71.2	568	54%	295	55%	273	54%
Black/African American, not Hispanic	489.0	349	33%	166	31%	183	36%
Hispanic/Latino	348.9	87	8%	44	8%	43	8%
Asian/Pacific Islander	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native	*	3	<1%	2	<1%	1	<1%
Unknown	*	43	4%	33	6%	10	2%
Race/Sex							
White, not Hispanic Males	118.1	459	44%	235	43%	224	44%
White, not Hispanic Females	26.7	109	10%	60	11%	49	10%
Black/African American, not Hispanic Males	654.6	230	22%	118	22%	112	22%
Black/African American, not Hispanic Females	328.3	119	11%	48	9%	71	14%
Hispanic/Latino Males	443.6	61	6%	28	5%	33	6%
Hispanic/Latina Females	232.5	26	2%	16	3%	10	2%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	43	4%	33	6%	10	2%
Total	114.4	1,051		541		510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	495	63%	244	61%	251	66%
Injection drug use (IDU)	50	6%	18	4%	32	8%
Male-to-male sexual contact and IDU	28	4%	13	3%	15	4%
Heterosexual contact	57	7%	29	7%	28	7%
Other/unknown	151	19%	98	24%	53	14%
Subtotal	781	100%	402	100%	379	100%
Female adult or adolescent						
Injection drug use	29	12%	12	10%	17	13%
Heterosexual contact	161	64%	71	57%	90	70%
Other/unknown	62	25%	41	33%	21	16%
Subtotal	252	100%	124	100%	128	100%
Child (<13 yrs at diagnosis)						
Perinatal	14	78%	11	73%	3	100%
Other/unknown	4	22%	4	27%	-	-
Subtotal	18	100%	15	100%	3	100%
Total	1,051		541		510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	461	44%	230	43%	231	45%
Injection drug use (IDU) only	51	5%	21	4%	30	6%
Heterosexual contact only	218	21%	100	18%	118	23%
MSM & IDU	22	2%	12	2%	10	2%
IDU & Heterosexual contact	28	3%	9	2%	19	4%
MSM & Heterosexual contact	34	3%	14	3%	20	4%
MSM & IDU & Heterosexual contact	6	1%	1	<1%	5	1%
Perinatal exposure	15	1%	12	2%	3	1%
Other/unknown	216	21%	142	26%	74	15%
Total	1,051		541		510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 4

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	328	72%	131	57%	24	39%	-	-	1	33%	11	34%
Injection drug use (IDU)	14	3%	21	9%	14	23%	-	-	-	-	1	3%
Male-to-male sexual contact and IDU	19	4%	7	3%	1	2%	-	-	-	-	1	3%
Heterosexual contact	20	4%	24	10%	6	10%	-	-	-	-	7	22%
Other/unknown	75	16%	46	20%	16	26%	-	-	2	67%	12	38%
Subtotal	456	100%	229	100%	61	100%	-	-	3	100%	32	100%
Female adult or adolescent												
Injection drug use	16	15%	12	10%	1	5%	-	-	-	-	-	-
Heterosexual contact	68	65%	75	65%	15	68%	1	100%	-	-	2	22%
Other/unknown	21	20%	28	24%	6	27%	-	-	-	-	7	78%
Subtotal	105	100%	115	100%	22	100%	1	100%	-	-	9	100%
Child (<13 yrs at diagnosis)												
Perinatal	7	100%	4	80%	3	75%	-	-	-	-	-	-
Other/unknown	-	-	1	20%	1	25%	-	-	-	-	2	100%
Subtotal	7	100%	5	100%	4	100%	-	-	-	-	2	100%
Total	568		349		87		1		3		43	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 4

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	311	55%	119	34%	20	23%	-	-	1	33%	10	19%
Injection drug use (IDU) only	22	4%	20	6%	8	9%	-	-	-	-	1	2%
Heterosexual contact only	88	15%	99	28%	21	24%	1	100%	-	-	9	17%
MSM & IDU	15	3%	5	1%	1	1%	-	-	-	-	1	2%
IDU & Heterosexual contact	8	1%	13	4%	7	8%	-	-	-	-	-	-
MSM & Heterosexual contact	17	3%	12	3%	4	5%	-	-	-	-	1	2%
MSM & IDU & Heterosexual contact	4	1%	2	1%	-	-	-	-	-	-	-	-
Perinatal exposure	7	1%	4	1%	4	5%	-	-	-	-	-	-
Other/unknown	96	17%	75	21%	22	25%	-	-	2	67%	31	58%
Total	568		349		87		1		3		53	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 4

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	619	73%	666	74%	715	74%	746	74%	785	75%
Females	227	27%	238	26%	248	26%	257	26%	266	25%
Age at end of year										
<13	2	<1%	2	<1%	4	<1%	5	<1%	4	<1%
13-14	1	<1%	-	-	1	<1%	1	<1%	1	<1%
15-19	12	1%	13	1%	8	1%	4	<1%	7	1%
20-24	40	5%	51	6%	50	5%	50	5%	46	4%
25-29	33	4%	48	5%	67	7%	72	7%	80	8%
30-34	48	6%	48	5%	51	5%	54	5%	63	6%
35-39	90	11%	78	9%	75	8%	74	7%	65	6%
40-44	144	17%	145	16%	140	15%	133	13%	128	12%
45-49	181	21%	171	19%	161	17%	171	17%	170	16%
50-54	141	17%	161	18%	194	20%	189	19%	201	19%
55-64	128	15%	149	16%	167	17%	190	19%	215	20%
65+	26	3%	38	4%	45	5%	60	6%	71	7%
Race/Ethnicity										
White, not Hispanic	454	54%	485	54%	519	54%	546	54%	568	54%
Black/African American, not Hispanic	284	34%	305	34%	322	33%	332	33%	349	33%
Hispanic/Latino	73	9%	76	8%	80	8%	81	8%	87	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	1	<1%	2	<1%	3	<1%	3	<1%	3	<1%
Unknown	34	4%	36	4%	39	4%	41	4%	43	4%
Race/Sex										
White, not Hispanic Males	364	43%	389	43%	418	43%	440	44%	459	44%
White, not Hispanic Females	90	11%	96	11%	101	10%	106	11%	109	10%
Black/African American, not Hispanic Males	178	21%	196	22%	209	22%	216	22%	230	22%
Black/African American, not Hispanic Females	106	13%	109	12%	113	12%	116	12%	119	11%
Hispanic/Latino Males	49	6%	51	6%	55	6%	56	6%	61	6%
Hispanic/Latina Females	24	3%	25	3%	25	3%	25	2%	26	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	1	<1%	2	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	34	4%	36	4%	39	4%	41	4%	43	4%
Total	846		904		963		1,003		1,051	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 4

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	380	62%	412	62%	444	62%	468	63%	495	63%
Injection drug use (IDU)	47	8%	47	7%	49	7%	50	7%	50	6%
Male-to-male sexual contact and IDU	24	4%	24	4%	26	4%	27	4%	28	4%
Heterosexual contact	49	8%	52	8%	53	7%	55	7%	57	7%
Other/unknown	117	19%	129	19%	139	20%	142	19%	151	19%
Subtotal	617	100%	664	100%	711	100%	742	100%	781	100%
Female adult or adolescent										
Injection drug use	25	12%	27	12%	28	12%	29	12%	29	12%
Heterosexual contact	133	62%	140	62%	146	62%	153	63%	161	64%
Other/unknown	58	27%	60	26%	62	26%	62	25%	62	25%
Subtotal	216	100%	227	100%	236	100%	244	100%	252	100%
Child (<13 yrs at diagnosis)										
Perinatal	12	92%	12	92%	14	88%	14	82%	14	78%
Other/unknown	1	8%	1	8%	2	13%	3	18%	4	22%
Subtotal	13	100%	13	100%	16	100%	17	100%	18	100%
Total	846		904		963		1,003		1,051	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 4

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	348	41%	379	42%	410	43%	434	43%	461	44%
Injection drug use (IDU) only	46	5%	47	5%	50	5%	51	5%	51	5%
Heterosexual contact only	182	22%	192	21%	199	21%	208	21%	218	21%
MSM & IDU	19	2%	19	2%	20	2%	21	2%	22	2%
IDU & Heterosexual contact	26	3%	27	3%	27	3%	28	3%	28	3%
MSM & Heterosexual contact	32	4%	33	4%	34	4%	34	3%	34	3%
MSM & IDU & Heterosexual contact	5	1%	5	1%	6	1%	6	1%	6	1%
Perinatal exposure	12	1%	12	1%	14	1%	14	1%	15	1%
Other/unknown	176	21%	190	21%	203	21%	207	21%	216	21%
Total	846		904		963		1,003		1,051	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 5

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 5

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	31.5	189	88%	128	88%	44	90%	17	85%	
Females	4.1	26	12%	18	12%	5	10%	3	15%	
Age at diagnosis (yr)										
<13	*	1	<1%	1	1%	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	7.6	6	3%	6	4%	-	-	-	-	
20-24	50.7	47	22%	33	23%	10	20%	4	20%	
25-29	38.8	46	21%	35	24%	7	14%	4	20%	
30-34	37.2	37	17%	22	15%	7	14%	8	40%	
35-39	29.7	25	12%	20	14%	5	10%	-	-	
40-44	21.2	17	8%	10	7%	7	14%	-	-	
45-49	27.4	21	10%	12	8%	7	14%	2	10%	
50-54	7.4	6	3%	2	1%	3	6%	1	5%	
55-64	5.0	7	3%	4	3%	2	4%	1	5%	
65+	*	2	1%	1	1%	1	2%	-	-	
Race/Ethnicity										
White, not Hispanic	10.5	85	40%	58	40%	23	47%	4	20%	
Black/African American, not Hispanic	38.6	103	48%	70	48%	20	41%	13	65%	
Hispanic/Latino	25.4	16	7%	11	8%	3	6%	2	10%	
Asian/Pacific Islander	*	4	2%	2	1%	1	2%	1	5%	
American Indian/Alaska Native	*	1	<1%	1	1%	-	-	-	-	
Unknown	*	6	3%	4	3%	2	4%	-	-	
Race/Sex										
White, not Hispanic Males	20.4	81	38%	56	38%	22	45%	3	15%	
White, not Hispanic Females	*	4	2%	2	1%	1	2%	1	5%	
Black/African American, not Hispanic Males	68.3	86	40%	59	40%	16	33%	11	55%	
Black/African American, not Hispanic Females	12.0	17	8%	11	8%	4	8%	2	10%	
Hispanic/Latino Males	42.5	14	7%	9	6%	3	6%	2	10%	
Hispanic/Latina Females	*	2	1%	2	1%	-	-	-	-	
Asian/Pacific Islander Males	*	3	1%	1	1%	1	2%	1	5%	
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	1	<1%	1	1%	-	-	-	-	
Unknown	*	6	3%	4	3%	2	4%	-	-	
Total	17.5	215		146		49		20		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	144	76%	101	79%	33	75%	10	59%
Injection drug use (IDU)	1	1%	1	1%	-	-	-	-
Male-to-male sexual contact and IDU	6	3%	6	5%	-	-	-	-
Heterosexual contact	5	3%	-	-	2	5%	3	18%
Other/unknown	33	17%	20	16%	9	20%	4	24%
Subtotal	189	100%	128	100%	44	100%	17	100%
Female adult or adolescent								
Injection drug use	1	4%	1	6%	-	-	-	-
Heterosexual contact	18	72%	13	76%	2	40%	3	100%
Other/unknown	6	24%	3	18%	3	60%	-	-
Subtotal	25	100%	17	100%	5	100%	3	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	215		146		49		20	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	138	64%	95	65%	33	67%	10	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	23	11%	13	9%	4	8%	6	30%
MSM & IDU	3	1%	3	2%	-	-	-	-
IDU & Heterosexual contact	2	1%	2	1%	-	-	-	-
MSM & Heterosexual contact	6	3%	6	4%	-	-	-	-
MSM & IDU & Heterosexual contact	3	1%	3	2%	-	-	-	-
Perinatal exposure	1	<1%	1	1%	-	-	-	-
Other/unknown	39	18%	23	16%	12	24%	4	20%
Total	215		146		49		20	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	63	78%	67	78%	8	57%	2	67%	-	-	4	80%
Injection drug use (IDU)	1	1%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	3	4%	1	1%	1	7%	-	-	-	-	1	20%
Heterosexual contact	1	1%	3	3%	1	7%	-	-	-	-	-	-
Other/unknown	13	16%	15	17%	4	29%	1	33%	-	-	-	-
Subtotal	81	100%	86	100%	14	100%	3	100%	-	-	5	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	50%	-	-	-	-	-	-
Heterosexual contact	3	75%	12	71%	1	50%	-	-	1	100%	1	100%
Other/unknown	1	25%	5	29%	-	-	-	-	-	-	-	-
Subtotal	4	100%	17	100%	2	100%	-	-	1	100%	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	-	-	-	-
Total	85		103		16		4		1		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	61	72%	64	62%	8	50%	1	25%	-	-	4	67%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	4	5%	15	15%	2	13%	-	-	1	100%	1	17%
MSM & IDU	1	1%	1	1%	1	6%	-	-	-	-	-	-
IDU & Heterosexual contact	1	1%	-	-	1	6%	-	-	-	-	-	-
MSM & Heterosexual contact	2	2%	3	3%	-	-	1	25%	-	-	-	-
MSM & IDU & Heterosexual contact	2	2%	-	-	-	-	-	-	-	-	1	17%
Perinatal exposure	-	-	-	-	-	-	1	25%	-	-	-	-
Other/unknown	14	16%	20	19%	4	25%	1	25%	-	-	-	-
Total	85		103		16		4		1		6	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 5

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	36.2	217	85%	35.3	212	81%	34.2	205	83%	30.8	185	81%
Females	5.9	37	15%	7.8	49	19%	6.5	41	17%	6.7	42	19%
Age at diagnosis (yr)												
<13	*	-	-	*	1	<1%	*	1	<1%	*	1	<1%
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	15.3	12	5%	20.3	16	6%	19.1	15	6%	19.1	15	7%
20-24	56.1	52	20%	53.9	50	19%	56.1	52	21%	60.4	56	25%
25-29	32.9	39	15%	35.5	42	16%	37.1	44	18%	27.0	32	14%
30-34	36.2	36	14%	42.2	42	16%	34.1	34	14%	22.1	22	10%
35-39	41.6	35	14%	42.8	36	14%	28.5	24	10%	22.6	19	8%
40-44	38.6	31	12%	34.9	28	11%	31.2	25	10%	31.2	25	11%
45-49	31.4	24	9%	22.2	17	7%	27.4	21	9%	28.7	22	10%
50-54	12.4	10	4%	18.5	15	6%	14.8	12	5%	24.7	20	9%
55-64	10.1	14	6%	7.9	11	4%	11.5	16	7%	9.3	13	6%
65+	*	1	<1%	*	3	1%	*	2	1%	*	2	1%
Race/Ethnicity												
White, not Hispanic	14.9	121	48%	14.2	115	44%	12.6	102	41%	11.5	93	41%
Black/African American, not Hispanic	40.8	109	43%	41.5	111	43%	43.4	116	47%	38.2	102	45%
Hispanic/Latino	20.7	13	5%	33.4	21	8%	12.7	8	3%	20.7	13	6%
Asian/Pacific Islander	*	2	1%	*	2	1%	*	4	2%	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	9	4%	*	12	5%	*	16	7%	*	17	7%
Race/Sex												
White, not Hispanic Males	27.8	110	43%	25.7	102	39%	24.5	97	39%	21.2	84	37%
White, not Hispanic Females	2.7	11	4%	3.1	13	5%	1.2	5	2%	2.2	9	4%
Black/African American, not Hispanic Males	68.3	86	34%	63.5	80	31%	67.5	85	35%	61.1	77	34%
Black/African American, not Hispanic Females	16.3	23	9%	22.0	31	12%	22.0	31	13%	17.7	25	11%
Hispanic/Latino Males	36.4	12	5%	51.6	17	7%	24.3	8	3%	27.3	9	4%
Hispanic/Latina Females	*	1	<1%	*	4	2%	*	-	-	*	4	2%
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	3	1%	*	1	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	9	4%	*	12	5%	*	16	7%	*	17	7%
Total	20.6	254		21.2	261		20.0	246		18.4	227	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 5

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	156	72%	150	71%	147	72%	132	71%
Injection drug use (IDU)	-	-	2	1%	1	<1%	-	-
Male-to-male sexual contact and IDU	8	4%	6	3%	2	1%	7	4%
Heterosexual contact	9	4%	3	1%	4	2%	7	4%
Other/unknown	44	20%	51	24%	50	25%	39	21%
Subtotal	217	100%	212	100%	204	100%	185	100%
Female adult or adolescent								
Injection drug use	3	8%	-	-	1	2%	5	12%
Heterosexual contact	27	73%	31	65%	23	56%	22	54%
Other/unknown	7	19%	17	35%	17	41%	14	34%
Subtotal	37	100%	48	100%	41	100%	41	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	1	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	1	100%	1	100%
Total	254		261		246		227	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 5

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	152	60%	148	57%	138	56%	123	54%
Injection drug use (IDU) only	-	-	2	1%	2	1%	1	<1%
Heterosexual contact only	36	14%	34	13%	27	11%	29	13%
MSM & IDU	8	3%	6	2%	2	1%	3	1%
IDU & Heterosexual contact	3	1%	-	-	-	-	4	2%
MSM & Heterosexual contact	4	2%	2	1%	9	4%	9	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	4	2%
Perinatal exposure	-	-	1	<1%	1	<1%	1	<1%
Other/unknown	51	20%	68	26%	67	27%	53	23%
Total	254		261		246		227	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 5

Year of HIV (not AIDS) Diagnosis

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	132	87%	132	81%	128	81%	128	83%	128	88%
Females	20	13%	31	19%	30	19%	27	17%	18	12%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	1	1%	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	9	6%	13	8%	13	8%	13	8%	6	4%
20-24	38	25%	36	22%	42	27%	43	28%	33	23%
25-29	30	20%	30	18%	26	16%	26	17%	35	24%
30-34	15	10%	28	17%	21	13%	16	10%	22	15%
35-39	21	14%	12	7%	13	8%	13	8%	20	14%
40-44	15	10%	17	10%	14	9%	15	10%	10	7%
45-49	11	7%	9	6%	14	9%	13	8%	12	8%
50-54	6	4%	10	6%	6	4%	6	4%	2	1%
55-64	6	4%	5	3%	9	6%	8	5%	4	3%
65+	1	1%	3	2%	-	-	1	1%	1	1%
Race/Ethnicity										
White, not Hispanic	74	49%	80	49%	68	43%	65	42%	58	40%
Black/African American, not Hispanic	64	42%	66	40%	77	49%	67	43%	70	48%
Hispanic/Latino	8	5%	10	6%	3	2%	10	6%	11	8%
Asian/Pacific Islander	2	1%	1	1%	4	3%	1	1%	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	1%
Unknown	4	3%	6	4%	6	4%	12	8%	4	3%
Race/Sex										
White, not Hispanic Males	67	44%	72	44%	64	41%	56	36%	56	38%
White, not Hispanic Females	7	5%	8	5%	4	3%	9	6%	2	1%
Black/African American, not Hispanic Males	53	35%	45	28%	54	34%	54	35%	59	40%
Black/African American, not Hispanic Females	11	7%	21	13%	23	15%	13	8%	11	8%
Hispanic/Latino Males	7	5%	9	6%	3	2%	7	5%	9	6%
Hispanic/Latina Females	1	1%	1	1%	-	-	3	2%	2	1%
Asian/Pacific Islander Males	1	1%	-	-	3	2%	1	1%	1	1%
Asian/Pacific Islander Females	1	1%	1	1%	1	1%	-	-	1	1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	1%
Unknown	4	3%	6	4%	6	4%	12	8%	4	3%
Total	152		163		158		155		146	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	99	75%	93	70%	98	77%	96	75%	101	79%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	1%
Male-to-male sexual contact and IDU	5	4%	4	3%	2	2%	7	5%	6	5%
Heterosexual contact	5	4%	1	1%	2	2%	3	2%	-	-
Other/unknown	23	17%	34	26%	26	20%	22	17%	20	16%
Subtotal	132	100%	132	100%	128	100%	128	100%	128	100%
Female adult or adolescent										
Injection drug use	1	5%	-	-	1	3%	5	19%	1	6%
Heterosexual contact	14	70%	22	71%	15	50%	15	58%	13	76%
Other/unknown	5	25%	9	29%	14	47%	6	23%	3	18%
Subtotal	20	100%	31	100%	30	100%	26	100%	17	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	1	100%
Total	152		163		158		155		146	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	98	64%	91	56%	92	58%	89	57%	95	65%
Injection drug use (IDU) only	-	-	-	-	1	1%	1	1%	-	-
Heterosexual contact only	19	13%	23	14%	17	11%	18	12%	13	9%
MSM & IDU	5	3%	4	2%	2	1%	3	2%	3	2%
IDU & Heterosexual contact	1	1%	-	-	-	-	4	3%	2	1%
MSM & Heterosexual contact	1	1%	2	1%	6	4%	7	5%	6	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	4	3%	3	2%
Perinatal exposure	-	-	-	-	-	-	1	1%	1	1%
Other/unknown	28	18%	43	26%	40	25%	28	18%	23	16%
Total	152		163		158		155		146	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 5

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	125	86%	117	80%	107	82%	109	83%	87	84%	3,279	87%
Females	21	14%	30	20%	23	18%	23	17%	16	16%	499	13%
Age at AIDS diagnosis (yr)												
<13	-	-	1	1%	-	-	-	-	-	-	23	1%
13-14	-	-	-	-	-	-	-	-	-	-	5	<1%
15-19	9	6%	8	5%	4	3%	3	2%	4	4%	75	2%
20-24	19	13%	18	12%	20	15%	32	24%	18	17%	379	10%
25-29	16	11%	16	11%	25	19%	16	12%	18	17%	644	17%
30-34	30	21%	24	16%	19	15%	16	12%	24	23%	851	23%
35-39	22	15%	34	23%	19	15%	18	14%	7	7%	664	18%
40-44	23	16%	21	14%	15	12%	13	10%	11	11%	486	13%
45-49	14	10%	13	9%	12	9%	12	9%	9	9%	327	9%
50-54	6	4%	6	4%	6	5%	13	10%	6	6%	178	5%
55-64	6	4%	6	4%	9	7%	8	6%	4	4%	123	3%
65+	1	1%	-	-	1	1%	1	1%	2	2%	23	1%
Race/Ethnicity												
White, not Hispanic	69	47%	62	42%	63	48%	56	42%	40	39%	2,287	61%
Black/African American, not Hispanic	69	47%	64	44%	47	36%	62	47%	50	49%	1,260	33%
Hispanic/Latino	4	3%	10	7%	8	6%	4	3%	6	6%	118	3%
Asian/Pacific Islander	-	-	1	1%	-	-	-	-	3	3%	16	<1%
American Indian/Alaska Native	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	4	3%	9	6%	12	9%	10	8%	4	4%	95	3%
Race/Sex												
White, not Hispanic Males	67	46%	56	38%	56	43%	51	39%	37	36%	2,146	57%
White, not Hispanic Females	2	1%	6	4%	7	5%	5	4%	3	3%	141	4%
Black/African American, not Hispanic Males	51	35%	45	31%	33	25%	46	35%	38	37%	935	25%
Black/African American, not Hispanic Females	18	12%	19	13%	14	11%	16	12%	12	12%	325	9%
Hispanic/Latino Males	4	3%	7	5%	8	6%	3	2%	6	6%	101	3%
Hispanic/Latina Females	-	-	3	2%	-	-	1	1%	-	-	17	<1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	-	-	2	2%	13	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	1%	3	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	4	3%	9	6%	12	9%	10	8%	4	4%	95	3%
Total	146		147		130		132		103		3,778	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	77	62%	80	68%	69	64%	70	64%	62	71%	2,403	74%
Injection drug use (IDU)	-	-	2	2%	2	2%	-	-	-	-	149	5%
Male-to-male sexual contact and IDU	6	5%	4	3%	2	2%	2	2%	-	-	118	4%
Heterosexual contact	9	7%	5	4%	3	3%	8	7%	7	8%	170	5%
Other/unknown	33	26%	26	22%	31	29%	29	27%	18	21%	428	13%
Subtotal	125	100%	117	100%	107	100%	109	100%	87	100%	3,268	100%
Female adult or adolescent												
Injection drug use	2	10%	-	-	1	4%	1	4%	1	6%	64	13%
Heterosexual contact	16	76%	20	69%	18	78%	12	52%	11	69%	327	68%
Other/unknown	3	14%	9	31%	4	17%	10	43%	4	25%	93	19%
Subtotal	21	100%	29	100%	23	100%	23	100%	16	100%	484	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	21	81%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	19%
Subtotal	-	-	1	100%	-	-	-	-	-	-	26	100%
Total	146		147		130		132		103		3,778	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	73	50%	78	53%	66	51%	68	52%	60	58%	2,325	62%
Injection drug use (IDU) only	-	-	2	1%	3	2%	-	-	-	-	151	4%
Heterosexual contact only	25	17%	25	17%	21	16%	20	15%	18	17%	497	13%
MSM & IDU	5	3%	4	3%	2	2%	2	2%	-	-	101	3%
IDU & Heterosexual contact	2	1%	-	-	-	-	1	1%	1	1%	62	2%
MSM & Heterosexual contact	4	3%	2	1%	3	2%	2	2%	2	2%	78	2%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-	-	-	17	<1%
Perinatal exposure	-	-	1	1%	-	-	-	-	-	-	22	1%
Other/unknown	36	25%	35	24%	35	27%	39	30%	22	21%	525	14%
Total	146		147		130		132		103		3,778	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 5

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 5

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	630.1	3,781	81%	2,015	81%	1,766	82%
Females	136.4	861	19%	483	19%	378	18%
Age at end of year							
<13	6.0	13	<1%	7	<1%	6	<1%
13-14	*	2	<1%	1	<1%	1	<1%
15-19	19.1	15	<1%	12	<1%	3	<1%
20-24	243.8	226	5%	171	7%	55	3%
25-29	376.5	446	10%	294	12%	152	7%
30-34	437.9	436	9%	284	11%	152	7%
35-39	561.0	472	10%	252	10%	220	10%
40-44	717.9	576	12%	312	12%	264	12%
45-49	1,008.6	772	17%	385	15%	387	18%
50-54	936.3	758	16%	335	13%	423	20%
55-64	530.5	739	16%	352	14%	387	18%
65+	139.8	187	4%	93	4%	94	4%
Race/Ethnicity							
White, not Hispanic	281.1	2,275	49%	1,255	50%	1,020	48%
Black/African American, not Hispanic	714.6	1,909	41%	1,016	41%	893	42%
Hispanic/Latino	362.4	228	5%	114	5%	114	5%
Asian/Pacific Islander	46.4	26	1%	14	1%	12	1%
American Indian/Alaska Native	202.0	5	<1%	4	<1%	1	<1%
Unknown	*	199	4%	95	4%	104	5%
Race/Sex							
White, not Hispanic Males	525.0	2,080	45%	1,135	45%	945	44%
White, not Hispanic Females	47.2	195	4%	120	5%	75	3%
Black/African American, not Hispanic Males	1,055.4	1,329	29%	704	28%	625	29%
Black/African American, not Hispanic Females	410.7	580	12%	312	12%	268	13%
Hispanic/Latino Males	540.1	178	4%	87	3%	91	4%
Hispanic/Latina Females	166.9	50	1%	27	1%	23	1%
Asian/Pacific Islander Males	76.0	21	<1%	11	<1%	10	<1%
Asian/Pacific Islander Females	17.6	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	199	4%	95	4%	104	5%
Total	377.0	4,642		2,498		2,144	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,702	72%	1,450	72%	1,252	71%
Injection drug use (IDU)	84	2%	24	1%	60	3%
Male-to-male sexual contact and IDU	121	3%	58	3%	63	4%
Heterosexual contact	196	5%	61	3%	135	8%
Other/unknown	663	18%	414	21%	249	14%
Subtotal	3,766	100%	2,007	100%	1,759	100%
Female adult or adolescent						
Injection drug use	71	9%	36	8%	35	10%
Heterosexual contact	547	66%	283	60%	264	72%
Other/unknown	217	26%	151	32%	66	18%
Subtotal	835	100%	470	100%	365	100%
Child (<13 yrs at diagnosis)						
Perinatal	34	83%	18	86%	16	80%
Other/unknown	7	17%	3	14%	4	20%
Subtotal	41	100%	21	100%	20	100%
Total	4,642		2,498		2,144	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,564	55%	1,384	55%	1,180	55%
Injection drug use (IDU) only	80	2%	26	1%	54	3%
Heterosexual contact only	743	16%	344	14%	399	19%
MSM & IDU	97	2%	44	2%	53	2%
IDU & Heterosexual contact	75	2%	34	1%	41	2%
MSM & Heterosexual contact	138	3%	66	3%	72	3%
MSM & IDU & Heterosexual contact	24	1%	14	1%	10	<1%
Perinatal exposure	36	1%	19	1%	17	1%
Other/unknown	885	19%	567	23%	318	15%
Total	4,642		2,498		2,144	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	1,649	79%	813	62%	112	64%	13	62%	3	100%	112	66%
Injection drug use (IDU)	26	1%	52	4%	4	2%	-	-	-	-	2	1%
Male-to-male sexual contact and IDU	70	3%	28	2%	9	5%	1	5%	-	-	13	8%
Heterosexual contact	45	2%	126	10%	18	10%	-	-	-	-	7	4%
Other/unknown	288	14%	299	23%	33	19%	7	33%	-	-	36	21%
Subtotal	2,078	100%	1,318	100%	176	100%	21	100%	3	100%	170	100%
Female adult or adolescent												
Injection drug use	19	10%	43	8%	7	15%	-	-	-	-	2	7%
Heterosexual contact	133	69%	358	64%	34	71%	2	50%	1	50%	19	68%
Other/unknown	41	21%	159	28%	7	15%	2	50%	1	50%	7	25%
Subtotal	193	100%	560	100%	48	100%	4	100%	2	100%	28	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	75%	26	84%	4	100%	1	100%	-	-	-	-
Other/unknown	1	25%	5	16%	-	-	-	-	-	-	1	100%
Subtotal	4	100%	31	100%	4	100%	1	100%	-	-	1	100%
Total	2,275		1,909		228		26		5		199	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 5

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	1,604	71%	738	39%	102	45%	12	46%	3	60%	105	53%
Injection drug use (IDU) only	20	1%	52	3%	7	3%	-	-	-	-	1	1%
Heterosexual contact only	178	8%	484	25%	52	23%	2	8%	1	20%	26	13%
MSM & IDU	54	2%	23	1%	9	4%	1	4%	-	-	10	5%
IDU & Heterosexual contact	25	1%	43	2%	4	2%	-	-	-	-	3	2%
MSM & Heterosexual contact	45	2%	75	4%	10	4%	1	4%	-	-	7	4%
MSM & IDU & Heterosexual contact	16	1%	5	0%	-	-	-	-	-	-	3	2%
Perinatal exposure	3	0%	27	1%	4	2%	1	4%	-	-	1	1%
Other/unknown	330	15%	462	24%	40	18%	9	35%	1	20%	43	22%
Total	2,275		1,909		228		26		5		199	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 5

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	3,009	81%	3,216	81%	3,410	81%	3,599	81%	3,781	81%
Females	704	19%	750	19%	789	19%	832	19%	861	19%
Age at end of year										
<13	12	<1%	12	<1%	13	<1%	14	<1%	13	<1%
13-14	2	<1%	3	<1%	1	<1%	-	-	2	<1%
15-19	31	1%	24	1%	23	1%	18	<1%	15	<1%
20-24	220	6%	229	6%	238	6%	246	6%	226	5%
25-29	298	8%	342	9%	374	9%	408	9%	446	10%
30-34	359	10%	403	10%	426	10%	422	10%	436	9%
35-39	439	12%	425	11%	432	10%	446	10%	472	10%
40-44	643	17%	630	16%	598	14%	588	13%	576	12%
45-49	727	20%	760	19%	775	18%	798	18%	772	17%
50-54	500	13%	564	14%	623	15%	677	15%	758	16%
55-64	407	11%	485	12%	586	14%	668	15%	739	16%
65+	75	2%	89	2%	110	3%	146	3%	187	4%
Race/Ethnicity										
White, not Hispanic	1,895	51%	2,005	51%	2,099	50%	2,194	50%	2,275	49%
Black/African American, not Hispanic	1,476	40%	1,584	40%	1,700	40%	1,805	41%	1,909	41%
Hispanic/Latino	173	5%	191	5%	198	5%	213	5%	228	5%
Asian/Pacific Islander	15	<1%	17	<1%	20	<1%	22	<1%	26	1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	5	<1%
Unknown	150	4%	165	4%	178	4%	193	4%	199	4%
Race/Sex										
White, not Hispanic Males	1,730	47%	1,827	46%	1,915	46%	2,003	45%	2,080	45%
White, not Hispanic Females	165	4%	178	4%	184	4%	191	4%	195	4%
Black/African American, not Hispanic Males	1,001	27%	1,081	27%	1,168	28%	1,245	28%	1,329	29%
Black/African American, not Hispanic Females	475	13%	503	13%	532	13%	560	13%	580	12%
Hispanic/Latino Males	134	4%	148	4%	155	4%	165	4%	178	4%
Hispanic/Latina Females	39	1%	43	1%	43	1%	48	1%	50	1%
Asian/Pacific Islander Males	14	<1%	15	<1%	17	<1%	18	<1%	21	<1%
Asian/Pacific Islander Females	1	<1%	2	<1%	3	<1%	4	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
Unknown	150	4%	165	4%	178	4%	193	4%	199	4%
Total	3,713		3,966		4,199		4,431		4,642	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 5

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	2,135	71%	2,286	71%	2,425	71%	2,563	72%	2,702	72%
Injection drug use (IDU)	78	3%	80	2%	83	2%	83	2%	84	2%
Male-to-male sexual contact and IDU	100	3%	104	3%	107	3%	114	3%	121	3%
Heterosexual contact	178	6%	181	6%	185	5%	192	5%	196	5%
Other/unknown	504	17%	551	17%	595	18%	632	18%	663	18%
Subtotal	2,995	100%	3,202	100%	3,395	100%	3,584	100%	3,766	100%
Female adult or adolescent										
Injection drug use	66	10%	66	9%	67	9%	70	9%	71	9%
Heterosexual contact	453	67%	482	66%	505	66%	528	65%	547	66%
Other/unknown	162	24%	178	25%	193	25%	209	26%	217	26%
Subtotal	681	100%	726	100%	765	100%	807	100%	835	100%
Child (<13 yrs at diagnosis)										
Perinatal	30	81%	31	82%	32	82%	33	83%	34	83%
Other/unknown	7	19%	7	18%	7	18%	7	18%	7	17%
Subtotal	37	100%	38	100%	39	100%	40	100%	41	100%
Total	3,713		3,966		4,199		4,431		4,642	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 5

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	2,023	54%	2,173	55%	2,300	55%	2,431	55%	2,564	55%
Injection drug use (IDU) only	75	2%	77	2%	80	2%	80	2%	80	2%
Heterosexual contact only	631	17%	663	17%	690	16%	720	16%	743	16%
MSM & IDU	83	2%	87	2%	89	2%	93	2%	97	2%
IDU & Heterosexual contact	69	2%	69	2%	70	2%	73	2%	75	2%
MSM & Heterosexual contact	112	3%	113	3%	125	3%	132	3%	138	3%
MSM & IDU & Heterosexual contact	17	<1%	17	<1%	18	<1%	21	<1%	24	1%
Perinatal exposure	32	1%	33	1%	34	1%	35	1%	36	1%
Other/unknown	671	18%	734	19%	793	19%	846	19%	885	19%
Total	3,713		3,966		4,199		4,431		4,642	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 6

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 6

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	30.7	119	80%	88	79%	24	83%	7	88%	
Females	6.9	29	20%	23	21%	5	17%	1	13%	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	15.2	8	5%	8	7%	-	-	-	-	
20-24	51.8	29	20%	23	21%	4	14%	2	25%	
25-29	57.3	36	24%	31	28%	5	17%	-	-	
30-34	32.7	18	12%	13	12%	4	14%	1	13%	
35-39	27.7	13	9%	11	10%	1	3%	1	13%	
40-44	25.4	12	8%	10	9%	1	3%	1	13%	
45-49	25.6	13	9%	5	5%	6	21%	2	25%	
50-54	10.4	6	4%	4	4%	2	7%	-	-	
55-64	11.3	12	8%	5	5%	6	21%	1	13%	
65+	*	1	1%	1	1%	-	-	-	-	
Race/Ethnicity										
White, not Hispanic	8.0	43	29%	33	30%	8	28%	2	25%	
Black/African American, not Hispanic	45.2	94	64%	67	60%	21	72%	6	75%	
Hispanic/Latino	25.6	6	4%	6	5%	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	5	3%	5	5%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	13.7	36	24%	26	23%	8	28%	2	25%	
White, not Hispanic Females	2.5	7	5%	7	6%	-	-	-	-	
Black/African American, not Hispanic Males	77.5	74	50%	53	48%	16	55%	5	63%	
Black/African American, not Hispanic Females	17.8	20	14%	14	13%	5	17%	1	13%	
Hispanic/Latino Males	41.2	5	3%	5	5%	-	-	-	-	
Hispanic/Latina Females	*	1	1%	1	1%	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	5	3%	5	5%	-	-	-	-	
Total	18.3	148		111		29		8		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	73	61%	52	59%	17	71%	4	57%
Injection drug use (IDU)	5	4%	4	5%	1	4%	-	-
Male-to-male sexual contact and IDU	3	3%	3	3%	-	-	-	-
Heterosexual contact	7	6%	4	5%	1	4%	2	29%
Other/unknown	31	26%	25	28%	5	21%	1	14%
Subtotal	119	100%	88	100%	24	100%	7	100%
Female adult or adolescent								
Injection drug use	5	17%	5	22%	-	-	-	-
Heterosexual contact	15	52%	11	48%	4	80%	-	-
Other/unknown	9	31%	7	30%	1	20%	1	100%
Subtotal	29	100%	23	100%	5	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	148		111		29		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	68	46%	47	42%	17	59%	4	50%
Injection drug use (IDU) only	2	1%	1	1%	1	3%	-	-
Heterosexual contact only	22	15%	15	14%	5	17%	2	25%
MSM & IDU	3	2%	3	3%	-	-	-	-
IDU & Heterosexual contact	8	5%	8	7%	-	-	-	-
MSM & Heterosexual contact	5	3%	5	5%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	40	27%	32	29%	6	21%	2	25%
Total	148		111		29		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	23	64%	44	59%	4	80%	-	-	-	-	2	50%
Injection drug use (IDU)	4	11%	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	6%	-	-	-	-	-	-	-	-	1	25%
Heterosexual contact	-	-	7	9%	-	-	-	-	-	-	-	-
Other/unknown	7	19%	22	30%	1	20%	-	-	-	-	1	25%
Subtotal	36	100%	74	100%	5	100%	-	-	-	-	4	100%
Female adult or adolescent												
Injection drug use	5	71%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	14%	12	60%	1	100%	-	-	-	-	1	100%
Other/unknown	1	14%	8	40%	-	-	-	-	-	-	-	-
Subtotal	7	100%	20	100%	1	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	43		94		6		0		0		5	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	23	53%	40	43%	3	50%	-	-	-	-	2	40%
Injection drug use (IDU) only	2	5%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	2%	19	20%	1	17%	-	-	-	-	1	20%
MSM & IDU	2	5%	-	-	-	-	-	-	-	-	1	20%
IDU & Heterosexual contact	7	16%	1	1%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	4	4%	1	17%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	8	19%	30	32%	1	17%	-	-	-	-	1	20%
Total	43		94		6		0		0		5	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 6

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	30.1	117	76%	25.2	98	72%	27.8	108	73%	33.8	131	80%
Females	8.6	36	24%	9.1	38	28%	9.3	39	27%	7.9	33	20%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	1	1%	*	1	1%
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	32.4	17	11%	22.9	12	9%	24.8	13	9%	28.6	15	9%
20-24	60.7	34	22%	58.9	33	24%	66.0	37	25%	75.0	42	26%
25-29	43.0	27	18%	33.5	21	15%	30.3	19	13%	49.4	31	19%
30-34	29.1	16	10%	29.1	16	12%	38.1	21	14%	30.9	17	10%
35-39	29.8	14	9%	21.3	10	7%	21.3	10	7%	14.9	7	4%
40-44	33.9	16	10%	21.2	10	7%	23.3	11	7%	25.4	12	7%
45-49	39.4	20	13%	25.6	13	10%	33.5	17	12%	25.6	13	8%
50-54	13.9	8	5%	22.5	13	10%	12.1	7	5%	19.1	11	7%
55-64	*	1	1%	5.6	6	4%	8.5	9	6%	12.2	13	8%
65+	*	-	-	*	2	1%	*	2	1%	*	2	1%
Race/Ethnicity												
White, not Hispanic	8.4	45	29%	7.6	41	30%	6.1	33	22%	10.8	58	35%
Black/African American, not Hispanic	50.0	104	68%	43.3	90	66%	50.0	104	71%	47.1	98	60%
Hispanic/Latino	*	-	-	*	3	2%	*	4	3%	*	4	2%
Asian/Pacific Islander	*	-	-	*	2	1%	*	-	-	*	3	2%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	3%	*	-	-	*	6	4%	*	1	1%
Race/Sex												
White, not Hispanic Males	13.3	35	23%	12.6	33	24%	10.3	27	18%	18.7	49	30%
White, not Hispanic Females	3.6	10	7%	2.9	8	6%	2.2	6	4%	3.3	9	5%
Black/African American, not Hispanic Males	83.8	80	52%	65.0	62	46%	76.5	73	50%	78.6	75	46%
Black/African American, not Hispanic Females	21.3	24	16%	24.9	28	21%	27.5	31	21%	20.4	23	14%
Hispanic/Latino Males	*	-	-	*	2	1%	*	3	2%	*	3	2%
Hispanic/Latina Females	*	-	-	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Males	*	-	-	*	1	1%	*	-	-	*	3	2%
Asian/Pacific Islander Females	*	-	-	*	1	1%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	3%	*	-	-	*	6	4%	*	1	1%
Total	19.0	153		16.9	136		18.2	147		20.3	164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 6

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	90	77%	64	65%	78	73%	93	72%
Injection drug use (IDU)	-	-	-	-	1	1%	5	4%
Male-to-male sexual contact and IDU	3	3%	2	2%	1	1%	6	5%
Heterosexual contact	10	9%	3	3%	2	2%	3	2%
Other/unknown	14	12%	29	30%	25	23%	23	18%
Subtotal	117	100%	98	100%	107	100%	130	100%
Female adult or adolescent								
Injection drug use	3	8%	1	3%	2	5%	5	16%
Heterosexual contact	29	81%	27	71%	30	79%	24	75%
Other/unknown	4	11%	10	26%	6	16%	3	9%
Subtotal	36	100%	38	100%	38	100%	32	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	1	50%
Other/unknown	-	-	-	-	2	100%	1	50%
Subtotal	-	-	-	-	2	100%	2	100%
Total	153		136		147		164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 6

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	87	57%	60	44%	72	49%	91	55%
Injection drug use (IDU) only	2	1%	1	1%	1	1%	3	2%
Heterosexual contact only	39	25%	30	22%	32	22%	27	16%
MSM & IDU	3	2%	2	1%	-	-	5	3%
IDU & Heterosexual contact	1	1%	-	-	2	1%	7	4%
MSM & Heterosexual contact	3	2%	4	3%	6	4%	2	1%
MSM & IDU & Heterosexual contact	-	-	-	-	1	1%	1	1%
Perinatal exposure	-	-	-	-	1	1%	2	1%
Other/unknown	18	12%	39	29%	32	22%	26	16%
Total	153		136		147		164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	61	71%	62	69%	61	69%	85	77%	88	79%
Females	25	29%	28	31%	27	31%	25	23%	23	21%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	1	1%	1	1%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	16	19%	11	12%	12	14%	14	13%	8	7%
20-24	24	28%	24	27%	28	32%	34	31%	23	21%
25-29	13	15%	16	18%	11	13%	19	17%	31	28%
30-34	9	10%	8	9%	11	13%	13	12%	13	12%
35-39	7	8%	5	6%	4	5%	3	3%	11	10%
40-44	6	7%	7	8%	6	7%	6	5%	10	9%
45-49	8	9%	9	10%	10	11%	7	6%	5	5%
50-54	3	3%	6	7%	1	1%	5	5%	4	4%
55-64	-	-	3	3%	3	3%	7	6%	5	5%
65+	-	-	1	1%	1	1%	1	1%	1	1%
Race/Ethnicity										
White, not Hispanic	30	35%	28	31%	19	22%	38	35%	33	30%
Black/African American, not Hispanic	53	62%	59	66%	63	72%	67	61%	67	60%
Hispanic/Latino	-	-	1	1%	3	3%	3	3%	6	5%
Asian/Pacific Islander	-	-	2	2%	-	-	1	1%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	-	-	3	3%	1	1%	5	5%
Race/Sex										
White, not Hispanic Males	22	26%	20	22%	13	15%	32	29%	26	23%
White, not Hispanic Females	8	9%	8	9%	6	7%	6	5%	7	6%
Black/African American, not Hispanic Males	38	44%	41	46%	43	49%	49	45%	53	48%
Black/African American, not Hispanic Females	15	17%	18	20%	20	23%	18	16%	14	13%
Hispanic/Latino Males	-	-	-	-	2	2%	2	2%	5	5%
Hispanic/Latina Females	-	-	1	1%	1	1%	1	1%	1	1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	1	1%	-	-
Asian/Pacific Islander Females	-	-	1	1%	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	-	-	3	3%	1	1%	5	5%
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	49	80%	39	63%	43	72%	60	71%	52	59%
Injection drug use (IDU)	-	-	-	-	-	-	4	5%	4	5%
Male-to-male sexual contact and IDU	2	3%	1	2%	1	2%	5	6%	3	3%
Heterosexual contact	7	11%	3	5%	2	3%	2	2%	4	5%
Other/unknown	3	5%	19	31%	14	23%	14	16%	25	28%
Subtotal	61	100%	62	100%	60	100%	85	100%	88	100%
Female adult or adolescent										
Injection drug use	3	12%	1	4%	2	8%	4	17%	5	22%
Heterosexual contact	20	80%	19	68%	21	81%	17	71%	11	48%
Other/unknown	2	8%	8	29%	3	12%	3	13%	7	30%
Subtotal	25	100%	28	100%	26	100%	24	100%	23	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	-	-
Other/unknown	-	-	-	-	2	100%	-	-	-	-
Subtotal	-	-	-	-	2	100%	1	100%	-	-
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	48	56%	35	39%	38	43%	59	54%	47	42%
Injection drug use (IDU) only	2	2%	1	1%	1	1%	3	3%	1	1%
Heterosexual contact only	27	31%	22	24%	23	26%	19	17%	15	14%
MSM & IDU	2	2%	1	1%	-	-	5	5%	3	3%
IDU & Heterosexual contact	1	1%	-	-	1	1%	5	5%	8	7%
MSM & Heterosexual contact	1	1%	4	4%	5	6%	1	1%	5	5%
MSM & IDU & Heterosexual contact	-	-	-	-	1	1%	-	-	-	-
Perinatal exposure	-	-	-	-	1	1%	1	1%	-	-
Other/unknown	5	6%	27	30%	18	20%	17	15%	32	29%
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	70	81%	56	76%	56	81%	62	85%	47	78%	2,156	86%
Females	16	19%	18	24%	13	19%	11	15%	13	22%	350	14%
Age at AIDS diagnosis (yr)												
<13	-	-	1	1%	-	-	-	-	-	-	19	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	3	3%	5	7%	2	3%	1	1%	2	3%	60	2%
20-24	12	14%	18	24%	13	19%	12	16%	12	20%	240	10%
25-29	18	21%	7	9%	8	12%	19	26%	7	12%	443	18%
30-34	12	14%	11	15%	8	12%	7	10%	5	8%	522	21%
35-39	6	7%	8	11%	9	13%	5	7%	6	10%	447	18%
40-44	10	12%	6	8%	9	13%	6	8%	3	5%	328	13%
45-49	13	15%	7	9%	7	10%	8	11%	11	18%	205	8%
50-54	7	8%	5	7%	6	9%	7	10%	6	10%	122	5%
55-64	4	5%	5	7%	6	9%	7	10%	8	13%	94	4%
65+	1	1%	1	1%	1	1%	1	1%	-	-	24	1%
Race/Ethnicity												
White, not Hispanic	25	29%	21	28%	16	23%	22	30%	16	27%	1,144	46%
Black/African American, not Hispanic	59	69%	51	69%	47	68%	47	64%	42	70%	1,297	52%
Hispanic/Latino	-	-	1	1%	2	3%	1	1%	1	2%	29	1%
Asian/Pacific Islander	-	-	-	-	-	-	2	3%	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	2	2%	1	1%	4	6%	1	1%	1	2%	26	1%
Race/Sex												
White, not Hispanic Males	23	27%	17	23%	16	23%	19	26%	14	23%	1,072	43%
White, not Hispanic Females	2	2%	4	5%	-	-	3	4%	2	3%	72	3%
Black/African American, not Hispanic Males	45	52%	37	50%	35	51%	39	53%	31	52%	1,030	41%
Black/African American, not Hispanic Females	14	16%	14	19%	12	17%	8	11%	11	18%	267	11%
Hispanic/Latino Males	-	-	1	1%	2	3%	1	1%	1	2%	23	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	6	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	2	3%	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	1	1%	4	6%	1	1%	1	2%	26	1%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	54	77%	39	70%	42	75%	47	77%	32	68%	1,576	74%
Injection drug use (IDU)	-	-	-	-	1	2%	1	2%	1	2%	91	4%
Male-to-male sexual contact and IDU	3	4%	1	2%	-	-	1	2%	1	2%	127	6%
Heterosexual contact	3	4%	1	2%	-	-	2	3%	5	11%	86	4%
Other/unknown	10	14%	15	27%	13	23%	10	16%	8	17%	262	12%
Subtotal	70	100%	56	100%	56	100%	61	100%	47	100%	2,142	100%
Female adult or adolescent												
Injection drug use	1	6%	2	12%	1	8%	-	-	1	8%	52	15%
Heterosexual contact	13	81%	13	76%	10	77%	11	100%	9	69%	249	72%
Other/unknown	2	13%	2	12%	2	15%	-	-	3	23%	43	13%
Subtotal	16	100%	17	100%	13	100%	11	100%	13	100%	344	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	14	70%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	6	30%
Subtotal	-	-	1	100%	-	-	1	100%	-	-	20	100%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	53	62%	39	53%	40	58%	46	63%	32	53%	1,508	60%
Injection drug use (IDU) only	-	-	2	3%	-	-	-	-	1	2%	85	3%
Heterosexual contact only	16	19%	14	19%	10	14%	13	18%	14	23%	335	13%
MSM & IDU	2	2%	1	1%	-	-	-	-	1	2%	112	4%
IDU & Heterosexual contact	1	1%	-	-	2	3%	1	1%	1	2%	58	2%
MSM & Heterosexual contact	1	1%	-	-	2	3%	1	1%	-	-	66	3%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	1	1%	-	-	15	1%
Perinatal exposure	-	-	1	1%	-	-	1	1%	-	-	15	1%
Other/unknown	12	14%	17	23%	15	22%	10	14%	11	18%	312	12%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 6

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	558.3	2,167	78%	1,039	76%	1,128	80%
Females	143.4	600	22%	326	24%	274	20%
Age at end of year							
<13	4.4	6	<1%	6	<1%	-	-
13-14	*	-	-	-	-	-	-
15-19	49.5	26	1%	24	2%	2	<1%
20-24	278.4	156	6%	126	9%	30	2%
25-29	468.3	294	11%	193	14%	101	7%
30-34	386.9	213	8%	111	8%	102	7%
35-39	540.9	254	9%	137	10%	117	8%
40-44	557.5	263	10%	128	9%	135	10%
45-49	843.8	428	15%	173	13%	255	18%
50-54	831.4	480	17%	220	16%	260	19%
55-64	486.7	517	19%	199	15%	318	23%
65+	113.8	130	5%	48	4%	82	6%
Race/Ethnicity							
White, not Hispanic	189.7	1,021	37%	515	38%	506	36%
Black/African American, not Hispanic	776.1	1,615	58%	783	57%	832	59%
Hispanic/Latino	256.1	60	2%	28	2%	32	2%
Asian/Pacific Islander	78.0	15	1%	8	1%	7	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	55	2%	31	2%	24	2%
Race/Sex							
White, not Hispanic Males	330.3	867	31%	418	31%	449	32%
White, not Hispanic Females	55.9	154	6%	97	7%	57	4%
Black/African American, not Hispanic Males	1,249.0	1,192	43%	568	42%	624	45%
Black/African American, not Hispanic Females	375.5	423	15%	215	16%	208	15%
Hispanic/Latino Males	395.7	48	2%	19	1%	29	2%
Hispanic/Latina Females	106.2	12	<1%	9	1%	3	<1%
Asian/Pacific Islander Males	147.6	14	1%	8	1%	6	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	55	2%	31	2%	24	2%
Total	343.0	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,496	69%	686	66%	810	72%
Injection drug use (IDU)	75	3%	31	3%	44	4%
Male-to-male sexual contact and IDU	98	5%	37	4%	61	5%
Heterosexual contact	119	6%	48	5%	71	6%
Other/unknown	368	17%	231	22%	137	12%
Subtotal	2,156	100%	1,033	100%	1,123	100%
Female adult or adolescent						
Injection drug use	70	12%	33	10%	37	14%
Heterosexual contact	417	71%	217	68%	200	74%
Other/unknown	100	17%	68	21%	32	12%
Subtotal	587	100%	318	100%	269	100%
Child (<13 yrs at diagnosis)						
Perinatal	17	71%	11	79%	6	60%
Other/unknown	7	29%	3	21%	4	40%
Subtotal	24	100%	14	100%	10	100%
Total	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	1,418	51%	652	48%	766	55%
Injection drug use (IDU) only	82	3%	31	2%	51	4%
Heterosexual contact only	536	19%	265	19%	271	19%
MSM & IDU	82	3%	31	2%	51	4%
IDU & Heterosexual contact	63	2%	33	2%	30	2%
MSM & Heterosexual contact	78	3%	34	2%	44	3%
MSM & IDU & Heterosexual contact	16	1%	6	<1%	10	1%
Perinatal exposure	19	1%	12	1%	7	<1%
Other/unknown	473	17%	301	22%	172	12%
Total	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	689	80%	739	62%	25	52%	11	79%	1	100%	31	69%
Injection drug use (IDU)	19	2%	48	4%	7	15%	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	50	6%	44	4%	3	6%	-	-	-	-	1	2%
Heterosexual contact	11	1%	102	9%	4	8%	1	7%	-	-	1	2%
Other/unknown	95	11%	251	21%	9	19%	2	14%	-	-	11	24%
Subtotal	864	100%	1,184	100%	48	100%	14	100%	1	100%	45	100%
Female adult or adolescent												
Injection drug use	36	24%	30	7%	3	25%	-	-	-	-	1	10%
Heterosexual contact	90	60%	310	75%	9	75%	1	100%	-	-	7	70%
Other/unknown	25	17%	73	18%	-	-	-	-	-	-	2	20%
Subtotal	151	100%	413	100%	12	100%	1	100%	-	-	10	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	50%	14	78%	-	-	-	-	-	-	-	-
Other/unknown	3	50%	4	22%	-	-	-	-	-	-	-	-
Subtotal	6	100%	18	100%	-	-	-	-	-	-	-	-
Total	1,021		1,615		60		15		1		55	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 6

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	666	65%	690	43%	21	35%	11	73%	1	100%	29	53%
Injection drug use (IDU) only	28	3%	46	3%	8	13%	-	-	-	-	-	-
Heterosexual contact only	101	10%	412	26%	13	22%	2	13%	-	-	8	15%
MSM & IDU	45	4%	34	2%	2	3%	-	-	-	-	1	2%
IDU & Heterosexual contact	27	3%	32	2%	2	3%	-	-	-	-	2	4%
MSM & Heterosexual contact	23	2%	49	3%	4	7%	-	-	-	-	2	4%
MSM & IDU & Heterosexual contact	5	0%	10	1%	1	2%	-	-	-	-	-	-
Perinatal exposure	3	0%	16	1%	-	-	-	-	-	-	-	-
Other/unknown	123	12%	326	20%	9	15%	2	13%	-	-	13	24%
Total	1,021		1,615		60		15		1		55	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 6

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	1,700	79%	1,802	78%	1,916	78%	2,046	78%	2,167	78%
Females	460	21%	499	22%	537	22%	570	22%	600	22%
Age at end of year										
<13	3	<1%	4	<1%	5	<1%	6	<1%	6	<1%
13-14	1	<1%	-	-	1	<1%	1	<1%	-	-
15-19	32	1%	35	2%	30	1%	28	1%	26	1%
20-24	124	6%	150	7%	177	7%	180	7%	156	6%
25-29	152	7%	169	7%	191	8%	239	9%	294	11%
30-34	176	8%	181	8%	187	8%	196	7%	213	8%
35-39	216	10%	225	10%	228	9%	228	9%	254	9%
40-44	348	16%	309	13%	297	12%	276	11%	263	10%
45-49	429	20%	446	19%	454	19%	453	17%	428	15%
50-54	337	16%	379	16%	418	17%	456	17%	480	17%
55-64	276	13%	327	14%	372	15%	441	17%	517	19%
65+	66	3%	76	3%	93	4%	112	4%	130	5%
Race/Ethnicity										
White, not Hispanic	833	39%	879	38%	916	37%	975	37%	1,021	37%
Black/African American, not Hispanic	1,232	57%	1,323	57%	1,426	58%	1,521	58%	1,615	58%
Hispanic/Latino	44	2%	46	2%	50	2%	54	2%	60	2%
Asian/Pacific Islander	11	1%	12	1%	12	<1%	15	1%	15	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	39	2%	40	2%	48	2%	50	2%	55	2%
Race/Sex										
White, not Hispanic Males	709	33%	746	32%	777	32%	827	32%	867	31%
White, not Hispanic Females	124	6%	133	6%	139	6%	148	6%	154	6%
Black/African American, not Hispanic Males	912	42%	974	42%	1,047	43%	1,119	43%	1,192	43%
Black/African American, not Hispanic Females	320	15%	349	15%	379	15%	402	15%	423	15%
Hispanic/Latino Males	37	2%	38	2%	41	2%	44	2%	48	2%
Hispanic/Latina Females	7	<1%	8	<1%	9	<1%	10	<1%	12	<1%
Asian/Pacific Islander Males	10	<1%	11	<1%	11	<1%	14	1%	14	1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	39	2%	40	2%	48	2%	50	2%	55	2%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 6

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,175	69%	1,245	69%	1,331	70%	1,424	70%	1,496	69%
Injection drug use (IDU)	63	4%	64	4%	66	3%	70	3%	75	3%
Male-to-male sexual contact and IDU	83	5%	86	5%	88	5%	94	5%	98	5%
Heterosexual contact	105	6%	107	6%	109	6%	112	6%	119	6%
Other/unknown	265	16%	291	16%	312	16%	335	16%	368	17%
Subtotal	1,691	100%	1,793	100%	1,906	100%	2,035	100%	2,156	100%
Female adult or adolescent										
Injection drug use	57	13%	58	12%	60	11%	65	12%	70	12%
Heterosexual contact	320	71%	348	71%	377	72%	401	72%	417	71%
Other/unknown	74	16%	83	17%	88	17%	91	16%	100	17%
Subtotal	451	100%	489	100%	525	100%	557	100%	587	100%
Child (<13 yrs at diagnosis)										
Perinatal	15	83%	16	84%	16	73%	17	71%	17	71%
Other/unknown	3	17%	3	16%	6	27%	7	29%	7	29%
Subtotal	18	100%	19	100%	22	100%	24	100%	24	100%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 6

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	1,115	52%	1,182	51%	1,260	51%	1,351	52%	1,418	51%
Injection drug use (IDU) only	74	3%	75	3%	77	3%	80	3%	82	3%
Heterosexual contact only	425	20%	455	20%	486	20%	513	20%	536	19%
MSM & IDU	69	3%	72	3%	73	3%	78	3%	82	3%
IDU & Heterosexual contact	46	2%	47	2%	49	2%	55	2%	63	2%
MSM & Heterosexual contact	60	3%	63	3%	71	3%	73	3%	78	3%
MSM & IDU & Heterosexual contact	14	1%	14	1%	15	1%	16	1%	16	1%
Perinatal exposure	15	1%	16	1%	17	1%	19	1%	19	1%
Other/unknown	342	16%	377	16%	405	17%	431	16%	473	17%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 7

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 7

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	11.1	72	77%	57	80%	13	68%	2	67%
Females	3.1	21	23%	14	20%	6	32%	1	33%
Age at diagnosis (yr)									
<13	*	1	1%	1	1%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	6.6	6	6%	6	8%	-	-	-	-
20-24	27.2	25	27%	23	32%	2	11%	-	-
25-29	16.5	13	14%	11	15%	2	11%	-	-
30-34	6.1	5	5%	4	6%	1	5%	-	-
35-39	18.0	14	15%	8	11%	4	21%	2	67%
40-44	11.8	10	11%	8	11%	1	5%	1	33%
45-49	6.8	6	6%	3	4%	3	16%	-	-
50-54	7.2	7	8%	3	4%	4	21%	-	-
55-64	*	3	3%	2	3%	1	5%	-	-
65+	*	3	3%	2	3%	1	5%	-	-
Race/Ethnicity									
White, not Hispanic	3.9	42	45%	30	42%	10	53%	2	67%
Black/African American, not Hispanic	29.9	45	48%	35	49%	9	47%	1	33%
Hispanic/Latino	*	4	4%	4	6%	-	-	-	-
Asian/Pacific Islander	*	1	1%	1	1%	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	1%	1	1%	-	-	-	-
Race/Sex									
White, not Hispanic Males	7.2	38	41%	28	39%	8	42%	2	67%
White, not Hispanic Females	*	4	4%	2	3%	2	11%	-	-
Black/African American, not Hispanic Males	40.1	29	31%	24	34%	5	26%	-	-
Black/African American, not Hispanic Females	20.4	16	17%	11	15%	4	21%	1	33%
Hispanic/Latino Males	*	3	3%	3	4%	-	-	-	-
Hispanic/Latina Females	*	1	1%	1	1%	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	1%	1	1%	-	-	-	-
Total	7.0	93		71		19		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	56	78%	46	81%	9	69%	1	50%
Injection drug use (IDU)	1	1%	-	-	1	8%	-	-
Male-to-male sexual contact and IDU	1	1%	1	2%	-	-	-	-
Heterosexual contact	1	1%	-	-	1	8%	-	-
Other/unknown	13	18%	10	18%	2	15%	1	50%
Subtotal	72	100%	57	100%	13	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	10	50%	8	62%	2	33%	-	-
Other/unknown	10	50%	5	38%	4	67%	1	100%
Subtotal	20	100%	13	100%	6	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	93		71		19		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	56	60%	46	65%	9	47%	1	33%
Injection drug use (IDU) only	1	1%	-	-	1	5%	-	-
Heterosexual contact only	11	12%	8	11%	3	16%	-	-
MSM & IDU	1	1%	1	1%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	1	1%	-	-	-	-
Other/unknown	23	25%	15	21%	6	32%	2	67%
Total	93		71		19		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	29	76%	24	83%	1	33%	1	100%	-	-	1	100%
Injection drug use (IDU)	1	3%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	3%	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	16%	5	17%	2	67%	-	-	-	-	-	-
Subtotal	38	100%	29	100%	3	100%	1	100%	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	50%	8	53%	-	-	-	-	-	-	-	-
Other/unknown	2	50%	7	47%	1	100%	-	-	-	-	-	-
Subtotal	4	100%	15	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-	-	-
Total	42		45		4		1		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	29	69%	24	53%	1	25%	1	100%	-	-	1	100%
Injection drug use (IDU) only	1	2%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	7%	8	18%	-	-	-	-	-	-	-	-
MSM & IDU	1	2%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	2%	-	-	-	-	-	-	-	-
Other/unknown	8	19%	12	27%	3	75%	-	-	-	-	-	-
Total	42		45		4		1		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 7

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex												
Males	13.0	84	82%	13.0	84	75%	10.0	65	81%	13.1	85	77%
Females	2.8	19	18%	4.1	28	25%	2.2	15	19%	3.7	25	23%
Age at diagnosis (yr)												
<13	*	2	2%	*	-	-	*	-	-	*	3	3%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	11.0	10	10%	5.5	5	4%	*	2	3%	8.8	8	7%
20-24	25.0	23	22%	30.4	28	25%	18.5	17	21%	20.6	19	17%
25-29	24.1	19	18%	15.2	12	11%	25.4	20	25%	29.2	23	21%
30-34	6.1	5	5%	20.8	17	15%	7.3	6	8%	13.5	11	10%
35-39	14.1	11	11%	15.4	12	11%	14.1	11	14%	23.1	18	16%
40-44	11.8	10	10%	14.1	12	11%	*	4	5%	13.0	11	10%
45-49	10.2	9	9%	12.5	11	10%	10.2	9	11%	7.9	7	6%
50-54	7.2	7	7%	10.3	10	9%	7.2	7	9%	5.1	5	5%
55-64	*	2	2%	*	3	3%	*	4	5%	*	4	4%
65+	2.5	5	5%	*	2	2%	*	-	-	*	1	1%
Race/Ethnicity												
White, not Hispanic	3.6	39	38%	4.0	43	38%	3.4	37	46%	3.6	39	35%
Black/African American, not Hispanic	39.2	59	57%	42.5	64	57%	24.6	37	46%	37.2	56	51%
Hispanic/Latino	*	1	1%	*	4	4%	*	4	5%	20.5	8	7%
Asian/Pacific Islander	*	1	1%	*	-	-	*	-	-	*	3	3%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	3%	*	1	1%	*	2	3%	*	4	4%
Race/Sex												
White, not Hispanic Males	6.6	35	34%	6.8	36	32%	5.9	31	39%	6.3	33	30%
White, not Hispanic Females	*	4	4%	1.3	7	6%	1.1	6	8%	1.1	6	5%
Black/African American, not Hispanic Males	63.7	46	45%	59.5	43	38%	38.8	28	35%	60.9	44	40%
Black/African American, not Hispanic Females	16.6	13	13%	26.8	21	19%	11.5	9	11%	15.3	12	11%
Hispanic/Latino Males	*	1	1%	*	4	4%	*	4	5%	*	4	4%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	4	4%
Asian/Pacific Islander Males	*	1	1%	*	-	-	*	-	-	*	2	2%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	1	1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	3%	*	1	1%	*	2	3%	*	4	4%
Total	7.8	103		8.4	112		6.0	80		8.3	110	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 7

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	63	77%	55	65%	47	72%	60	72%
Injection drug use (IDU)	1	1%	1	1%	-	-	1	1%
Male-to-male sexual contact and IDU	1	1%	-	-	2	3%	-	-
Heterosexual contact	3	4%	7	8%	4	6%	3	4%
Other/unknown	14	17%	21	25%	12	18%	19	23%
Subtotal	82	100%	84	100%	65	100%	83	100%
Female adult or adolescent								
Injection drug use	1	5%	3	11%	-	-	3	13%
Heterosexual contact	14	74%	19	68%	9	60%	11	46%
Other/unknown	4	21%	6	21%	6	40%	10	42%
Subtotal	19	100%	28	100%	15	100%	24	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	50%	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	3	100%
Subtotal	2	100%	-	-	-	-	3	100%
Total	103		112		80		110	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 7

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	63	61%	52	46%	45	56%	57	52%
Injection drug use (IDU) only	1	1%	1	1%	-	-	1	1%
Heterosexual contact only	17	17%	26	23%	13	16%	14	13%
MSM & IDU	-	-	-	-	2	3%	-	-
IDU & Heterosexual contact	1	1%	3	3%	-	-	3	3%
MSM & Heterosexual contact	-	-	3	3%	2	3%	3	3%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-
Perinatal exposure	1	1%	-	-	-	-	-	-
Other/unknown	19	18%	27	24%	18	23%	32	29%
Total	103		112		80		110	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 7

Year of HIV (not AIDS) Diagnosis

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	54	84%	48	74%	36	75%	59	74%	57	80%
Females	10	16%	17	26%	12	25%	21	26%	14	20%
Age at HIV diagnosis (yr)										
<13	2	3%	-	-	-	-	3	4%	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	8	13%	4	6%	2	4%	7	9%	6	8%
20-24	20	31%	16	25%	13	27%	15	19%	23	32%
25-29	11	17%	7	11%	14	29%	17	21%	11	15%
30-34	2	3%	12	18%	3	6%	7	9%	4	6%
35-39	6	9%	6	9%	8	17%	13	16%	8	11%
40-44	6	9%	8	12%	-	-	8	10%	8	11%
45-49	3	5%	3	5%	3	6%	5	6%	3	4%
50-54	3	5%	6	9%	3	6%	3	4%	3	4%
55-64	1	2%	2	3%	2	4%	1	1%	2	3%
65+	2	3%	1	2%	-	-	1	1%	2	3%
Race/Ethnicity										
White, not Hispanic	21	33%	27	42%	19	40%	30	38%	30	42%
Black/African American, not Hispanic	40	63%	35	54%	26	54%	39	49%	35	49%
Hispanic/Latino	1	2%	2	3%	1	2%	5	6%	4	6%
Asian/Pacific Islander	1	2%	-	-	-	-	3	4%	1	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	2	4%	3	4%	1	1%
Race/Sex										
White, not Hispanic Males	20	31%	22	34%	15	31%	24	30%	28	39%
White, not Hispanic Females	1	2%	5	8%	4	8%	6	8%	2	3%
Black/African American, not Hispanic Males	32	50%	23	35%	18	38%	29	36%	24	34%
Black/African American, not Hispanic Females	8	13%	12	18%	8	17%	10	13%	11	15%
Hispanic/Latino Males	1	2%	2	3%	1	2%	2	3%	3	4%
Hispanic/Latina Females	-	-	-	-	-	-	3	4%	1	1%
Asian/Pacific Islander Males	1	2%	-	-	-	-	2	3%	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	1	1%	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	2	4%	3	4%	1	1%
Total	64		65		48		80		71	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	40	77%	30	63%	27	75%	42	74%	46	81%
Injection drug use (IDU)	-	-	1	2%	-	-	1	2%	-	-
Male-to-male sexual contact and IDU	1	2%	-	-	1	3%	-	-	1	2%
Heterosexual contact	2	4%	3	6%	2	6%	2	4%	-	-
Other/unknown	9	17%	14	29%	6	17%	12	21%	10	18%
Subtotal	52	100%	48	100%	36	100%	57	100%	57	100%
Female adult or adolescent										
Injection drug use	1	10%	3	18%	-	-	3	15%	-	-
Heterosexual contact	6	60%	9	53%	6	50%	7	35%	8	62%
Other/unknown	3	30%	5	29%	6	50%	10	50%	5	38%
Subtotal	10	100%	17	100%	12	100%	20	100%	13	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	50%	-	-	-	-	-	-	1	100%
Other/unknown	1	50%	-	-	-	-	3	100%	-	-
Subtotal	2	100%	-	-	-	-	3	100%	1	100%
Total	64		65		48		80		71	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	40	63%	27	42%	26	54%	39	49%	46	65%
Injection drug use (IDU) only	-	-	1	2%	-	-	1	1%	-	-
Heterosexual contact only	8	13%	12	18%	8	17%	9	11%	8	11%
MSM & IDU	-	-	-	-	1	2%	-	-	1	1%
IDU & Heterosexual contact	1	2%	3	5%	-	-	3	4%	-	-
MSM & Heterosexual contact	-	-	3	5%	1	2%	3	4%	-	-
MSM & IDU & Heterosexual contact	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	-	-	-	-	-	-	1	1%
Other/unknown	13	20%	19	29%	12	25%	25	31%	15	21%
Total	64		65		48		80		71	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 7

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	36	75%	47	81%	44	92%	37	80%	33	73%	1,533	85%
Females	12	25%	11	19%	4	8%	9	20%	12	27%	278	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	2%	-	-	23	1%
13-14	-	-	-	-	-	-	-	-	-	-	3	<1%
15-19	3	6%	4	7%	2	4%	1	2%	3	7%	40	2%
20-24	4	8%	11	19%	4	8%	6	13%	7	16%	162	9%
25-29	9	19%	7	12%	9	19%	11	24%	6	13%	317	18%
30-34	10	21%	9	16%	8	17%	3	7%	5	11%	336	19%
35-39	4	8%	7	12%	7	15%	8	17%	8	18%	325	18%
40-44	4	8%	4	7%	4	8%	7	15%	5	11%	249	14%
45-49	6	13%	8	14%	7	15%	3	7%	3	7%	172	9%
50-54	4	8%	5	9%	5	10%	4	9%	5	11%	98	5%
55-64	1	2%	2	3%	2	4%	2	4%	2	4%	62	3%
65+	3	6%	1	2%	-	-	-	-	1	2%	24	1%
Race/Ethnicity												
White, not Hispanic	23	48%	27	47%	24	50%	14	30%	23	51%	1,081	60%
Black/African American, not Hispanic	24	50%	25	43%	19	40%	27	59%	19	42%	675	37%
Hispanic/Latino	-	-	2	3%	5	10%	4	9%	-	-	31	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	2%	5	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	1	2%	4	7%	-	-	1	2%	2	4%	18	1%
Race/Sex												
White, not Hispanic Males	20	42%	23	40%	22	46%	13	28%	18	40%	980	54%
White, not Hispanic Females	3	6%	4	7%	2	4%	1	2%	5	11%	101	6%
Black/African American, not Hispanic Males	15	31%	20	34%	17	35%	20	43%	14	31%	507	28%
Black/African American, not Hispanic Females	9	19%	5	9%	2	4%	7	15%	5	11%	168	9%
Hispanic/Latino Males	-	-	2	3%	5	10%	3	7%	-	-	29	2%
Hispanic/Latina Females	-	-	-	-	-	-	1	2%	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	1	2%	4	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	4	7%	-	-	1	2%	2	4%	18	1%
Total	48		58		48		46		45		1,811	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	28	78%	33	70%	29	66%	29	78%	21	64%	1,095	72%
Injection drug use (IDU)	1	3%	-	-	1	2%	-	-	2	6%	62	4%
Male-to-male sexual contact and IDU	2	6%	1	2%	1	2%	-	-	-	-	92	6%
Heterosexual contact	1	3%	5	11%	4	9%	1	3%	3	9%	86	6%
Other/unknown	4	11%	8	17%	9	20%	7	19%	7	21%	184	12%
Subtotal	36	100%	47	100%	44	100%	37	100%	33	100%	1,519	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	1	13%	-	-	43	16%
Heterosexual contact	11	92%	11	100%	4	100%	6	75%	6	50%	196	73%
Other/unknown	1	8%	-	-	-	-	1	13%	6	50%	30	11%
Subtotal	12	100%	11	100%	4	100%	8	100%	12	100%	269	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	1	100%	-	-	16	70%
Other/unknown	-	-	-	-	-	-	-	-	-	-	7	30%
Subtotal	-	-	-	-	-	-	1	100%	-	-	23	100%
Total	48		58		48		46		45		1,811	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	28	58%	33	57%	28	58%	28	61%	21	47%	1,048	58%
Injection drug use (IDU) only	1	2%	-	-	1	2%	1	2%	2	4%	64	4%
Heterosexual contact only	12	25%	16	28%	8	17%	7	15%	9	20%	282	16%
MSM & IDU	1	2%	1	2%	1	2%	-	-	-	-	77	4%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	40	2%
MSM & Heterosexual contact	-	-	-	-	1	2%	1	2%	-	-	47	3%
MSM & IDU & Heterosexual contact	1	2%	-	-	-	-	-	-	-	-	15	1%
Perinatal exposure	-	-	-	-	-	-	1	2%	-	-	16	1%
Other/unknown	5	10%	8	14%	9	19%	8	17%	13	29%	222	12%
Total	48		58		48		46		45		1,811	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 7

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 7

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	234.5	1,521	80%	677	76%	844	83%
Females	57.7	391	20%	214	24%	177	17%
Age at end of year							
<13	3.1	7	<1%	7	1%	-	-
13-14	*	1	<1%	1	<1%	-	-
15-19	14.2	13	1%	9	1%	4	<1%
20-24	92.3	85	4%	64	7%	21	2%
25-29	191.7	151	8%	104	12%	47	5%
30-34	194.7	159	8%	88	10%	71	7%
35-39	244.2	190	10%	99	11%	91	9%
40-44	259.1	220	12%	111	12%	109	11%
45-49	320.4	283	15%	115	13%	168	16%
50-54	362.3	353	18%	127	14%	226	22%
55-64	200.0	347	18%	130	15%	217	21%
65+	51.1	103	5%	36	4%	67	7%
Race/Ethnicity							
White, not Hispanic	87.9	945	49%	418	47%	527	52%
Black/African American, not Hispanic	551.9	831	43%	408	46%	423	41%
Hispanic/Latino	195.0	76	4%	35	4%	41	4%
Asian/Pacific Islander	46.7	16	1%	9	1%	7	1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	43	2%	21	2%	22	2%
Race/Sex							
White, not Hispanic Males	155.9	821	43%	354	40%	467	46%
White, not Hispanic Females	22.6	124	6%	64	7%	60	6%
Black/African American, not Hispanic Males	819.4	592	31%	273	31%	319	31%
Black/African American, not Hispanic Females	305.1	239	13%	135	15%	104	10%
Hispanic/Latino Males	318.1	63	3%	27	3%	36	4%
Hispanic/Latina Females	67.8	13	1%	8	1%	5	<1%
Asian/Pacific Islander Males	86.6	14	1%	8	1%	6	1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	43	2%	21	2%	22	2%
Total	144.1	1,912		891		1,021	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,085	72%	462	69%	623	74%
Injection drug use (IDU)	40	3%	18	3%	22	3%
Male-to-male sexual contact and IDU	54	4%	17	3%	37	4%
Heterosexual contact	86	6%	32	5%	54	6%
Other/unknown	242	16%	139	21%	103	12%
Subtotal	1,507	100%	668	100%	839	100%
Female adult or adolescent						
Injection drug use	38	10%	19	9%	19	11%
Heterosexual contact	266	70%	132	64%	134	78%
Other/unknown	75	20%	56	27%	19	11%
Subtotal	379	100%	207	100%	172	100%
Child (<13 yrs at diagnosis)						
Perinatal	17	65%	10	63%	7	70%
Other/unknown	9	35%	6	38%	3	30%
Subtotal	26	100%	16	100%	10	100%
Total	1,912		891		1,021	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	1,039	54%	447	50%	592	58%
Injection drug use (IDU) only	37	2%	16	2%	21	2%
Heterosexual contact only	352	18%	164	18%	188	18%
MSM & IDU	47	2%	15	2%	32	3%
IDU & Heterosexual contact	41	2%	21	2%	20	2%
MSM & Heterosexual contact	46	2%	15	2%	31	3%
MSM & IDU & Heterosexual contact	7	<1%	2	<1%	5	<1%
Perinatal exposure	18	1%	11	1%	7	1%
Other/unknown	325	17%	200	22%	125	12%
Total	1,912		891		1,021	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	642	79%	372	64%	44	70%	7	54%	-	-	20	67%
Injection drug use (IDU)	9	1%	25	4%	5	8%	-	-	1	100%	-	-
Male-to-male sexual contact and IDU	31	4%	20	3%	2	3%	-	-	-	-	1	3%
Heterosexual contact	22	3%	57	10%	5	8%	2	15%	-	-	-	-
Other/unknown	112	14%	110	19%	7	11%	4	31%	-	-	9	30%
Subtotal	816	100%	584	100%	63	100%	13	100%	1	100%	30	100%
Female adult or adolescent												
Injection drug use	20	17%	16	7%	1	8%	1	100%	-	-	-	-
Heterosexual contact	77	64%	170	74%	8	62%	-	-	-	-	11	85%
Other/unknown	24	20%	45	19%	4	31%	-	-	-	-	2	15%
Subtotal	121	100%	231	100%	13	100%	1	100%	-	-	13	100%
Child (<13 yrs at diagnosis)												
Perinatal	6	75%	11	69%	-	-	-	-	-	-	-	-
Other/unknown	2	25%	5	31%	-	-	2	100%	-	-	-	-
Subtotal	8	100%	16	100%	-	-	2	100%	-	-	-	-
Total	945		831		76		16		1		43	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 7

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	627	66%	348	42%	39	51%	6	38%	-	-	19	44%
Injection drug use (IDU) only	12	1%	21	3%	2	3%	1	6%	1	100%	-	-
Heterosexual contact only	99	10%	227	27%	13	17%	2	13%	-	-	11	26%
MSM & IDU	27	3%	18	2%	1	1%	-	-	-	-	1	2%
IDU & Heterosexual contact	17	2%	20	2%	4	5%	-	-	-	-	-	-
MSM & Heterosexual contact	15	2%	24	3%	5	7%	1	6%	-	-	1	2%
MSM & IDU & Heterosexual contact	4	0%	2	0%	1	1%	-	-	-	-	-	-
Perinatal exposure	6	1%	12	1%	-	-	-	-	-	-	-	-
Other/unknown	138	15%	159	19%	11	14%	6	38%	-	-	11	26%
Total	945		831		76		16		1		43	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 7

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	1,221	80%	1,307	80%	1,368	80%	1,457	80%	1,521	80%
Females	314	20%	335	20%	348	20%	372	20%	391	20%
Age at end of year										
<13	6	<1%	5	<1%	4	<1%	7	<1%	7	<1%
13-14	2	<1%	1	<1%	2	<1%	1	<1%	1	<1%
15-19	21	1%	20	1%	16	1%	18	1%	13	1%
20-24	54	4%	81	5%	84	5%	88	5%	85	4%
25-29	104	7%	111	7%	121	7%	140	8%	151	8%
30-34	144	9%	142	9%	144	8%	150	8%	159	8%
35-39	164	11%	149	9%	167	10%	186	10%	190	10%
40-44	233	15%	237	14%	218	13%	218	12%	220	12%
45-49	325	21%	339	21%	320	19%	295	16%	283	15%
50-54	223	15%	259	16%	290	17%	336	18%	353	18%
55-64	217	14%	243	15%	281	16%	306	17%	347	18%
65+	42	3%	55	3%	69	4%	84	5%	103	5%
Race/Ethnicity										
White, not Hispanic	798	52%	838	51%	869	51%	910	50%	945	49%
Black/African American, not Hispanic	637	41%	698	43%	733	43%	791	43%	831	43%
Hispanic/Latino	52	3%	57	3%	63	4%	70	4%	76	4%
Asian/Pacific Islander	11	1%	11	1%	11	1%	15	1%	16	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	36	2%	37	2%	39	2%	42	2%	43	2%
Race/Sex										
White, not Hispanic Males	695	45%	730	44%	755	44%	790	43%	821	43%
White, not Hispanic Females	103	7%	108	7%	114	7%	120	7%	124	6%
Black/African American, not Hispanic Males	446	29%	491	30%	519	30%	566	31%	592	31%
Black/African American, not Hispanic Females	191	12%	207	13%	214	12%	225	12%	239	13%
Hispanic/Latino Males	44	3%	49	3%	55	3%	58	3%	63	3%
Hispanic/Latina Females	8	1%	8	<1%	8	<1%	12	1%	13	1%
Asian/Pacific Islander Males	10	1%	10	1%	10	1%	13	1%	14	1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	36	2%	37	2%	39	2%	42	2%	43	2%
Total	1,535		1,642		1,716		1,829		1,912	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 7

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	857	71%	923	71%	966	71%	1,032	72%	1,085	72%
Injection drug use (IDU)	39	3%	39	3%	39	3%	40	3%	40	3%
Male-to-male sexual contact and IDU	53	4%	53	4%	54	4%	54	4%	54	4%
Heterosexual contact	71	6%	78	6%	82	6%	85	6%	86	6%
Other/unknown	189	16%	202	16%	215	16%	232	16%	242	16%
Subtotal	1,209	100%	1,295	100%	1,356	100%	1,443	100%	1,507	100%
Female adult or adolescent										
Injection drug use	32	11%	34	10%	35	10%	38	11%	38	10%
Heterosexual contact	224	74%	239	74%	247	73%	258	71%	266	70%
Other/unknown	48	16%	52	16%	56	17%	65	18%	75	20%
Subtotal	304	100%	325	100%	338	100%	361	100%	379	100%
Child (<13 yrs at diagnosis)										
Perinatal	16	73%	16	73%	16	73%	16	64%	17	65%
Other/unknown	6	27%	6	27%	6	27%	9	36%	9	35%
Subtotal	22	100%	22	100%	22	100%	25	100%	26	100%
Total	1,535		1,642		1,716		1,829		1,912	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 7

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	819	53%	882	54%	924	54%	987	54%	1,039	54%
Injection drug use (IDU) only	35	2%	36	2%	37	2%	37	2%	37	2%
Heterosexual contact only	295	19%	317	19%	329	19%	343	19%	352	18%
MSM & IDU	46	3%	46	3%	47	3%	47	3%	47	2%
IDU & Heterosexual contact	36	2%	37	2%	37	2%	41	2%	41	2%
MSM & Heterosexual contact	38	2%	41	2%	42	2%	45	2%	46	2%
MSM & IDU & Heterosexual contact	7	<1%	7	<1%	7	<1%	7	<1%	7	<1%
Perinatal exposure	17	1%	17	1%	17	1%	17	1%	18	1%
Other/unknown	242	16%	259	16%	276	16%	305	17%	325	17%
Total	1,535		1,642		1,716		1,829		1,912	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 8A

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 8A

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	3.6	16	70%	10	67%	3	75%	3	75%	
Females	1.5	7	30%	5	33%	1	25%	1	25%	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	10.5	6	26%	5	33%	1	25%	-	-	
25-29	12.4	6	26%	5	33%	1	25%	-	-	
30-34	*	2	9%	-	-	1	25%	1	25%	
35-39	*	4	17%	2	13%	-	-	2	50%	
40-44	*	1	4%	1	7%	-	-	-	-	
45-49	*	2	9%	1	7%	1	25%	-	-	
50-54	*	-	-	-	-	-	-	-	-	
55-64	*	2	9%	1	7%	-	-	1	25%	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity										
White, not Hispanic	2.0	16	70%	10	67%	3	75%	3	75%	
Black/African American, not Hispanic	19.2	6	26%	4	27%	1	25%	1	25%	
Hispanic/Latino	*	1	4%	1	7%	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	-	-	-	-	-	-	-	-	
Race/Sex										
White, not Hispanic Males	3.0	12	52%	8	53%	2	50%	2	50%	
White, not Hispanic Females	*	4	17%	2	13%	1	25%	1	25%	
Black/African American, not Hispanic Males	*	4	17%	2	13%	1	25%	1	25%	
Black/African American, not Hispanic Females	*	2	9%	2	13%	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	1	4%	1	7%	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	-	-	-	-	-	-	-	-	
Total	2.6	23		15		4		4		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	8	50%	5	50%	2	67%	1	33%
Injection drug use (IDU)	1	6%	1	10%	-	-	-	-
Male-to-male sexual contact and IDU	2	13%	2	20%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	5	31%	2	20%	1	33%	2	67%
Subtotal	16	100%	10	100%	3	100%	3	100%
Female adult or adolescent								
Injection drug use	1	14%	1	20%	-	-	-	-
Heterosexual contact	6	86%	4	80%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	7	100%	5	100%	1	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	23		15		4		4	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	30%	5	33%	1	25%	1	25%
Injection drug use (IDU) only	2	9%	2	13%	-	-	-	-
Heterosexual contact only	6	26%	4	27%	1	25%	1	25%
MSM & IDU	2	9%	2	13%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	4%	-	-	1	25%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	5	22%	2	13%	1	25%	2	50%
Total	23		15		4		4	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	50%	2	50%	-	-	-	-	-	-	-	-
Injection drug use (IDU)	1	8%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	17%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	25%	2	50%	-	-	-	-	-	-	-	-
Subtotal	12	100%	4	100%	-	-	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	1	100%	-	-	-	-	-	-
Heterosexual contact	4	100%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	2	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	16		6		1		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	6	38%	1	17%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	1	6%	-	-	1	100%	-	-	-	-	-	-
Heterosexual contact only	4	25%	2	33%	-	-	-	-	-	-	-	-
MSM & IDU	2	13%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	1	17%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	19%	2	33%	-	-	-	-	-	-	-	-
Total	16		6		1		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 8A

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	6.7	30	91%	4.9	22	85%	5.8	26	90%	7.4	33	83%
Females	*	3	9%	*	4	15%	*	3	10%	1.5	7	18%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	3%	*	1	4%	*	1	3%	*	1	3%
20-24	10.5	6	18%	8.8	5	19%	*	2	7%	15.8	9	23%
25-29	12.4	6	18%	*	1	4%	*	3	10%	*	4	10%
30-34	*	3	9%	*	4	15%	*	3	10%	*	4	10%
35-39	*	4	12%	*	4	15%	*	3	10%	*	4	10%
40-44	*	2	6%	9.9	6	23%	*	3	10%	8.3	5	13%
45-49	8.2	5	15%	*	4	15%	9.8	6	21%	*	1	3%
50-54	*	3	9%	*	-	-	7.5	5	17%	*	3	8%
55-64	*	3	9%	*	1	4%	*	3	10%	5.8	7	18%
65+	*	-	-	*	-	-	*	-	-	*	2	5%
Race/Ethnicity												
White, not Hispanic	3.1	25	76%	2.2	18	69%	2.2	18	62%	3.1	25	63%
Black/African American, not Hispanic	*	2	6%	16.0	5	19%	22.4	7	24%	35.2	11	28%
Hispanic/Latino	*	2	6%	*	2	8%	*	-	-	*	1	3%
Asian/Pacific Islander	*	-	-	*	1	4%	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	12%	*	-	-	*	4	14%	*	3	8%
Race/Sex												
White, not Hispanic Males	5.7	23	70%	4.0	16	62%	4.5	18	62%	5.2	21	53%
White, not Hispanic Females	*	2	6%	*	2	8%	*	-	-	*	4	10%
Black/African American, not Hispanic Males	*	2	6%	*	3	12%	*	4	14%	52.0	9	23%
Black/African American, not Hispanic Females	*	-	-	*	2	8%	*	3	10%	*	2	5%
Hispanic/Latino Males	*	1	3%	*	2	8%	*	-	-	*	1	3%
Hispanic/Latina Females	*	1	3%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	1	4%	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	12%	*	-	-	*	4	14%	*	3	8%
Total	3.7	33		2.9	26		3.2	29		4.4	40	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 8A

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	16	53%	13	59%	18	69%	22	67%
Injection drug use (IDU)	-	-	1	5%	2	8%	2	6%
Male-to-male sexual contact and IDU	1	3%	-	-	-	-	-	-
Heterosexual contact	1	3%	-	-	1	4%	1	3%
Other/unknown	12	40%	8	36%	5	19%	8	24%
Subtotal	30	100%	22	100%	26	100%	33	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	100%	4	100%	2	67%	7	100%
Other/unknown	-	-	-	-	1	33%	-	-
Subtotal	3	100%	4	100%	3	100%	7	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	33		26		29		40	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 8A

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	48%	12	46%	11	38%	19	48%
Injection drug use (IDU) only	-	-	-	-	1	3%	-	-
Heterosexual contact only	4	12%	4	15%	3	10%	8	20%
MSM & IDU	1	3%	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	4%	1	3%	2	5%
MSM & Heterosexual contact	-	-	1	4%	7	24%	3	8%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	12	36%	8	31%	6	21%	8	20%
Total	33		26		29		40	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 8A

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	19	100%	15	83%	14	82%	18	75%	10	67%
Females	-	-	3	17%	3	18%	6	25%	5	33%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	1	6%	-	-	1	4%	-	-
20-24	6	32%	3	17%	2	12%	7	29%	5	33%
25-29	3	16%	1	6%	3	18%	1	4%	5	33%
30-34	1	5%	3	17%	3	18%	4	17%	-	-
35-39	2	11%	2	11%	1	6%	3	13%	2	13%
40-44	2	11%	3	17%	1	6%	2	8%	1	7%
45-49	2	11%	4	22%	2	12%	-	-	1	7%
50-54	1	5%	-	-	4	24%	2	8%	-	-
55-64	2	11%	1	6%	1	6%	3	13%	1	7%
65+	-	-	-	-	-	-	1	4%	-	-
Race/Ethnicity										
White, not Hispanic	13	68%	13	72%	9	53%	15	63%	10	67%
Black/African American, not Hispanic	2	11%	2	11%	5	29%	7	29%	4	27%
Hispanic/Latino	1	5%	2	11%	-	-	-	-	1	7%
Asian/Pacific Islander	-	-	1	6%	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	3	16%	-	-	3	18%	2	8%	-	-
Race/Sex										
White, not Hispanic Males	13	68%	11	61%	9	53%	11	46%	8	53%
White, not Hispanic Females	-	-	2	11%	-	-	4	17%	2	13%
Black/African American, not Hispanic Males	2	11%	1	6%	2	12%	6	25%	2	13%
Black/African American, not Hispanic Females	-	-	1	6%	3	18%	1	4%	2	13%
Hispanic/Latino Males	1	5%	2	11%	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-	-	-	1	7%
Asian/Pacific Islander Males	-	-	1	6%	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	16%	-	-	3	18%	2	8%	-	-
Total	19		18		17		24		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	12	63%	8	53%	11	79%	13	72%	5	50%
Injection drug use (IDU)	-	-	1	7%	2	14%	1	6%	1	10%
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-	2	20%
Heterosexual contact	-	-	-	-	1	7%	1	6%	-	-
Other/unknown	6	32%	6	40%	-	-	3	17%	2	20%
Subtotal	19	100%	15	100%	14	100%	18	100%	10	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	1	20%
Heterosexual contact	-	-	3	100%	2	67%	6	100%	4	80%
Other/unknown	-	-	-	-	1	33%	-	-	-	-
Subtotal	-	-	3	100%	3	100%	6	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	19		18		17		24		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	63%	7	39%	6	35%	11	46%	5	33%
Injection drug use (IDU) only	-	-	-	-	1	6%	-	-	2	13%
Heterosexual contact only	-	-	3	17%	3	18%	7	29%	4	27%
MSM & IDU	1	5%	-	-	-	-	-	-	2	13%
IDU & Heterosexual contact	-	-	1	6%	1	6%	1	4%	-	-
MSM & Heterosexual contact	-	-	1	6%	5	29%	2	8%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	32%	6	33%	1	6%	3	13%	2	13%
Total	19		18		17		24		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 8A

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	13	87%	12	86%	17	85%	23	100%	8	67%	468	85%
Females	2	13%	2	14%	3	15%	-	-	4	33%	82	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	4%	-	-	7	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	-	-	1	7%	3	15%	-	-	-	-	10	2%
20-24	-	-	1	7%	-	-	2	9%	1	8%	38	7%
25-29	4	27%	3	21%	1	5%	4	17%	1	8%	86	16%
30-34	3	20%	2	14%	1	5%	2	9%	2	17%	104	19%
35-39	3	20%	3	21%	4	20%	2	9%	3	25%	102	19%
40-44	1	7%	2	14%	3	15%	6	26%	-	-	73	13%
45-49	2	13%	-	-	4	20%	2	9%	2	17%	59	11%
50-54	2	13%	1	7%	2	10%	1	4%	-	-	34	6%
55-64	-	-	1	7%	2	10%	3	13%	2	17%	27	5%
65+	-	-	-	-	-	-	-	-	1	8%	8	1%
Race/Ethnicity												
White, not Hispanic	13	87%	9	64%	15	75%	16	70%	8	67%	453	82%
Black/African American, not Hispanic	-	-	2	14%	4	20%	4	17%	4	33%	68	12%
Hispanic/Latino	-	-	2	14%	-	-	1	4%	-	-	16	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	2	13%	1	7%	1	5%	2	9%	-	-	12	2%
Race/Sex												
White, not Hispanic Males	11	73%	9	64%	14	70%	16	70%	5	42%	402	73%
White, not Hispanic Females	2	13%	-	-	1	5%	-	-	3	25%	51	9%
Black/African American, not Hispanic Males	-	-	1	7%	2	10%	4	17%	3	25%	43	8%
Black/African American, not Hispanic Females	-	-	1	7%	2	10%	-	-	1	8%	25	5%
Hispanic/Latino Males	-	-	1	7%	-	-	1	4%	-	-	12	2%
Hispanic/Latina Females	-	-	1	7%	-	-	-	-	-	-	4	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	13%	1	7%	1	5%	2	9%	-	-	12	2%
Total	15		14		20		23		12		550	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	8	62%	9	75%	10	59%	15	68%	3	38%	311	67%
Injection drug use (IDU)	-	-	-	-	-	-	2	9%	-	-	21	5%
Male-to-male sexual contact and IDU	-	-	-	-	1	6%	1	5%	-	-	20	4%
Heterosexual contact	1	8%	-	-	-	-	-	-	-	-	24	5%
Other/unknown	4	31%	3	25%	6	35%	4	18%	5	63%	89	19%
Subtotal	13	100%	12	100%	17	100%	22	100%	8	100%	465	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	14	18%
Heterosexual contact	2	100%	2	100%	3	100%	-	-	4	100%	56	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	7	9%
Subtotal	2	100%	2	100%	3	100%	-	-	4	100%	77	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	50%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	4	50%
Subtotal	-	-	-	-	-	-	1	100%	-	-	8	100%
Total	15		14		20		23		12		550	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	8	53%	9	64%	8	40%	13	57%	2	17%	289	53%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	16	3%
Heterosexual contact only	3	20%	2	14%	3	15%	-	-	4	33%	80	15%
MSM & IDU	-	-	-	-	-	-	1	4%	-	-	16	3%
IDU & Heterosexual contact	-	-	-	-	-	-	2	9%	-	-	19	3%
MSM & Heterosexual contact	-	-	-	-	2	10%	2	9%	1	8%	22	4%
MSM & IDU & Heterosexual contact	-	-	-	-	1	5%	-	-	-	-	4	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	4	1%
Other/unknown	4	27%	3	21%	6	30%	5	22%	5	42%	100	18%
Total	15		14		20		23		12		550	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 8A

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 8A

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	112.1	503	77%	220	75%	283	80%
Females	32.5	147	23%	75	25%	72	20%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	3	<1%	1	<1%	2	1%
20-24	29.9	17	3%	12	4%	5	1%
25-29	76.2	37	6%	25	8%	12	3%
30-34	90.0	48	7%	29	10%	19	5%
35-39	120.8	67	10%	29	10%	38	11%
40-44	153.8	93	14%	36	12%	57	16%
45-49	160.8	98	15%	38	13%	60	17%
50-54	191.9	128	20%	59	20%	69	19%
55-64	99.3	119	18%	49	17%	70	20%
65+	29.8	40	6%	17	6%	23	6%
Race/Ethnicity							
White, not Hispanic	58.8	477	73%	216	73%	261	74%
Black/African American, not Hispanic	380.7	119	18%	58	20%	61	17%
Hispanic/Latino	118.3	29	4%	11	4%	18	5%
Asian/Pacific Islander	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	23	4%	9	3%	14	4%
Race/Sex							
White, not Hispanic Males	97.0	390	60%	173	59%	217	61%
White, not Hispanic Females	21.2	87	13%	43	15%	44	12%
Black/African American, not Hispanic Males	404.5	70	11%	32	11%	38	11%
Black/African American, not Hispanic Females	351.1	49	8%	26	9%	23	6%
Hispanic/Latino Males	162.0	21	3%	6	2%	15	4%
Hispanic/Latina Females	69.3	8	1%	5	2%	3	1%
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	23	4%	9	3%	14	4%
Total	72.2	650		295		355	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	326	65%	139	63%	187	67%
Injection drug use (IDU)	18	4%	6	3%	12	4%
Male-to-male sexual contact and IDU	29	6%	7	3%	22	8%
Heterosexual contact	37	7%	11	5%	26	9%
Other/unknown	90	18%	57	26%	33	12%
Subtotal	500	100%	220	100%	280	100%
Female adult or adolescent						
Injection drug use	24	17%	10	13%	14	20%
Heterosexual contact	99	68%	51	68%	48	69%
Other/unknown	22	15%	14	19%	8	11%
Subtotal	145	100%	75	100%	70	100%
Child (<13 yrs at diagnosis)						
Perinatal	5	100%	-	-	5	100%
Other/unknown	-	-	-	-	-	-
Subtotal	5	100%	-	-	5	100%
Total	650		295		355	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	301	46%	128	43%	173	49%
Injection drug use (IDU) only	20	3%	9	3%	11	3%
Heterosexual contact only	136	21%	62	21%	74	21%
MSM & IDU	23	4%	5	2%	18	5%
IDU & Heterosexual contact	22	3%	7	2%	15	4%
MSM & Heterosexual contact	25	4%	11	4%	14	4%
MSM & IDU & Heterosexual contact	6	1%	2	1%	4	1%
Perinatal exposure	5	1%	-	-	5	1%
Other/unknown	112	17%	71	24%	41	12%
Total	650		295		355	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 8A

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	272	70%	33	48%	7	37%	-	-	-	-	14	74%
Injection drug use (IDU)	9	2%	6	9%	3	16%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	21	5%	3	4%	3	16%	-	-	-	-	2	11%
Heterosexual contact	19	5%	13	19%	4	21%	1	50%	-	-	-	-
Other/unknown	69	18%	14	20%	2	11%	1	50%	-	-	3	16%
Subtotal	390	100%	69	100%	19	100%	2	100%	-	-	19	100%
Female adult or adolescent												
Injection drug use	14	16%	9	18%	1	13%	-	-	-	-	-	-
Heterosexual contact	56	66%	35	71%	6	75%	-	-	-	-	2	67%
Other/unknown	15	18%	5	10%	1	13%	-	-	-	-	1	33%
Subtotal	85	100%	49	100%	8	100%	-	-	-	-	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	1	100%	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	1	100%	2	100%	-	-	-	-	-	-
Total	477		119		29		2		0		22	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 8A

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	253	53%	29	24%	7	24%	-	-	-	-	12	52%
Injection drug use (IDU) only	12	3%	6	5%	2	7%	-	-	-	-	-	-
Heterosexual contact only	75	16%	48	40%	10	34%	1	50%	-	-	2	9%
MSM & IDU	17	4%	3	3%	2	7%	-	-	-	-	1	4%
IDU & Heterosexual contact	11	2%	9	8%	2	7%	-	-	-	-	-	-
MSM & Heterosexual contact	19	4%	4	3%	-	-	-	-	-	-	2	9%
MSM & IDU & Heterosexual contact	4	1%	-	-	1	3%	-	-	-	-	1	4%
Perinatal exposure	2	0%	1	1%	2	7%	-	-	-	-	-	-
Other/unknown	84	18%	19	16%	3	10%	1	50%	-	-	5	22%
Total	477		119		29		2		0		23	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 8A

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	417	77%	443	77%	463	78%	487	78%	503	77%
Females	126	23%	129	23%	134	22%	141	22%	147	23%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	1	<1%	1	<1%	-	-	-	-	-	-
15-19	3	1%	3	1%	4	1%	4	1%	3	<1%
20-24	16	3%	16	3%	16	3%	18	3%	17	3%
25-29	31	6%	32	6%	32	5%	36	6%	37	6%
30-34	48	9%	52	9%	50	8%	45	7%	48	7%
35-39	66	12%	66	12%	65	11%	66	11%	67	10%
40-44	91	17%	88	15%	95	16%	93	15%	93	14%
45-49	114	21%	126	22%	116	19%	118	19%	98	15%
50-54	82	15%	84	15%	104	17%	106	17%	128	20%
55-64	74	14%	82	14%	88	15%	111	18%	119	18%
65+	17	3%	22	4%	27	5%	31	5%	40	6%
Race/Ethnicity										
White, not Hispanic	405	75%	424	74%	441	74%	460	73%	477	73%
Black/African American, not Hispanic	96	18%	103	18%	107	18%	115	18%	119	18%
Hispanic/Latino	25	5%	26	5%	27	5%	28	4%	29	4%
Asian/Pacific Islander	1	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	16	3%	17	3%	20	3%	23	4%	23	4%
Race/Sex										
White, not Hispanic Males	328	60%	346	60%	361	60%	376	60%	390	60%
White, not Hispanic Females	77	14%	78	14%	80	13%	84	13%	87	13%
Black/African American, not Hispanic Males	56	10%	61	11%	62	10%	68	11%	70	11%
Black/African American, not Hispanic Females	40	7%	42	7%	45	8%	47	7%	49	8%
Hispanic/Latino Males	18	3%	19	3%	20	3%	21	3%	21	3%
Hispanic/Latina Females	7	1%	7	1%	7	1%	7	1%	8	1%
Asian/Pacific Islander Males	1	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	16	3%	17	3%	20	3%	23	4%	23	4%
Total	543		572		597		628		650	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 8A

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	266	64%	285	65%	301	65%	316	65%	326	65%
Injection drug use (IDU)	16	4%	17	4%	17	4%	18	4%	18	4%
Male-to-male sexual contact and IDU	27	7%	28	6%	28	6%	28	6%	29	6%
Heterosexual contact	33	8%	34	8%	35	8%	37	8%	37	7%
Other/unknown	72	17%	76	17%	79	17%	85	18%	90	18%
Subtotal	414	100%	440	100%	460	100%	484	100%	500	100%
Female adult or adolescent										
Injection drug use	21	17%	21	17%	22	17%	23	17%	24	17%
Heterosexual contact	83	67%	86	68%	88	67%	94	68%	99	68%
Other/unknown	20	16%	20	16%	22	17%	22	16%	22	15%
Subtotal	124	100%	127	100%	132	100%	139	100%	145	100%
Child (<13 yrs at diagnosis)										
Perinatal	5	100%	5	100%	5	100%	5	100%	5	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	543		572		597		628		650	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 8A

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	249	46%	266	47%	280	47%	292	46%	301	46%
Injection drug use (IDU) only	18	3%	19	3%	19	3%	19	3%	20	3%
Heterosexual contact only	116	21%	120	21%	123	21%	131	21%	136	21%
MSM & IDU	22	4%	22	4%	22	4%	22	4%	23	4%
IDU & Heterosexual contact	19	3%	19	3%	20	3%	22	4%	22	3%
MSM & Heterosexual contact	17	3%	19	3%	21	4%	24	4%	25	4%
MSM & IDU & Heterosexual contact	5	1%	6	1%	6	1%	6	1%	6	1%
Perinatal exposure	5	1%	5	1%	5	1%	5	1%	5	1%
Other/unknown	92	17%	96	17%	101	17%	107	17%	112	17%
Total	543		572		597		628		650	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 8B

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 8B

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	2.9	16	80%	13	81%	1	50%	2	100%
Females	*	4	20%	3	19%	1	50%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	1	5%	1	6%	-	-	-	-
20-24	*	4	20%	4	25%	-	-	-	-
25-29	*	4	20%	3	19%	-	-	1	50%
30-34	*	1	5%	1	6%	-	-	-	-
35-39	*	4	20%	4	25%	-	-	-	-
40-44	*	3	15%	2	13%	1	50%	-	-
45-49	*	1	5%	-	-	-	-	1	50%
50-54	*	1	5%	1	6%	-	-	-	-
55-64	*	-	-	-	-	-	-	-	-
65+	*	1	5%	-	-	1	50%	-	-
Race/Ethnicity									
White, not Hispanic	0.8	8	40%	6	38%	1	50%	1	50%
Black/African American, not Hispanic	*	3	15%	2	13%	-	-	1	50%
Hispanic/Latino	15.4	7	35%	7	44%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	2	10%	1	6%	1	50%	-	-
Race/Sex									
White, not Hispanic Males	1.0	5	25%	4	25%	-	-	1	50%
White, not Hispanic Females	*	3	15%	2	13%	1	50%	-	-
Black/African American, not Hispanic Males	*	2	10%	1	6%	-	-	1	50%
Black/African American, not Hispanic Females	*	1	5%	1	6%	-	-	-	-
Hispanic/Latino Males	30.1	7	35%	7	44%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	2	10%	1	6%	1	50%	-	-
Total	1.8	20		16		2		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	7	44%	7	54%	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	6%	-	-	-	-	1	50%
Other/unknown	8	50%	6	46%	1	100%	1	50%
Subtotal	16	100%	13	100%	1	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	75%	2	67%	1	100%	-	-
Other/unknown	1	25%	1	33%	-	-	-	-
Subtotal	4	100%	3	100%	1	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	20		16		2		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	35%	7	44%	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	20%	2	13%	1	50%	1	50%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	9	45%	7	44%	1	50%	1	50%
Total	20		16		2		2	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	3	60%	-	-	3	43%	-	-	-	-	1	50%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	50%	-	-	-	-	-	-	-	-
Other/unknown	2	40%	1	50%	4	57%	-	-	-	-	1	50%
Subtotal	5	100%	2	100%	7	100%	-	-	-	-	2	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	100%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	3	100%	1	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	8		3		7		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	3	38%	-	-	3	43%	-	-	-	-	1	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	38%	1	33%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	25%	2	67%	4	57%	-	-	-	-	1	50%
Total	8		3		7		0		0		2	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 8B

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	4.1	23	85%	4.9	27	75%	5.9	33	80%	6.3	35	78%
Females	*	4	15%	1.6	9	25%	1.4	8	20%	1.8	10	22%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	1	2%	*	1	2%
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	4%	*	1	3%	*	3	7%	*	-	-
20-24	8.5	6	22%	11.4	8	22%	15.6	11	27%	11.4	8	18%
25-29	8.2	5	19%	9.8	6	17%	9.8	6	15%	8.2	5	11%
30-34	9.4	6	22%	*	2	6%	7.9	5	12%	7.9	5	11%
35-39	7.8	5	19%	7.8	5	14%	*	2	5%	14.0	9	20%
40-44	*	-	-	*	3	8%	*	3	7%	7.0	5	11%
45-49	*	2	7%	*	4	11%	*	-	-	6.7	5	11%
50-54	*	1	4%	*	3	8%	6.0	5	12%	*	1	2%
55-64	*	1	4%	*	3	8%	3.2	5	12%	3.2	5	11%
65+	*	-	-	*	1	3%	*	-	-	*	1	2%
Race/Ethnicity												
White, not Hispanic	1.8	18	67%	2.2	22	61%	2.8	28	68%	2.8	28	62%
Black/African American, not Hispanic	14.3	7	26%	12.2	6	17%	18.4	9	22%	24.5	12	27%
Hispanic/Latino	*	2	7%	11.0	5	14%	*	4	10%	11.0	5	11%
Asian/Pacific Islander	*	-	-	*	1	3%	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	2	6%	*	-	-	*	-	-
Race/Sex												
White, not Hispanic Males	3.3	16	59%	3.1	15	42%	4.9	24	59%	4.7	23	51%
White, not Hispanic Females	*	2	7%	1.4	7	19%	*	4	10%	1.0	5	11%
Black/African American, not Hispanic Males	22.1	6	22%	22.1	6	17%	18.4	5	12%	29.5	8	18%
Black/African American, not Hispanic Females	*	1	4%	*	-	-	*	4	10%	*	4	9%
Hispanic/Latino Males	*	1	4%	*	4	11%	*	4	10%	*	4	9%
Hispanic/Latina Females	*	1	4%	*	1	3%	*	-	-	*	1	2%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	1	3%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	2	6%	*	-	-	*	-	-
Total	2.4	27		3.2	36		3.6	41		4.0	45	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 8B

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	18	78%	16	59%	23	70%	26	74%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	9%	2	7%	1	3%	-	-
Heterosexual contact	-	-	2	7%	-	-	3	9%
Other/unknown	3	13%	7	26%	9	27%	6	17%
Subtotal	23	100%	27	100%	33	100%	35	100%
Female adult or adolescent								
Injection drug use	-	-	3	33%	-	-	-	-
Heterosexual contact	4	100%	4	44%	2	29%	5	56%
Other/unknown	-	-	2	22%	5	71%	4	44%
Subtotal	4	100%	9	100%	7	100%	9	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	1	100%	-	-
Subtotal	-	-	-	-	1	100%	1	100%
Total	27		36		41		45	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 8B

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	17	63%	16	44%	23	56%	25	56%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	15%	6	17%	2	5%	8	18%
MSM & IDU	1	4%	1	3%	1	2%	-	-
IDU & Heterosexual contact	-	-	3	8%	-	-	-	-
MSM & Heterosexual contact	1	4%	-	-	-	-	1	2%
MSM & IDU & Heterosexual contact	1	4%	1	3%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	1	2%
Other/unknown	3	11%	9	25%	15	37%	10	22%
Total	27		36		41		45	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 8B

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	14	70%	22	81%	17	81%	24	75%	13	81%
Females	6	30%	5	19%	4	19%	8	25%	3	19%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	1	5%	1	3%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	5%	1	4%	2	10%	-	-	1	6%
20-24	4	20%	7	26%	6	29%	8	25%	4	25%
25-29	3	15%	3	11%	5	24%	4	13%	3	19%
30-34	4	20%	2	7%	3	14%	4	13%	1	6%
35-39	2	10%	4	15%	-	-	6	19%	4	25%
40-44	1	5%	2	7%	-	-	2	6%	2	13%
45-49	3	15%	3	11%	-	-	4	13%	-	-
50-54	1	5%	1	4%	3	14%	1	3%	1	6%
55-64	1	5%	3	11%	1	5%	2	6%	-	-
65+	-	-	1	4%	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	13	65%	17	63%	14	67%	18	56%	6	38%
Black/African American, not Hispanic	4	20%	4	15%	4	19%	11	34%	2	13%
Hispanic/Latino	3	15%	3	11%	3	14%	3	9%	7	44%
Asian/Pacific Islander	-	-	1	4%	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	7%	-	-	-	-	1	6%
Race/Sex										
White, not Hispanic Males	9	45%	13	48%	12	57%	14	44%	4	25%
White, not Hispanic Females	4	20%	4	15%	2	10%	4	13%	2	13%
Black/African American, not Hispanic Males	3	15%	4	15%	2	10%	8	25%	1	6%
Black/African American, not Hispanic Females	1	5%	-	-	2	10%	3	9%	1	6%
Hispanic/Latino Males	2	10%	3	11%	3	14%	2	6%	7	44%
Hispanic/Latina Females	1	5%	-	-	-	-	1	3%	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	1	4%	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	7%	-	-	-	-	1	6%
Total	20		27		21		32		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	9	64%	14	64%	13	76%	18	75%	7	54%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	14%	2	9%	-	-	-	-	-	-
Heterosexual contact	-	-	2	9%	-	-	1	4%	-	-
Other/unknown	3	21%	4	18%	4	24%	5	21%	6	46%
Subtotal	14	100%	22	100%	17	100%	24	100%	13	100%
Female adult or adolescent										
Injection drug use	-	-	1	20%	-	-	-	-	-	-
Heterosexual contact	4	67%	3	60%	2	67%	4	57%	2	67%
Other/unknown	2	33%	1	20%	1	33%	3	43%	1	33%
Subtotal	6	100%	5	100%	3	100%	7	100%	3	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	-	-
Total	20		27		21		32		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	9	45%	14	52%	13	62%	17	53%	7	44%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	4	20%	5	19%	2	10%	5	16%	2	13%
MSM & IDU	2	10%	1	4%	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	4%	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	1	3%	-	-
MSM & IDU & Heterosexual contact	-	-	1	4%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	1	3%	-	-
Other/unknown	5	25%	5	19%	6	29%	8	25%	7	44%
Total	20		27		21		32		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 8B

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	10	77%	12	71%	14	70%	18	86%	5	71%	595	81%
Females	3	23%	5	29%	6	30%	3	14%	2	29%	142	19%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	11	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	-	-	1	6%	2	10%	1	5%	-	-	16	2%
20-24	3	23%	5	29%	4	20%	2	10%	1	14%	67	9%
25-29	2	15%	3	18%	1	5%	3	14%	1	14%	125	17%
30-34	4	31%	1	6%	2	10%	3	14%	-	-	120	16%
35-39	2	15%	2	12%	3	15%	3	14%	-	-	151	20%
40-44	1	8%	2	12%	2	10%	3	14%	2	29%	102	14%
45-49	-	-	-	-	1	5%	1	5%	2	29%	69	9%
50-54	-	-	3	18%	2	10%	-	-	-	-	29	4%
55-64	1	8%	-	-	3	15%	4	19%	-	-	38	5%
65+	-	-	-	-	-	-	1	5%	1	14%	7	1%
Race/Ethnicity												
White, not Hispanic	8	62%	9	53%	14	70%	18	86%	3	43%	526	71%
Black/African American, not Hispanic	4	31%	5	29%	4	20%	1	5%	2	29%	123	17%
Hispanic/Latino	1	8%	2	12%	2	10%	1	5%	1	14%	76	10%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	1	6%	-	-	1	5%	1	14%	8	1%
Race/Sex												
White, not Hispanic Males	6	46%	7	41%	10	50%	16	76%	2	29%	448	61%
White, not Hispanic Females	2	15%	2	12%	4	20%	2	10%	1	14%	78	11%
Black/African American, not Hispanic Males	3	23%	3	18%	3	15%	-	-	1	14%	93	13%
Black/African American, not Hispanic Females	1	8%	2	12%	1	5%	1	5%	1	14%	30	4%
Hispanic/Latino Males	1	8%	1	6%	1	5%	1	5%	1	14%	47	6%
Hispanic/Latina Females	-	-	1	6%	1	5%	-	-	-	-	29	4%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	1	6%	-	-	1	5%	1	14%	8	1%
Total	13		17		20		21		7		737	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	90%	7	58%	9	64%	14	78%	2	40%	382	65%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	41	7%
Male-to-male sexual contact and IDU	-	-	1	8%	1	7%	-	-	-	-	43	7%
Heterosexual contact	1	10%	-	-	-	-	2	11%	1	20%	34	6%
Other/unknown	-	-	4	33%	4	29%	2	11%	2	40%	89	15%
Subtotal	10	100%	12	100%	14	100%	18	100%	5	100%	589	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	2	33%	1	33%	-	-	22	16%
Heterosexual contact	2	67%	3	60%	1	17%	1	33%	1	50%	93	68%
Other/unknown	1	33%	2	40%	3	50%	1	33%	1	50%	22	16%
Subtotal	3	100%	5	100%	6	100%	3	100%	2	100%	137	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	8	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	27%
Subtotal	-	-	-	-	-	-	-	-	-	-	11	100%
Total	13		17		20		21		7		737	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	7	54%	7	41%	8	40%	13	62%	2	29%	368	50%
Injection drug use (IDU) only	-	-	-	-	1	5%	-	-	-	-	38	5%
Heterosexual contact only	3	23%	3	18%	1	5%	3	14%	2	29%	127	17%
MSM & IDU	-	-	-	-	1	5%	-	-	-	-	37	5%
IDU & Heterosexual contact	-	-	-	-	1	5%	1	5%	-	-	25	3%
MSM & Heterosexual contact	2	15%	-	-	1	5%	1	5%	-	-	14	2%
MSM & IDU & Heterosexual contact	-	-	1	6%	-	-	-	-	-	-	6	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	8	1%
Other/unknown	1	8%	6	35%	7	35%	3	14%	3	43%	114	15%
Total	13		17		20		21		7		737	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 8B

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 8B

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	114.3	636	77%	295	76%	341	78%
Females	34.0	193	23%	94	24%	99	23%
Age at end of year							
<13	*	3	<1%	3	1%	-	-
13-14	*	2	<1%	-	-	2	<1%
15-19	*	4	<1%	3	1%	1	<1%
20-24	42.6	30	4%	24	6%	6	1%
25-29	108.3	66	8%	46	12%	20	5%
30-34	88.2	56	7%	32	8%	24	5%
35-39	128.7	83	10%	42	11%	41	9%
40-44	166.0	118	14%	57	15%	61	14%
45-49	195.4	146	18%	60	15%	86	20%
50-54	157.9	132	16%	48	12%	84	19%
55-64	93.3	146	18%	55	14%	91	21%
65+	23.0	43	5%	19	5%	24	5%
Race/Ethnicity							
White, not Hispanic	50.2	502	61%	233	60%	269	61%
Black/African American, not Hispanic	387.5	190	23%	86	22%	104	24%
Hispanic/Latino	265.3	121	15%	62	16%	59	13%
Asian/Pacific Islander	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	13	2%	7	2%	6	1%
Race/Sex							
White, not Hispanic Males	81.0	398	48%	181	47%	217	49%
White, not Hispanic Females	20.5	104	13%	52	13%	52	12%
Black/African American, not Hispanic Males	526.4	143	17%	63	16%	80	18%
Black/African American, not Hispanic Females	215.0	47	6%	23	6%	24	5%
Hispanic/Latino Males	365.3	85	10%	46	12%	39	9%
Hispanic/Latina Females	161.2	36	4%	16	4%	20	5%
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	13	2%	7	2%	6	1%
Total	73.7	829		389		440	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	400	63%	183	62%	217	64%
Injection drug use (IDU)	30	5%	14	5%	16	5%
Male-to-male sexual contact and IDU	34	5%	8	3%	26	8%
Heterosexual contact	51	8%	15	5%	36	11%
Other/unknown	119	19%	74	25%	45	13%
Subtotal	634	100%	294	100%	340	100%
Female adult or adolescent						
Injection drug use	18	10%	8	9%	10	11%
Heterosexual contact	132	72%	63	71%	69	73%
Other/unknown	34	18%	18	20%	16	17%
Subtotal	184	100%	89	100%	95	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	91%	6	100%	4	80%
Other/unknown	1	9%	-	-	1	20%
Subtotal	11	100%	6	100%	5	100%
Total	829		389		440	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	381	46%	175	45%	206	47%
Injection drug use (IDU) only	27	3%	14	4%	13	3%
Heterosexual contact only	183	22%	78	20%	105	24%
MSM & IDU	26	3%	5	1%	21	5%
IDU & Heterosexual contact	21	3%	8	2%	13	3%
MSM & Heterosexual contact	19	2%	8	2%	11	3%
MSM & IDU & Heterosexual contact	8	1%	3	1%	5	1%
Perinatal exposure	12	1%	6	2%	6	1%
Other/unknown	152	18%	92	24%	60	14%
Total	829		389		440	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 8B

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	276	69%	72	51%	45	53%	1	50%	-	-	6	75%
Injection drug use (IDU)	12	3%	8	6%	10	12%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	22	6%	8	6%	4	5%	-	-	-	-	-	-
Heterosexual contact	18	5%	21	15%	11	13%	-	-	-	-	1	13%
Other/unknown	70	18%	32	23%	15	18%	1	50%	-	-	1	13%
Subtotal	398	100%	141	100%	85	100%	2	100%	-	-	8	100%
Female adult or adolescent												
Injection drug use	12	12%	1	2%	4	11%	-	-	-	-	1	25%
Heterosexual contact	70	71%	36	80%	22	63%	-	-	1	100%	3	75%
Other/unknown	17	17%	8	18%	9	26%	-	-	-	-	-	-
Subtotal	99	100%	45	100%	35	100%	-	-	1	100%	4	100%
Child (<13 yrs at diagnosis)												
Perinatal	5	100%	4	100%	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	5	100%	4	100%	1	100%	-	-	-	-	1	100%
Total	502		190		121		2		1		13	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 8B

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	265	53%	67	35%	42	35%	1	50%	-	-	6	46%
Injection drug use (IDU) only	11	2%	6	3%	10	8%	-	-	-	-	-	-
Heterosexual contact only	88	18%	57	30%	33	27%	-	-	1	100%	4	31%
MSM & IDU	18	4%	6	3%	2	2%	-	-	-	-	-	-
IDU & Heterosexual contact	13	3%	3	2%	4	3%	-	-	-	-	1	8%
MSM & Heterosexual contact	11	2%	5	3%	3	2%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	4	1%	2	1%	2	2%	-	-	-	-	-	-
Perinatal exposure	5	1%	5	3%	1	1%	-	-	-	-	1	8%
Other/unknown	87	17%	39	21%	24	20%	1	50%	-	-	1	8%
Total	502		190		121		2		1		13	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 8B

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	517	76%	542	76%	584	77%	619	77%	636	77%
Females	164	24%	171	24%	178	23%	188	23%	193	23%
Age at end of year										
<13	5	1%	4	1%	4	1%	4	<1%	3	<1%
13-14	1	<1%	2	<1%	1	<1%	1	<1%	2	<1%
15-19	3	<1%	3	<1%	5	1%	4	<1%	4	<1%
20-24	34	5%	35	5%	41	5%	34	4%	30	4%
25-29	38	6%	44	6%	50	7%	57	7%	66	8%
30-34	58	9%	52	7%	55	7%	60	7%	56	7%
35-39	94	14%	94	13%	85	11%	89	11%	83	10%
40-44	126	19%	129	18%	124	16%	125	15%	118	14%
45-49	131	19%	125	18%	138	18%	135	17%	146	18%
50-54	89	13%	106	15%	121	16%	137	17%	132	16%
55-64	82	12%	95	13%	105	14%	122	15%	146	18%
65+	20	3%	24	3%	33	4%	39	5%	43	5%
Race/Ethnicity										
White, not Hispanic	414	61%	432	61%	465	61%	492	61%	502	61%
Black/African American, not Hispanic	156	23%	164	23%	174	23%	185	23%	190	23%
Hispanic/Latino	97	14%	102	14%	108	14%	115	14%	121	15%
Asian/Pacific Islander	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	11	2%	12	2%	12	2%	12	1%	13	2%
Race/Sex										
White, not Hispanic Males	329	48%	341	48%	370	49%	392	49%	398	48%
White, not Hispanic Females	85	12%	91	13%	95	12%	100	12%	104	13%
Black/African American, not Hispanic Males	117	17%	125	18%	132	17%	139	17%	143	17%
Black/African American, not Hispanic Females	39	6%	39	5%	42	6%	46	6%	47	6%
Hispanic/Latino Males	63	9%	67	9%	73	10%	79	10%	85	10%
Hispanic/Latina Females	34	5%	35	5%	35	5%	36	4%	36	4%
Asian/Pacific Islander Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	11	2%	12	2%	12	2%	12	1%	13	2%
Total	681		713		762		807		829	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 8B

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	328	64%	340	63%	369	63%	392	64%	400	63%
Injection drug use (IDU)	29	6%	29	5%	29	5%	30	5%	30	5%
Male-to-male sexual contact and IDU	29	6%	31	6%	34	6%	34	6%	34	5%
Heterosexual contact	44	9%	46	9%	47	8%	50	8%	51	8%
Other/unknown	85	17%	94	17%	103	18%	111	18%	119	19%
Subtotal	515	100%	540	100%	582	100%	617	100%	634	100%
Female adult or adolescent										
Injection drug use	15	10%	17	10%	18	11%	18	10%	18	10%
Heterosexual contact	118	76%	121	74%	123	72%	128	72%	132	72%
Other/unknown	23	15%	25	15%	29	17%	33	18%	34	18%
Subtotal	156	100%	163	100%	170	100%	179	100%	184	100%
Child (<13 yrs at diagnosis)										
Perinatal	9	90%	9	90%	9	90%	10	91%	10	91%
Other/unknown	1	10%	1	10%	1	10%	1	9%	1	9%
Subtotal	10	100%	10	100%	10	100%	11	100%	11	100%
Total	681		713		762		807		829	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 8B

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	313	46%	325	46%	352	46%	373	46%	381	46%
Injection drug use (IDU) only	25	4%	25	4%	26	3%	27	3%	27	3%
Heterosexual contact only	162	24%	167	23%	170	22%	178	22%	183	22%
MSM & IDU	24	4%	24	3%	26	3%	26	3%	26	3%
IDU & Heterosexual contact	19	3%	21	3%	21	3%	21	3%	21	3%
MSM & Heterosexual contact	15	2%	15	2%	17	2%	19	2%	19	2%
MSM & IDU & Heterosexual contact	5	1%	7	1%	8	1%	8	1%	8	1%
Perinatal exposure	11	2%	11	2%	11	1%	12	1%	12	1%
Other/unknown	107	16%	118	17%	131	17%	143	18%	152	18%
Total	681		713		762		807		829	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 8C

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 8C

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	5.5	29	91%	23	92%	3	75%	3	100%
Females	*	3	9%	2	8%	1	25%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	3	9%	3	12%	-	-	-	-
20-24	10.1	7	22%	7	28%	-	-	-	-
25-29	9.3	6	19%	5	20%	-	-	1	33%
30-34	7.7	5	16%	3	12%	1	25%	1	33%
35-39	*	2	6%	1	4%	1	25%	-	-
40-44	*	1	3%	-	-	1	25%	-	-
45-49	*	3	9%	2	8%	1	25%	-	-
50-54	*	2	6%	2	8%	-	-	-	-
55-64	*	3	9%	2	8%	-	-	1	33%
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	2.2	21	66%	17	68%	2	50%	2	67%
Black/African American, not Hispanic	18.7	8	25%	6	24%	1	25%	1	33%
Hispanic/Latino	*	3	9%	2	8%	1	25%	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Race/Sex									
White, not Hispanic Males	3.8	18	56%	15	60%	1	25%	2	67%
White, not Hispanic Females	*	3	9%	2	8%	1	25%	-	-
Black/African American, not Hispanic Males	32.2	8	25%	6	24%	1	25%	1	33%
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	3	9%	2	8%	1	25%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Total	3.0	32		25		4		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 8C

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	17	59%	13	57%	2	67%	2	67%
Injection drug use (IDU)	1	3%	1	4%	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	1	4%	-	-	-	-
Heterosexual contact	1	3%	1	4%	-	-	-	-
Other/unknown	9	31%	7	30%	1	33%	1	33%
Subtotal	29	100%	23	100%	3	100%	3	100%
Female adult or adolescent								
Injection drug use	2	67%	1	50%	1	100%	-	-
Heterosexual contact	1	33%	1	50%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	1	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	32		25		4		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 8C

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	50%	12	48%	2	50%	2	67%
Injection drug use (IDU) only	2	6%	2	8%	-	-	-	-
Heterosexual contact only	2	6%	2	8%	-	-	-	-
MSM & IDU	1	3%	1	4%	-	-	-	-
IDU & Heterosexual contact	1	3%	-	-	1	25%	-	-
MSM & Heterosexual contact	1	3%	1	4%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	9	28%	7	28%	1	25%	1	33%
Total	32		25		4		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 8C

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	11	61%	5	63%	1	33%	-	-	-	-	-	-
Injection drug use (IDU)	1	6%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	6%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	6%	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	22%	3	38%	2	67%	-	-	-	-	-	-
Subtotal	18	100%	8	100%	3	100%	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	2	67%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	33%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	21		8		3		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 8C

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	48%	5	63%	1	33%	-	-	-	-	-	-
Injection drug use (IDU) only	2	10%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	10%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	1	5%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	19%	3	38%	2	67%	-	-	-	-	-	-
Total	21		8		3		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 8C

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	5.8	31	86%	6.0	32	74%	7.0	37	86%	5.8	31	86%
Females	0.9	5	14%	2.1	11	26%	1.1	6	14%	0.9	5	14%
Age at diagnosis (yr)												
<13	*	2	6%	*	-	-	*	2	5%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	7.1	5	14%	*	1	2%	*	3	7%	*	1	3%
20-24	10.1	7	19%	14.5	10	23%	11.6	8	19%	17.3	12	33%
25-29	*	4	11%	*	4	9%	*	4	9%	7.7	5	14%
30-34	7.7	5	14%	12.4	8	19%	13.9	9	21%	*	3	8%
35-39	*	4	11%	14.1	9	21%	*	4	9%	9.4	6	17%
40-44	8.7	6	17%	*	4	9%	*	3	7%	*	-	-
45-49	*	1	3%	*	4	9%	*	3	7%	*	3	8%
50-54	*	2	6%	*	2	5%	7.6	6	14%	*	2	6%
55-64	*	-	-	*	-	-	*	1	2%	*	3	8%
65+	*	-	-	*	1	2%	*	-	-	*	1	3%
Race/Ethnicity												
White, not Hispanic	2.5	24	67%	3.3	32	74%	2.6	25	58%	2.9	28	78%
Black/African American, not Hispanic	21.0	9	25%	18.7	8	19%	28.0	12	28%	14.0	6	17%
Hispanic/Latino	*	1	3%	*	2	5%	*	2	5%	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	1	2%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	6%	*	1	2%	*	3	7%	*	2	6%
Race/Sex												
White, not Hispanic Males	4.6	22	61%	5.0	24	56%	4.8	23	53%	5.4	26	72%
White, not Hispanic Females	*	2	6%	1.6	8	19%	*	2	5%	*	2	6%
Black/African American, not Hispanic Males	24.1	6	17%	24.1	6	14%	32.2	8	19%	20.1	5	14%
Black/African American, not Hispanic Females	*	3	8%	*	2	5%	*	4	9%	*	1	3%
Hispanic/Latino Males	*	1	3%	*	2	5%	*	2	5%	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	1	2%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	6%	*	1	2%	*	3	7%	*	2	6%
Total	3.4	36		4.0	43		4.0	43		3.4	36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 8C

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	25	83%	19	59%	23	62%	24	77%
Injection drug use (IDU)	-	-	2	6%	1	3%	2	6%
Male-to-male sexual contact and IDU	1	3%	-	-	2	5%	-	-
Heterosexual contact	-	-	1	3%	1	3%	1	3%
Other/unknown	4	13%	10	31%	10	27%	4	13%
Subtotal	30	100%	32	100%	37	100%	31	100%
Female adult or adolescent								
Injection drug use	-	-	2	18%	-	-	2	40%
Heterosexual contact	3	75%	6	55%	1	25%	2	40%
Other/unknown	1	25%	3	27%	3	75%	1	20%
Subtotal	4	100%	11	100%	4	100%	5	100%
Child (<13 yrs at diagnosis)								
Perinatal	2	100%	-	-	2	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	2	100%	-	-	2	100%	-	-
Total	36		43		43		36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 8C

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	25	69%	19	44%	22	51%	24	67%
Injection drug use (IDU) only	-	-	1	2%	1	2%	1	3%
Heterosexual contact only	3	8%	7	16%	2	5%	3	8%
MSM & IDU	1	3%	-	-	1	2%	-	-
IDU & Heterosexual contact	-	-	3	7%	-	-	3	8%
MSM & Heterosexual contact	-	-	-	-	1	2%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	1	2%	-	-
Perinatal exposure	2	6%	-	-	2	5%	-	-
Other/unknown	5	14%	13	30%	13	30%	5	14%
Total	36		43		43		36	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the highest category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 8C

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	18	86%	21	81%	20	80%	19	86%	23	92%
Females	3	14%	5	19%	5	20%	3	14%	2	8%
Age at HIV diagnosis (yr)										
<13	2	10%	-	-	2	8%	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	14%	1	4%	2	8%	1	5%	3	12%
20-24	5	24%	7	27%	7	28%	9	41%	7	28%
25-29	2	10%	2	8%	3	12%	4	18%	5	20%
30-34	3	14%	3	12%	5	20%	1	5%	3	12%
35-39	3	14%	7	27%	2	8%	4	18%	1	4%
40-44	3	14%	2	8%	1	4%	-	-	-	-
45-49	-	-	1	4%	-	-	1	5%	2	8%
50-54	-	-	2	8%	3	12%	1	5%	2	8%
55-64	-	-	-	-	-	-	1	5%	2	8%
65+	-	-	1	4%	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	15	71%	17	65%	15	60%	19	86%	17	68%
Black/African American, not Hispanic	5	24%	7	27%	8	32%	2	9%	6	24%
Hispanic/Latino	-	-	2	8%	1	4%	-	-	2	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	-	-	1	4%	1	5%	-	-
Race/Sex										
White, not Hispanic Males	14	67%	13	50%	13	52%	17	77%	15	60%
White, not Hispanic Females	1	5%	4	15%	2	8%	2	9%	2	8%
Black/African American, not Hispanic Males	3	14%	6	23%	5	20%	2	9%	6	24%
Black/African American, not Hispanic Females	2	10%	1	4%	3	12%	-	-	-	-
Hispanic/Latino Males	-	-	2	8%	1	4%	-	-	2	8%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	-	-	1	4%	1	5%	-	-
Total	21		26		25		22		25	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 8C

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	15	88%	12	57%	15	75%	16	84%	13	57%
Injection drug use (IDU)	-	-	2	10%	1	5%	2	11%	1	4%
Male-to-male sexual contact and IDU	1	6%	-	-	-	-	-	-	1	4%
Heterosexual contact	-	-	-	-	1	5%	-	-	1	4%
Other/unknown	1	6%	7	33%	3	15%	1	5%	7	30%
Subtotal	17	100%	21	100%	20	100%	19	100%	23	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	2	67%	1	50%
Heterosexual contact	1	50%	3	60%	1	33%	-	-	1	50%
Other/unknown	1	50%	2	40%	2	67%	1	33%	-	-
Subtotal	2	100%	5	100%	3	100%	3	100%	2	100%
Child (<13 yrs at diagnosis)										
Perinatal	2	100%	-	-	2	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	-	-	2	100%	-	-	-	-
Total	21		26		25		22		25	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 8C

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	15	71%	12	46%	15	60%	16	73%	12	48%
Injection drug use (IDU) only	-	-	-	-	1	4%	1	5%	2	8%
Heterosexual contact only	1	5%	3	12%	2	8%	-	-	2	8%
MSM & IDU	1	5%	-	-	-	-	-	-	1	4%
IDU & Heterosexual contact	-	-	2	8%	-	-	3	14%	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	1	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	2	10%	-	-	2	8%	-	-	-	-
Other/unknown	2	10%	9	35%	5	20%	2	9%	7	28%
Total	21		26		25		22		25	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 8C

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	16	80%	13	72%	21	91%	16	84%	14	88%	721	87%
Females	4	20%	5	28%	2	9%	3	16%	2	13%	108	13%
Age at AIDS diagnosis (yr)												
<13	1	5%	-	-	-	-	-	-	-	-	10	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	2	10%	-	-	2	9%	-	-	-	-	17	2%
20-24	4	20%	3	17%	2	9%	4	21%	1	6%	75	9%
25-29	2	10%	1	6%	1	4%	3	16%	2	13%	133	16%
30-34	3	15%	5	28%	5	22%	2	11%	3	19%	168	20%
35-39	2	10%	2	11%	3	13%	4	21%	4	25%	142	17%
40-44	3	15%	4	22%	2	9%	-	-	2	13%	137	17%
45-49	1	5%	2	11%	3	13%	2	11%	3	19%	63	8%
50-54	2	10%	1	6%	3	13%	1	5%	-	-	43	5%
55-64	-	-	-	-	2	9%	2	11%	1	6%	28	3%
65+	-	-	-	-	-	-	1	5%	-	-	13	2%
Race/Ethnicity												
White, not Hispanic	12	60%	17	94%	11	48%	14	74%	10	63%	660	80%
Black/African American, not Hispanic	6	30%	1	6%	7	30%	4	21%	3	19%	131	16%
Hispanic/Latino	1	5%	-	-	1	4%	-	-	2	13%	18	2%
Asian/Pacific Islander	-	-	-	-	1	4%	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	-	-	3	13%	1	5%	1	6%	16	2%
Race/Sex												
White, not Hispanic Males	10	50%	13	72%	11	48%	13	68%	8	50%	581	70%
White, not Hispanic Females	2	10%	4	22%	-	-	1	5%	2	13%	79	10%
Black/African American, not Hispanic Males	4	20%	-	-	6	26%	3	16%	3	19%	106	13%
Black/African American, not Hispanic Females	2	10%	1	6%	1	4%	1	5%	-	-	25	3%
Hispanic/Latino Males	1	5%	-	-	1	4%	-	-	2	13%	17	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	-	-	-	-	1	4%	-	-	-	-	4	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	-	-	3	13%	1	5%	1	6%	16	2%
Total	20		18		23		19		16		829	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 8C

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	12	80%	9	69%	11	52%	11	69%	6	43%	491	69%
Injection drug use (IDU)	-	-	1	8%	-	-	-	-	-	-	42	6%
Male-to-male sexual contact and IDU	-	-	-	-	3	14%	-	-	1	7%	32	4%
Heterosexual contact	-	-	1	8%	1	5%	1	6%	2	14%	40	6%
Other/unknown	3	20%	2	15%	6	29%	4	25%	5	36%	110	15%
Subtotal	15	100%	13	100%	21	100%	16	100%	14	100%	715	100%
Female adult or adolescent												
Injection drug use	-	-	1	20%	-	-	1	33%	1	50%	13	13%
Heterosexual contact	3	75%	3	60%	1	50%	2	67%	1	50%	75	72%
Other/unknown	1	25%	1	20%	1	50%	-	-	-	-	16	15%
Subtotal	4	100%	5	100%	2	100%	3	100%	2	100%	104	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	50%
Other/unknown	1	100%	-	-	-	-	-	-	-	-	5	50%
Subtotal	1	100%	-	-	-	-	-	-	-	-	10	100%
Total	20		18		23		19		16		829	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 8C

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	12	60%	9	50%	10	43%	11	58%	6	38%	470	57%
Injection drug use (IDU) only	-	-	1	6%	-	-	1	5%	-	-	40	5%
Heterosexual contact only	3	15%	4	22%	2	9%	3	16%	3	19%	115	14%
MSM & IDU	-	-	-	-	2	9%	-	-	-	-	23	3%
IDU & Heterosexual contact	-	-	1	6%	-	-	-	-	1	6%	15	2%
MSM & Heterosexual contact	-	-	-	-	1	4%	-	-	-	-	21	3%
MSM & IDU & Heterosexual contact	-	-	-	-	1	4%	-	-	1	6%	9	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	5	25%	3	17%	7	30%	4	21%	5	31%	131	16%
Total	20		18		23		19		16		829	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 8C

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 8C

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	141.5	751	82%	332	79%	419	84%
Females	31.2	167	18%	87	21%	80	16%
Age at end of year							
<13	*	4	<1%	2	<1%	2	<1%
13-14	*	1	<1%	1	<1%	-	-
15-19	*	3	<1%	2	<1%	1	<1%
20-24	73.7	51	6%	42	10%	9	2%
25-29	104.9	68	7%	49	12%	19	4%
30-34	103.7	67	7%	37	9%	30	6%
35-39	158.7	101	11%	56	13%	45	9%
40-44	168.1	116	13%	46	11%	70	14%
45-49	210.2	151	16%	48	11%	103	21%
50-54	198.7	156	17%	47	11%	109	22%
55-64	106.0	155	17%	67	16%	88	18%
65+	26.8	45	5%	22	5%	23	5%
Race/Ethnicity							
White, not Hispanic	68.0	660	72%	298	71%	362	73%
Black/African American, not Hispanic	462.8	198	22%	95	23%	103	21%
Hispanic/Latino	154.2	29	3%	14	3%	15	3%
Asian/Pacific Islander	*	4	<1%	1	<1%	3	1%
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	25	3%	10	2%	15	3%
Race/Sex							
White, not Hispanic Males	114.4	549	60%	240	57%	309	62%
White, not Hispanic Females	22.6	111	12%	58	14%	53	11%
Black/African American, not Hispanic Males	614.9	153	17%	73	17%	80	16%
Black/African American, not Hispanic Females	251.4	45	5%	22	5%	23	5%
Hispanic/Latino Males	271.4	27	3%	12	3%	15	3%
Hispanic/Latina Females	*	2	<1%	2	<1%	-	-
Asian/Pacific Islander Males	*	3	<1%	-	-	3	1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	25	3%	10	2%	15	3%
Total	86.1	918		419		499	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 8C

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	514	69%	225	68%	289	70%
Injection drug use (IDU)	31	4%	13	4%	18	4%
Male-to-male sexual contact and IDU	38	5%	13	4%	25	6%
Heterosexual contact	52	7%	18	5%	34	8%
Other/unknown	108	15%	61	18%	47	11%
Subtotal	743	100%	330	100%	413	100%
Female adult or adolescent						
Injection drug use	20	13%	10	12%	10	13%
Heterosexual contact	102	64%	45	55%	57	73%
Other/unknown	38	24%	27	33%	11	14%
Subtotal	160	100%	82	100%	78	100%
Child (<13 yrs at diagnosis)						
Perinatal	11	73%	7	100%	4	50%
Other/unknown	4	27%	-	-	4	50%
Subtotal	15	100%	7	100%	8	100%
Total	918		419		499	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 8C

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	484	53%	214	51%	270	54%
Injection drug use (IDU) only	24	3%	12	3%	12	2%
Heterosexual contact only	154	17%	63	15%	91	18%
MSM & IDU	29	3%	13	3%	16	3%
IDU & Heterosexual contact	27	3%	11	3%	16	3%
MSM & Heterosexual contact	30	3%	11	3%	19	4%
MSM & IDU & Heterosexual contact	9	1%	-	-	9	2%
Perinatal exposure	11	1%	7	2%	4	1%
Other/unknown	150	16%	88	21%	62	12%
Total	918		419		499	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 8C

Living with a Diagnosis of HIV Infection in 2014												
Transmission Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	405	74%	79	53%	15	56%	1	33%	1	100%	13	76%
Injection drug use (IDU)	20	4%	8	5%	3	11%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	27	5%	11	7%	-	-	-	-	-	-	-	-
Heterosexual contact	24	4%	23	15%	3	11%	1	33%	-	-	1	6%
Other/unknown	69	13%	29	19%	6	22%	1	33%	-	-	3	18%
Subtotal	545	100%	150	100%	27	100%	3	100%	1	100%	17	100%
Female adult or adolescent												
Injection drug use	15	14%	4	10%	-	-	-	-	-	-	1	14%
Heterosexual contact	69	63%	26	65%	1	100%	1	100%	1	100%	4	57%
Other/unknown	26	24%	10	25%	-	-	-	-	-	-	2	29%
Subtotal	110	100%	40	100%	1	100%	1	100%	1	100%	7	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	40%	7	88%	1	100%	-	-	-	-	1	100%
Other/unknown	3	60%	1	13%	-	-	-	-	-	-	-	-
Subtotal	5	100%	8	100%	1	100%	-	-	-	-	1	100%
Total	660		198		29		4		2		25	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 8C

Living with a Diagnosis of HIV Infection in 2014												
Exposure Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	387	59%	71	36%	13	45%	1	25%	1	50%	11	44%
Injection drug use (IDU) only	19	3%	5	3%	-	-	-	-	-	-	-	-
Heterosexual contact only	93	14%	49	25%	4	14%	2	50%	1	50%	5	20%
MSM & IDU	22	3%	7	4%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	16	2%	7	4%	3	10%	-	-	-	-	1	4%
MSM & Heterosexual contact	18	3%	8	4%	2	7%	-	-	-	-	2	8%
MSM & IDU & Heterosexual contact	5	1%	4	2%	-	-	-	-	-	-	-	-
Perinatal exposure	2	0%	7	4%	1	3%	-	-	-	-	1	4%
Other/unknown	98	15%	40	20%	6	21%	1	25%	-	-	5	20%
Total	660		198		29		4		2		25	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 8C

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	623	82%	654	81%	692	81%	719	81%	751	82%
Females	136	18%	151	19%	158	19%	164	19%	167	18%
Age at end of year										
<13	3	<1%	3	<1%	5	1%	5	1%	4	<1%
13-14	1	<1%	1	<1%	-	-	-	-	1	<1%
15-19	12	2%	7	1%	8	1%	3	<1%	3	<1%
20-24	39	5%	48	6%	47	6%	56	6%	51	6%
25-29	40	5%	41	5%	47	6%	55	6%	68	7%
30-34	74	10%	77	10%	79	9%	70	8%	67	7%
35-39	87	11%	99	12%	105	12%	96	11%	101	11%
40-44	128	17%	119	15%	120	14%	123	14%	116	13%
45-49	150	20%	160	20%	159	19%	153	17%	151	16%
50-54	117	15%	121	15%	134	16%	149	17%	156	17%
55-64	84	11%	102	13%	115	14%	134	15%	155	17%
65+	24	3%	27	3%	31	4%	39	4%	45	5%
Race/Ethnicity										
White, not Hispanic	557	73%	588	73%	613	72%	638	72%	660	72%
Black/African American, not Hispanic	157	21%	167	21%	182	21%	188	21%	198	22%
Hispanic/Latino	23	3%	25	3%	26	3%	26	3%	29	3%
Asian/Pacific Islander	2	<1%	3	<1%	4	<1%	4	<1%	4	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	18	2%	20	2%	23	3%	25	3%	25	3%
Race/Sex										
White, not Hispanic Males	464	61%	485	60%	508	60%	530	60%	549	60%
White, not Hispanic Females	93	12%	103	13%	105	12%	108	12%	111	12%
Black/African American, not Hispanic Males	121	16%	128	16%	138	16%	143	16%	153	17%
Black/African American, not Hispanic Females	36	5%	39	5%	44	5%	45	5%	45	5%
Hispanic/Latino Males	21	3%	23	3%	24	3%	24	3%	27	3%
Hispanic/Latina Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Asian/Pacific Islander Males	2	<1%	2	<1%	3	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	18	2%	20	2%	23	3%	25	3%	25	3%
Total	759		805		850		883		918	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 8C

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	423	69%	444	69%	472	69%	493	69%	514	69%
Injection drug use (IDU)	27	4%	29	4%	29	4%	30	4%	31	4%
Male-to-male sexual contact and IDU	35	6%	35	5%	36	5%	37	5%	38	5%
Heterosexual contact	51	8%	52	8%	52	8%	52	7%	52	7%
Other/unknown	79	13%	86	13%	95	14%	99	14%	108	15%
Subtotal	615	100%	646	100%	684	100%	711	100%	743	100%
Female adult or adolescent										
Injection drug use	13	10%	15	10%	15	10%	17	11%	20	13%
Heterosexual contact	90	69%	98	67%	99	66%	102	65%	102	64%
Other/unknown	28	21%	33	23%	37	25%	38	24%	38	24%
Subtotal	131	100%	146	100%	151	100%	157	100%	160	100%
Child (<13 yrs at diagnosis)										
Perinatal	9	69%	9	69%	11	73%	11	73%	11	73%
Other/unknown	4	31%	4	31%	4	27%	4	27%	4	27%
Subtotal	13	100%	13	100%	15	100%	15	100%	15	100%
Total	759		805		850		883		918	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 8C

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	397	52%	417	52%	444	52%	464	53%	484	53%
Injection drug use (IDU) only	20	3%	21	3%	21	2%	22	2%	24	3%
Heterosexual contact only	141	19%	150	19%	151	18%	154	17%	154	17%
MSM & IDU	26	3%	26	3%	27	3%	28	3%	29	3%
IDU & Heterosexual contact	20	3%	23	3%	23	3%	25	3%	27	3%
MSM & Heterosexual contact	26	3%	27	3%	28	3%	29	3%	30	3%
MSM & IDU & Heterosexual contact	9	1%	9	1%	9	1%	9	1%	9	1%
Perinatal exposure	9	1%	9	1%	11	1%	11	1%	11	1%
Other/unknown	111	15%	123	15%	136	16%	141	16%	150	16%
Total	759		805		850		883		918	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Region 8D

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Region 8D

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	5.7	23	92%	15	94%	7	88%	1	100%
Females	*	2	8%	1	6%	1	13%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	1	4%	1	6%	-	-	-	-
20-24	*	3	12%	3	19%	-	-	-	-
25-29	*	1	4%	1	6%	-	-	-	-
30-34	*	4	16%	3	19%	1	13%	-	-
35-39	*	3	12%	2	13%	1	13%	-	-
40-44	11.6	6	24%	5	31%	-	-	1	100%
45-49	*	2	8%	-	-	2	25%	-	-
50-54	*	2	8%	1	6%	1	13%	-	-
55-64	*	2	8%	-	-	2	25%	-	-
65+	*	1	4%	-	-	1	13%	-	-
Race/Ethnicity									
White, not Hispanic	2.5	19	76%	11	69%	7	88%	1	100%
Black/African American, not Hispanic	*	4	16%	3	19%	1	13%	-	-
Hispanic/Latino	*	2	8%	2	13%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Race/Sex									
White, not Hispanic Males	4.6	17	68%	10	63%	6	75%	1	100%
White, not Hispanic Females	*	2	8%	1	6%	1	13%	-	-
Black/African American, not Hispanic Males	*	4	16%	3	19%	1	13%	-	-
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	2	8%	2	13%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Total	3.1	25		16		8		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Region 8D

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	12	52%	10	67%	1	14%	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	9%	2	13%	-	-	-	-
Heterosexual contact	1	4%	1	7%	-	-	-	-
Other/unknown	8	35%	2	13%	6	86%	-	-
Subtotal	23	100%	15	100%	7	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	2	100%	1	100%	1	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	25		16		8		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Region 8D

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	11	44%	10	63%	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	3	12%	2	13%	1	13%	-	-
MSM & IDU	2	8%	2	13%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	4%	-	-	1	13%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	8	32%	2	13%	6	75%	-	-
Total	25		16		8		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Region 8D

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	8	47%	2	50%	2	100%	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	12%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	25%	-	-	-	-	-	-	-	-
Other/unknown	7	41%	1	25%	-	-	-	-	-	-	-	-
Subtotal	17	100%	4	100%	2	100%	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	19		4		2		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Region 8D

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	37%	2	50%	2	100%	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	11%	1	25%	-	-	-	-	-	-	-	-
MSM & IDU	2	11%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	7	37%	1	25%	-	-	-	-	-	-	-	-
Total	19		4		2		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Region 8D

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	6.4	26	79%	5.2	21	84%	4.7	19	68%	4.9	20	87%
Females	1.7	7	21%	*	4	16%	2.2	9	32%	*	3	13%
Age at diagnosis (yr)												
<13	*	1	3%	*	-	-	*	1	4%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	3%	*	1	4%	*	1	4%	*	-	-
20-24	15.3	9	27%	15.3	9	36%	10.2	6	21%	8.5	5	22%
25-29	*	3	9%	*	3	12%	*	3	11%	*	4	17%
30-34	10.6	5	15%	*	3	12%	*	4	14%	*	2	9%
35-39	*	2	6%	*	2	8%	*	1	4%	*	2	9%
40-44	17.4	9	27%	*	-	-	*	4	14%	*	3	13%
45-49	*	1	3%	*	2	8%	*	3	11%	*	2	9%
50-54	*	1	3%	*	2	8%	*	2	7%	*	4	17%
55-64	*	1	3%	*	3	12%	*	3	11%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	1	4%
Race/Ethnicity												
White, not Hispanic	2.7	20	61%	2.8	21	84%	2.8	21	75%	2.3	17	74%
Black/African American, not Hispanic	31.7	9	27%	*	2	8%	*	3	11%	*	4	17%
Hispanic/Latino	*	2	6%	*	1	4%	*	-	-	*	1	4%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	4%
Unknown	*	2	6%	*	1	4%	*	4	14%	*	-	-
Race/Sex												
White, not Hispanic Males	4.6	17	52%	4.6	17	68%	4.3	16	57%	4.3	16	70%
White, not Hispanic Females	*	3	9%	*	4	16%	1.3	5	18%	*	1	4%
Black/African American, not Hispanic Males	31.2	5	15%	*	2	8%	*	1	4%	*	3	13%
Black/African American, not Hispanic Females	*	4	12%	*	-	-	*	2	7%	*	1	4%
Hispanic/Latino Males	*	2	6%	*	1	4%	*	-	-	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	1	4%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	4%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	6%	*	1	4%	*	4	14%	*	-	-
Total	4.0	33		3.1	25		3.4	28		2.8	23	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AII infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Region 8D

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	20	80%	15	71%	17	89%	16	80%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	4%	2	10%	-	-	-	-
Other/unknown	4	16%	4	19%	2	11%	4	20%
Subtotal	25	100%	21	100%	19	100%	20	100%
Female adult or adolescent								
Injection drug use	-	-	2	50%	-	-	-	-
Heterosexual contact	3	43%	2	50%	5	63%	2	67%
Other/unknown	4	57%	-	-	3	38%	1	33%
Subtotal	7	100%	4	100%	8	100%	3	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	-	-	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%	-	-
Total	33		25		28		23	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Region 8D

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	19	58%	15	60%	17	61%	15	65%
Injection drug use (IDU) only	-	-	1	4%	-	-	-	-
Heterosexual contact only	4	12%	4	16%	5	18%	2	9%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	4%	-	-	-	-
MSM & Heterosexual contact	1	3%	-	-	-	-	1	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	1	3%	-	-	1	4%	-	-
Other/unknown	8	24%	4	16%	5	18%	5	22%
Total	33		25		28		23	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Region 8D

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	13	81%	11	85%	9	64%	16	84%	15	94%
Females	3	19%	2	15%	5	36%	3	16%	1	6%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	6%	1	8%	1	7%	-	-	1	6%
20-24	6	38%	6	46%	5	36%	4	21%	3	19%
25-29	2	13%	3	23%	1	7%	4	21%	1	6%
30-34	4	25%	1	8%	-	-	2	11%	3	19%
35-39	1	6%	1	8%	-	-	2	11%	2	13%
40-44	2	13%	-	-	2	14%	3	16%	5	31%
45-49	-	-	-	-	2	14%	2	11%	-	-
50-54	-	-	1	8%	1	7%	2	11%	1	6%
55-64	-	-	-	-	2	14%	-	-	-	-
65+	-	-	-	-	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	7	44%	10	77%	13	93%	14	74%	11	69%
Black/African American, not Hispanic	5	31%	1	8%	-	-	3	16%	3	19%
Hispanic/Latino	2	13%	1	8%	-	-	1	5%	2	13%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	1	5%	-	-
Unknown	2	13%	1	8%	1	7%	-	-	-	-
Race/Sex										
White, not Hispanic Males	6	38%	8	62%	8	57%	13	68%	10	63%
White, not Hispanic Females	1	6%	2	15%	5	36%	1	5%	1	6%
Black/African American, not Hispanic Males	3	19%	1	8%	-	-	2	11%	3	19%
Black/African American, not Hispanic Females	2	13%	-	-	-	-	1	5%	-	-
Hispanic/Latino Males	2	13%	1	8%	-	-	-	-	2	13%
Hispanic/Latina Females	-	-	-	-	-	-	1	5%	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	1	5%	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	13%	1	8%	1	7%	-	-	-	-
Total	16		13		14		19		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Region 8D

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	11	85%	8	73%	8	89%	14	88%	10	67%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	2	13%
Heterosexual contact	1	8%	2	18%	-	-	-	-	1	7%
Other/unknown	1	8%	1	9%	1	11%	2	13%	2	13%
Subtotal	13	100%	11	100%	9	100%	16	100%	15	100%
Female adult or adolescent										
Injection drug use	-	-	2	100%	-	-	-	-	-	-
Heterosexual contact	2	67%	-	-	4	80%	2	67%	1	100%
Other/unknown	1	33%	-	-	1	20%	1	33%	-	-
Subtotal	3	100%	2	100%	5	100%	3	100%	1	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	16		13		14		19		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Region 8D

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	63%	8	62%	8	57%	13	68%	10	63%
Injection drug use (IDU) only	-	-	1	8%	-	-	-	-	-	-
Heterosexual contact only	3	19%	2	15%	4	29%	2	11%	2	13%
MSM & IDU	-	-	-	-	-	-	-	-	2	13%
IDU & Heterosexual contact	-	-	1	8%	-	-	-	-	-	-
MSM & Heterosexual contact	1	6%	-	-	-	-	1	5%	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	13%	1	8%	2	14%	3	16%	2	13%
Total	16		13		14		19		16	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 8D

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	15	75%	11	85%	11	61%	11	85%	9	90%	472	85%
Females	5	25%	2	15%	7	39%	2	15%	1	10%	84	15%
Age at AIDS diagnosis (yr)												
<13	1	5%	-	-	1	6%	-	-	-	-	11	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	1	5%	-	-	1	6%	-	-	-	-	8	1%
20-24	2	10%	3	23%	1	6%	2	15%	1	10%	47	8%
25-29	1	5%	-	-	2	11%	4	31%	-	-	73	13%
30-34	2	10%	2	15%	4	22%	-	-	-	-	104	19%
35-39	3	15%	2	15%	2	11%	3	23%	2	20%	104	19%
40-44	7	35%	2	15%	2	11%	-	-	1	10%	91	16%
45-49	1	5%	-	-	4	22%	1	8%	2	20%	43	8%
50-54	1	5%	1	8%	1	6%	2	15%	1	10%	35	6%
55-64	1	5%	3	23%	-	-	1	8%	2	20%	34	6%
65+	-	-	-	-	-	-	-	-	1	10%	5	1%
Race/Ethnicity												
White, not Hispanic	15	75%	13	100%	12	67%	11	85%	8	80%	483	87%
Black/African American, not Hispanic	5	25%	-	-	3	17%	2	15%	2	20%	51	9%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	10	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	3	17%	-	-	-	-	11	2%
Race/Sex												
White, not Hispanic Males	13	65%	11	85%	9	50%	9	69%	7	70%	417	75%
White, not Hispanic Females	2	10%	2	15%	3	17%	2	15%	1	10%	66	12%
Black/African American, not Hispanic Males	2	10%	-	-	1	6%	2	15%	2	20%	39	7%
Black/African American, not Hispanic Females	3	15%	-	-	2	11%	-	-	-	-	12	2%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	8	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	3	17%	-	-	-	-	11	2%
Total	20		13		18		13		10		556	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 8D

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	64%	8	73%	9	82%	7	64%	3	33%	298	64%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	23	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	9%	-	-	21	5%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	30	6%
Other/unknown	5	36%	3	27%	2	18%	3	27%	6	67%	92	20%
Subtotal	14	100%	11	100%	11	100%	11	100%	9	100%	464	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	10	12%
Heterosexual contact	2	40%	2	100%	4	67%	2	100%	1	100%	57	70%
Other/unknown	3	60%	-	-	2	33%	-	-	-	-	14	17%
Subtotal	5	100%	2	100%	6	100%	2	100%	1	100%	81	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	1	100%	-	-	-	-	8	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	27%
Subtotal	1	100%	-	-	1	100%	-	-	-	-	11	100%
Total	20		13		18		13		10		556	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 8D

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	9	45%	8	62%	9	50%	7	54%	2	20%	288	52%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	20	4%
Heterosexual contact only	2	10%	2	15%	4	22%	2	15%	1	10%	87	16%
MSM & IDU	-	-	-	-	-	-	1	8%	-	-	18	3%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	13	2%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	1	10%	10	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	1%
Perinatal exposure	1	5%	-	-	1	6%	-	-	-	-	8	1%
Other/unknown	8	40%	3	23%	4	22%	3	23%	6	60%	109	20%
Total	20		13		18		13		10		556	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Region 8D

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Region 8D

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	130.9	530	80%	230	78%	300	82%
Females	31.9	131	20%	64	22%	67	18%
Age at end of year							
<13	4.0	5	1%	1	<1%	4	1%
13-14	*	1	<1%	1	<1%	-	-
15-19	*	4	1%	4	1%	-	-
20-24	42.4	25	4%	21	7%	4	1%
25-29	47.3	22	3%	17	6%	5	1%
30-34	120.7	57	9%	38	13%	19	5%
35-39	101.2	47	7%	22	7%	25	7%
40-44	158.4	82	12%	40	14%	42	11%
45-49	225.4	121	18%	45	15%	76	21%
50-54	185.3	112	17%	39	13%	73	20%
55-64	135.4	153	23%	55	19%	98	27%
65+	24.2	32	5%	11	4%	21	6%
Race/Ethnicity							
White, not Hispanic	69.9	526	80%	232	79%	294	80%
Black/African American, not Hispanic	313.3	89	13%	39	13%	50	14%
Hispanic/Latino	144.3	16	2%	9	3%	7	2%
Asian/Pacific Islander	*	3	<1%	1	<1%	2	1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	26	4%	12	4%	14	4%
Race/Sex							
White, not Hispanic Males	115.7	430	65%	184	63%	246	67%
White, not Hispanic Females	25.2	96	15%	48	16%	48	13%
Black/African American, not Hispanic Males	430.7	69	10%	29	10%	40	11%
Black/African American, not Hispanic Females	161.5	20	3%	10	3%	10	3%
Hispanic/Latino Males	181.3	10	2%	6	2%	4	1%
Hispanic/Latina Females	107.8	6	1%	3	1%	3	1%
Asian/Pacific Islander Males	*	3	<1%	1	<1%	2	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	26	4%	12	4%	14	4%
Total	81.0	661		294		367	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Region 8D

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	348	67%	152	67%	196	67%
Injection drug use (IDU)	21	4%	2	1%	19	6%
Male-to-male sexual contact and IDU	25	5%	11	5%	14	5%
Heterosexual contact	35	7%	14	6%	21	7%
Other/unknown	91	18%	47	21%	44	15%
Subtotal	520	100%	226	100%	294	100%
Female adult or adolescent						
Injection drug use	11	9%	5	8%	6	9%
Heterosexual contact	84	68%	32	54%	52	80%
Other/unknown	29	23%	22	37%	7	11%
Subtotal	124	100%	59	100%	65	100%
Child (<13 yrs at diagnosis)						
Perinatal	13	76%	7	78%	6	75%
Other/unknown	4	24%	2	22%	2	25%
Subtotal	17	100%	9	100%	8	100%
Total	661		294		367	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Region 8D

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	334	51%	147	50%	187	51%
Injection drug use (IDU) only	20	3%	3	1%	17	5%
Heterosexual contact only	119	18%	46	16%	73	20%
MSM & IDU	20	3%	9	3%	11	3%
IDU & Heterosexual contact	12	2%	4	1%	8	2%
MSM & Heterosexual contact	14	2%	5	2%	9	2%
MSM & IDU & Heterosexual contact	5	1%	2	1%	3	1%
Perinatal exposure	14	2%	8	3%	6	2%
Other/unknown	123	19%	70	24%	53	14%
Total	661		294		367	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Region 8D

Living with a Diagnosis of HIV Infection in 2014												
Transmission Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	300	71%	27	42%	7	70%	1	33%	1	100%	12	71%
Injection drug use (IDU)	14	3%	6	9%	1	10%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	19	4%	5	8%	-	-	1	33%	-	-	-	-
Heterosexual contact	25	6%	8	12%	1	10%	-	-	-	-	1	6%
Other/unknown	66	16%	19	29%	1	10%	1	33%	-	-	4	24%
Subtotal	424	100%	65	100%	10	100%	3	100%	1	100%	17	100%
Female adult or adolescent												
Injection drug use	9	10%	-	-	1	25%	-	-	-	-	1	13%
Heterosexual contact	66	71%	13	68%	2	50%	-	-	-	-	3	38%
Other/unknown	18	19%	6	32%	1	25%	-	-	-	-	4	50%
Subtotal	93	100%	19	100%	4	100%	-	-	-	-	8	100%
Child (<13 yrs at diagnosis)												
Perinatal	5	56%	5	100%	2	100%	-	-	-	-	1	100%
Other/unknown	4	44%	-	-	-	-	-	-	-	-	-	-
Subtotal	9	100%	5	100%	2	100%	-	-	-	-	1	100%
Total	526		89		16		3		1		26	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Region 8D

Living with a Diagnosis of HIV Infection in 2014												
Exposure Category ^a	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	291	55%	23	26%	7	44%	1	33%	1	100%	11	42%
Injection drug use (IDU) only	13	2%	4	4%	2	13%	-	-	-	-	1	4%
Heterosexual contact only	91	17%	21	24%	3	19%	-	-	-	-	4	15%
MSM & IDU	15	3%	4	4%	-	-	1	33%	-	-	-	-
IDU & Heterosexual contact	10	2%	2	2%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	9	2%	4	4%	-	-	-	-	-	-	1	4%
MSM & IDU & Heterosexual contact	4	1%	1	1%	-	-	-	-	-	-	-	-
Perinatal exposure	6	1%	5	6%	2	13%	-	-	-	-	1	4%
Other/unknown	87	17%	25	28%	2	13%	1	33%	-	-	8	31%
Total	526		89		16		3		1		26	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Region 8D

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	434	79%	460	80%	488	80%	506	80%	530	80%
Females	112	21%	117	20%	124	20%	128	20%	131	20%
Age at end of year										
<13	5	1%	5	1%	6	1%	6	1%	5	1%
13-14	2	<1%	2	<1%	-	-	-	-	1	<1%
15-19	3	1%	2	<1%	5	1%	6	1%	4	1%
20-24	10	2%	12	2%	19	3%	24	4%	25	4%
25-29	29	5%	29	5%	22	4%	21	3%	22	3%
30-34	32	6%	40	7%	53	9%	53	8%	57	9%
35-39	64	12%	54	9%	46	8%	43	7%	47	7%
40-44	112	21%	103	18%	92	15%	87	14%	82	12%
45-49	103	19%	123	21%	138	23%	131	21%	121	18%
50-54	96	18%	88	15%	89	15%	105	17%	112	17%
55-64	74	14%	101	18%	121	20%	132	21%	153	23%
65+	16	3%	18	3%	21	3%	26	4%	32	5%
Race/Ethnicity										
White, not Hispanic	437	80%	461	80%	486	79%	502	79%	526	80%
Black/African American, not Hispanic	72	13%	78	14%	83	14%	87	14%	89	13%
Hispanic/Latino	14	3%	14	2%	14	2%	15	2%	16	2%
Asian/Pacific Islander	3	1%	3	1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	1	<1%	1	<1%
Unknown	20	4%	21	4%	26	4%	26	4%	26	4%
Race/Sex										
White, not Hispanic Males	354	65%	374	65%	395	65%	409	65%	430	65%
White, not Hispanic Females	83	15%	87	15%	91	15%	93	15%	96	15%
Black/African American, not Hispanic Males	55	10%	60	10%	64	10%	67	11%	69	10%
Black/African American, not Hispanic Females	17	3%	18	3%	19	3%	20	3%	20	3%
Hispanic/Latino Males	9	2%	9	2%	9	1%	9	1%	10	2%
Hispanic/Latina Females	5	1%	5	1%	5	1%	6	1%	6	1%
Asian/Pacific Islander Males	3	1%	3	1%	3	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	20	4%	21	4%	26	4%	26	4%	26	4%
Total	546		577		612		634		661	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Region 8D

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	283	67%	297	66%	319	67%	334	67%	348	67%
Injection drug use (IDU)	20	5%	21	5%	21	4%	21	4%	21	4%
Male-to-male sexual contact and IDU	21	5%	23	5%	23	5%	23	5%	25	5%
Heterosexual contact	30	7%	31	7%	34	7%	34	7%	35	7%
Other/unknown	70	17%	78	17%	81	17%	84	17%	91	18%
Subtotal	424	100%	450	100%	478	100%	496	100%	520	100%
Female adult or adolescent										
Injection drug use	8	8%	9	8%	9	8%	10	8%	11	9%
Heterosexual contact	72	68%	76	68%	80	68%	82	68%	84	68%
Other/unknown	26	25%	26	23%	28	24%	29	24%	29	23%
Subtotal	106	100%	111	100%	117	100%	121	100%	124	100%
Child (<13 yrs at diagnosis)										
Perinatal	12	75%	12	75%	13	76%	13	76%	13	76%
Other/unknown	4	25%	4	25%	4	24%	4	24%	4	24%
Subtotal	16	100%	16	100%	17	100%	17	100%	17	100%
Total	546		577		612		634		661	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Region 8D

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	272	50%	286	50%	307	50%	321	51%	334	51%
Injection drug use (IDU) only	19	3%	20	3%	20	3%	20	3%	20	3%
Heterosexual contact only	102	19%	107	19%	114	19%	116	18%	119	18%
MSM & IDU	16	3%	18	3%	18	3%	18	3%	20	3%
IDU & Heterosexual contact	9	2%	10	2%	10	2%	11	2%	12	2%
MSM & Heterosexual contact	11	2%	11	2%	12	2%	13	2%	14	2%
MSM & IDU & Heterosexual contact	5	1%	5	1%	5	1%	5	1%	5	1%
Perinatal exposure	13	2%	13	2%	14	2%	14	2%	14	2%
Other/unknown	99	18%	107	19%	112	18%	116	18%	123	19%
Total	546		577		612		634		661	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.