

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Hamilton

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Hamilton

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	30.7	119	80%	88	79%	24	83%	7	88%	
Females	6.9	29	20%	23	21%	5	17%	1	13%	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	15.2	8	5%	8	7%	-	-	-	-	
20-24	51.8	29	20%	23	21%	4	14%	2	25%	
25-29	57.3	36	24%	31	28%	5	17%	-	-	
30-34	32.7	18	12%	13	12%	4	14%	1	13%	
35-39	27.7	13	9%	11	10%	1	3%	1	13%	
40-44	25.4	12	8%	10	9%	1	3%	1	13%	
45-49	25.6	13	9%	5	5%	6	21%	2	25%	
50-54	10.4	6	4%	4	4%	2	7%	-	-	
55-64	11.3	12	8%	5	5%	6	21%	1	13%	
65+	*	1	1%	1	1%	-	-	-	-	
Race/Ethnicity										
White, not Hispanic	8.0	43	29%	33	30%	8	28%	2	25%	
Black/African American, not Hispanic	45.2	94	64%	67	60%	21	72%	6	75%	
Hispanic/Latino	25.6	6	4%	6	5%	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	5	3%	5	5%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	13.7	36	24%	26	23%	8	28%	2	25%	
White, not Hispanic Females	2.5	7	5%	7	6%	-	-	-	-	
Black/African American, not Hispanic Males	77.5	74	50%	53	48%	16	55%	5	63%	
Black/African American, not Hispanic Females	17.8	20	14%	14	13%	5	17%	1	13%	
Hispanic/Latino Males	41.2	5	3%	5	5%	-	-	-	-	
Hispanic/Latina Females	*	1	1%	1	1%	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	5	3%	5	5%	-	-	-	-	
Total	18.3	148		111		29		8		

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	73	61%	52	59%	17	71%	4	57%
Injection drug use (IDU)	5	4%	4	5%	1	4%	-	-
Male-to-male sexual contact and IDU	3	3%	3	3%	-	-	-	-
Heterosexual contact	7	6%	4	5%	1	4%	2	29%
Other/unknown	31	26%	25	28%	5	21%	1	14%
Subtotal	119	100%	88	100%	24	100%	7	100%
Female adult or adolescent								
Injection drug use	5	17%	5	22%	-	-	-	-
Heterosexual contact	15	52%	11	48%	4	80%	-	-
Other/unknown	9	31%	7	30%	1	20%	1	100%
Subtotal	29	100%	23	100%	5	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	148		111		29		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	68	46%	47	42%	17	59%	4	50%
Injection drug use (IDU) only	2	1%	1	1%	1	3%	-	-
Heterosexual contact only	22	15%	15	14%	5	17%	2	25%
MSM & IDU	3	2%	3	3%	-	-	-	-
IDU & Heterosexual contact	8	5%	8	7%	-	-	-	-
MSM & Heterosexual contact	5	3%	5	5%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	40	27%	32	29%	6	21%	2	25%
Total	148		111		29		8	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	23	64%	44	59%	4	80%	-	-	-	-	2	50%
Injection drug use (IDU)	4	11%	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	6%	-	-	-	-	-	-	-	-	1	25%
Heterosexual contact	-	-	7	9%	-	-	-	-	-	-	-	-
Other/unknown	7	19%	22	30%	1	20%	-	-	-	-	1	25%
Subtotal	36	100%	74	100%	5	100%	-	-	-	-	4	100%
Female adult or adolescent												
Injection drug use	5	71%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	14%	12	60%	1	100%	-	-	-	-	1	100%
Other/unknown	1	14%	8	40%	-	-	-	-	-	-	-	-
Subtotal	7	100%	20	100%	1	100%	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	43		94		6		0		0		5	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	23	53%	40	43%	3	50%	-	-	-	-	2	40%
Injection drug use (IDU) only	2	5%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	2%	19	20%	1	17%	-	-	-	-	1	20%
MSM & IDU	2	5%	-	-	-	-	-	-	-	-	1	20%
IDU & Heterosexual contact	7	16%	1	1%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	4	4%	1	17%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	8	19%	30	32%	1	17%	-	-	-	-	1	20%
Total	43		94		6		0		0		5	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Hamilton

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	30.1	117	76%	25.2	98	72%	27.8	108	73%	33.8	131	80%
Females	8.6	36	24%	9.1	38	28%	9.3	39	27%	7.9	33	20%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	1	1%	*	1	1%
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	32.4	17	11%	22.9	12	9%	24.8	13	9%	28.6	15	9%
20-24	60.7	34	22%	58.9	33	24%	66.0	37	25%	75.0	42	26%
25-29	43.0	27	18%	33.5	21	15%	30.3	19	13%	49.4	31	19%
30-34	29.1	16	10%	29.1	16	12%	38.1	21	14%	30.9	17	10%
35-39	29.8	14	9%	21.3	10	7%	21.3	10	7%	14.9	7	4%
40-44	33.9	16	10%	21.2	10	7%	23.3	11	7%	25.4	12	7%
45-49	39.4	20	13%	25.6	13	10%	33.5	17	12%	25.6	13	8%
50-54	13.9	8	5%	22.5	13	10%	12.1	7	5%	19.1	11	7%
55-64	*	1	1%	5.6	6	4%	8.5	9	6%	12.2	13	8%
65+	*	-	-	*	2	1%	*	2	1%	*	2	1%
Race/Ethnicity												
White, not Hispanic	8.4	45	29%	7.6	41	30%	6.1	33	22%	10.8	58	35%
Black/African American, not Hispanic	50.0	104	68%	43.3	90	66%	50.0	104	71%	47.1	98	60%
Hispanic/Latino	*	-	-	*	3	2%	*	4	3%	*	4	2%
Asian/Pacific Islander	*	-	-	*	2	1%	*	-	-	*	3	2%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	3%	*	-	-	*	6	4%	*	1	1%
Race/Sex												
White, not Hispanic Males	13.3	35	23%	12.6	33	24%	10.3	27	18%	18.7	49	30%
White, not Hispanic Females	3.6	10	7%	2.9	8	6%	2.2	6	4%	3.3	9	5%
Black/African American, not Hispanic Males	83.8	80	52%	65.0	62	46%	76.5	73	50%	78.6	75	46%
Black/African American, not Hispanic Females	21.3	24	16%	24.9	28	21%	27.5	31	21%	20.4	23	14%
Hispanic/Latino Males	*	-	-	*	2	1%	*	3	2%	*	3	2%
Hispanic/Latina Females	*	-	-	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Males	*	-	-	*	1	1%	*	-	-	*	3	2%
Asian/Pacific Islander Females	*	-	-	*	1	1%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	4	3%	*	-	-	*	6	4%	*	1	1%
Total	19.0	153		16.9	136		18.2	147		20.3	164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Hamilton

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	90	77%	64	65%	78	73%	93	72%
Injection drug use (IDU)	-	-	-	-	1	1%	5	4%
Male-to-male sexual contact and IDU	3	3%	2	2%	1	1%	6	5%
Heterosexual contact	10	9%	3	3%	2	2%	3	2%
Other/unknown	14	12%	29	30%	25	23%	23	18%
Subtotal	117	100%	98	100%	107	100%	130	100%
Female adult or adolescent								
Injection drug use	3	8%	1	3%	2	5%	5	16%
Heterosexual contact	29	81%	27	71%	30	79%	24	75%
Other/unknown	4	11%	10	26%	6	16%	3	9%
Subtotal	36	100%	38	100%	38	100%	32	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	1	50%
Other/unknown	-	-	-	-	2	100%	1	50%
Subtotal	-	-	-	-	2	100%	2	100%
Total	153		136		147		164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Hamilton

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	87	57%	60	44%	72	49%	91	55%
Injection drug use (IDU) only	2	1%	1	1%	1	1%	3	2%
Heterosexual contact only	39	25%	30	22%	32	22%	27	16%
MSM & IDU	3	2%	2	1%	-	-	5	3%
IDU & Heterosexual contact	1	1%	-	-	2	1%	7	4%
MSM & Heterosexual contact	3	2%	4	3%	6	4%	2	1%
MSM & IDU & Heterosexual contact	-	-	-	-	1	1%	1	1%
Perinatal exposure	-	-	-	-	1	1%	2	1%
Other/unknown	18	12%	39	29%	32	22%	26	16%
Total	153		136		147		164	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Hamilton

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	61	71%	62	69%	61	69%	85	77%	88	79%
Females	25	29%	28	31%	27	31%	25	23%	23	21%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	1	1%	1	1%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	16	19%	11	12%	12	14%	14	13%	8	7%
20-24	24	28%	24	27%	28	32%	34	31%	23	21%
25-29	13	15%	16	18%	11	13%	19	17%	31	28%
30-34	9	10%	8	9%	11	13%	13	12%	13	12%
35-39	7	8%	5	6%	4	5%	3	3%	11	10%
40-44	6	7%	7	8%	6	7%	6	5%	10	9%
45-49	8	9%	9	10%	10	11%	7	6%	5	5%
50-54	3	3%	6	7%	1	1%	5	5%	4	4%
55-64	-	-	3	3%	3	3%	7	6%	5	5%
65+	-	-	1	1%	1	1%	1	1%	1	1%
Race/Ethnicity										
White, not Hispanic	30	35%	28	31%	19	22%	38	35%	33	30%
Black/African American, not Hispanic	53	62%	59	66%	63	72%	67	61%	67	60%
Hispanic/Latino	-	-	1	1%	3	3%	3	3%	6	5%
Asian/Pacific Islander	-	-	2	2%	-	-	1	1%	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	-	-	3	3%	1	1%	5	5%
Race/Sex										
White, not Hispanic Males	22	26%	20	22%	13	15%	32	29%	26	23%
White, not Hispanic Females	8	9%	8	9%	6	7%	6	5%	7	6%
Black/African American, not Hispanic Males	38	44%	41	46%	43	49%	49	45%	53	48%
Black/African American, not Hispanic Females	15	17%	18	20%	20	23%	18	16%	14	13%
Hispanic/Latino Males	-	-	-	-	2	2%	2	2%	5	5%
Hispanic/Latina Females	-	-	1	1%	1	1%	1	1%	1	1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	1	1%	-	-
Asian/Pacific Islander Females	-	-	1	1%	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	3%	-	-	3	3%	1	1%	5	5%
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	49	80%	39	63%	43	72%	60	71%	52	59%
Injection drug use (IDU)	-	-	-	-	-	-	4	5%	4	5%
Male-to-male sexual contact and IDU	2	3%	1	2%	1	2%	5	6%	3	3%
Heterosexual contact	7	11%	3	5%	2	3%	2	2%	4	5%
Other/unknown	3	5%	19	31%	14	23%	14	16%	25	28%
Subtotal	61	100%	62	100%	60	100%	85	100%	88	100%
Female adult or adolescent										
Injection drug use	3	12%	1	4%	2	8%	4	17%	5	22%
Heterosexual contact	20	80%	19	68%	21	81%	17	71%	11	48%
Other/unknown	2	8%	8	29%	3	12%	3	13%	7	30%
Subtotal	25	100%	28	100%	26	100%	24	100%	23	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	-	-
Other/unknown	-	-	-	-	2	100%	-	-	-	-
Subtotal	-	-	-	-	2	100%	1	100%	-	-
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	48	56%	35	39%	38	43%	59	54%	47	42%
Injection drug use (IDU) only	2	2%	1	1%	1	1%	3	3%	1	1%
Heterosexual contact only	27	31%	22	24%	23	26%	19	17%	15	14%
MSM & IDU	2	2%	1	1%	-	-	5	5%	3	3%
IDU & Heterosexual contact	1	1%	-	-	1	1%	5	5%	8	7%
MSM & Heterosexual contact	1	1%	4	4%	5	6%	1	1%	5	5%
MSM & IDU & Heterosexual contact	-	-	-	-	1	1%	-	-	-	-
Perinatal exposure	-	-	-	-	1	1%	1	1%	-	-
Other/unknown	5	6%	27	30%	18	20%	17	15%	32	29%
Total	86		90		88		110		111	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Hamilton

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	70	81%	56	76%	56	81%	62	85%	47	78%	2,156	86%
Females	16	19%	18	24%	13	19%	11	15%	13	22%	350	14%
Age at AIDS diagnosis (yr)												
<13	-	-	1	1%	-	-	-	-	-	-	19	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	3	3%	5	7%	2	3%	1	1%	2	3%	60	2%
20-24	12	14%	18	24%	13	19%	12	16%	12	20%	240	10%
25-29	18	21%	7	9%	8	12%	19	26%	7	12%	443	18%
30-34	12	14%	11	15%	8	12%	7	10%	5	8%	522	21%
35-39	6	7%	8	11%	9	13%	5	7%	6	10%	447	18%
40-44	10	12%	6	8%	9	13%	6	8%	3	5%	328	13%
45-49	13	15%	7	9%	7	10%	8	11%	11	18%	205	8%
50-54	7	8%	5	7%	6	9%	7	10%	6	10%	122	5%
55-64	4	5%	5	7%	6	9%	7	10%	8	13%	94	4%
65+	1	1%	1	1%	1	1%	1	1%	-	-	24	1%
Race/Ethnicity												
White, not Hispanic	25	29%	21	28%	16	23%	22	30%	16	27%	1,144	46%
Black/African American, not Hispanic	59	69%	51	69%	47	68%	47	64%	42	70%	1,297	52%
Hispanic/Latino	-	-	1	1%	2	3%	1	1%	1	2%	29	1%
Asian/Pacific Islander	-	-	-	-	-	-	2	3%	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	2	2%	1	1%	4	6%	1	1%	1	2%	26	1%
Race/Sex												
White, not Hispanic Males	23	27%	17	23%	16	23%	19	26%	14	23%	1,072	43%
White, not Hispanic Females	2	2%	4	5%	-	-	3	4%	2	3%	72	3%
Black/African American, not Hispanic Males	45	52%	37	50%	35	51%	39	53%	31	52%	1,030	41%
Black/African American, not Hispanic Females	14	16%	14	19%	12	17%	8	11%	11	18%	267	11%
Hispanic/Latino Males	-	-	1	1%	2	3%	1	1%	1	2%	23	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	6	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	2	3%	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	2%	1	1%	4	6%	1	1%	1	2%	26	1%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	54	77%	39	70%	42	75%	47	77%	32	68%	1,576	74%
Injection drug use (IDU)	-	-	-	-	1	2%	1	2%	1	2%	91	4%
Male-to-male sexual contact and IDU	3	4%	1	2%	-	-	1	2%	1	2%	127	6%
Heterosexual contact	3	4%	1	2%	-	-	2	3%	5	11%	86	4%
Other/unknown	10	14%	15	27%	13	23%	10	16%	8	17%	262	12%
Subtotal	70	100%	56	100%	56	100%	61	100%	47	100%	2,142	100%
Female adult or adolescent												
Injection drug use	1	6%	2	12%	1	8%	-	-	1	8%	52	15%
Heterosexual contact	13	81%	13	76%	10	77%	11	100%	9	69%	249	72%
Other/unknown	2	13%	2	12%	2	15%	-	-	3	23%	43	13%
Subtotal	16	100%	17	100%	13	100%	11	100%	13	100%	344	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	14	70%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	6	30%
Subtotal	-	-	1	100%	-	-	1	100%	-	-	20	100%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	53	62%	39	53%	40	58%	46	63%	32	53%	1,508	60%
Injection drug use (IDU) only	-	-	2	3%	-	-	-	-	1	2%	85	3%
Heterosexual contact only	16	19%	14	19%	10	14%	13	18%	14	23%	335	13%
MSM & IDU	2	2%	1	1%	-	-	-	-	1	2%	112	4%
IDU & Heterosexual contact	1	1%	-	-	2	3%	1	1%	1	2%	58	2%
MSM & Heterosexual contact	1	1%	-	-	2	3%	1	1%	-	-	66	3%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	1	1%	-	-	15	1%
Perinatal exposure	-	-	1	1%	-	-	1	1%	-	-	15	1%
Other/unknown	12	14%	17	23%	15	22%	10	14%	11	18%	312	12%
Total	86		74		69		73		60		2,506	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Hamilton

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Hamilton

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	558.3	2,167	78%	1,039	76%	1,128	80%
Females	143.4	600	22%	326	24%	274	20%
Age at end of year							
<13	4.4	6	<1%	6	<1%	-	-
13-14	*	-	-	-	-	-	-
15-19	49.5	26	1%	24	2%	2	<1%
20-24	278.4	156	6%	126	9%	30	2%
25-29	468.3	294	11%	193	14%	101	7%
30-34	386.9	213	8%	111	8%	102	7%
35-39	540.9	254	9%	137	10%	117	8%
40-44	557.5	263	10%	128	9%	135	10%
45-49	843.8	428	15%	173	13%	255	18%
50-54	831.4	480	17%	220	16%	260	19%
55-64	486.7	517	19%	199	15%	318	23%
65+	113.8	130	5%	48	4%	82	6%
Race/Ethnicity							
White, not Hispanic	189.7	1,021	37%	515	38%	506	36%
Black/African American, not Hispanic	776.1	1,615	58%	783	57%	832	59%
Hispanic/Latino	256.1	60	2%	28	2%	32	2%
Asian/Pacific Islander	78.0	15	1%	8	1%	7	<1%
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Unknown	*	55	2%	31	2%	24	2%
Race/Sex							
White, not Hispanic Males	330.3	867	31%	418	31%	449	32%
White, not Hispanic Females	55.9	154	6%	97	7%	57	4%
Black/African American, not Hispanic Males	1,249.0	1,192	43%	568	42%	624	45%
Black/African American, not Hispanic Females	375.5	423	15%	215	16%	208	15%
Hispanic/Latino Males	395.7	48	2%	19	1%	29	2%
Hispanic/Latina Females	106.2	12	<1%	9	1%	3	<1%
Asian/Pacific Islander Males	147.6	14	1%	8	1%	6	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	55	2%	31	2%	24	2%
Total	343.0	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,496	69%	686	66%	810	72%
Injection drug use (IDU)	75	3%	31	3%	44	4%
Male-to-male sexual contact and IDU	98	5%	37	4%	61	5%
Heterosexual contact	119	6%	48	5%	71	6%
Other/unknown	368	17%	231	22%	137	12%
Subtotal	2,156	100%	1,033	100%	1,123	100%
Female adult or adolescent						
Injection drug use	70	12%	33	10%	37	14%
Heterosexual contact	417	71%	217	68%	200	74%
Other/unknown	100	17%	68	21%	32	12%
Subtotal	587	100%	318	100%	269	100%
Child (<13 yrs at diagnosis)						
Perinatal	17	71%	11	79%	6	60%
Other/unknown	7	29%	3	21%	4	40%
Subtotal	24	100%	14	100%	10	100%
Total	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	1,418	51%	652	48%	766	55%
Injection drug use (IDU) only	82	3%	31	2%	51	4%
Heterosexual contact only	536	19%	265	19%	271	19%
MSM & IDU	82	3%	31	2%	51	4%
IDU & Heterosexual contact	63	2%	33	2%	30	2%
MSM & Heterosexual contact	78	3%	34	2%	44	3%
MSM & IDU & Heterosexual contact	16	1%	6	<1%	10	1%
Perinatal exposure	19	1%	12	1%	7	<1%
Other/unknown	473	17%	301	22%	172	12%
Total	2,767		1,365		1,402	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Hamilton

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	689	80%	739	62%	25	52%	11	79%	1	100%	31	69%
Injection drug use (IDU)	19	2%	48	4%	7	15%	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	50	6%	44	4%	3	6%	-	-	-	-	1	2%
Heterosexual contact	11	1%	102	9%	4	8%	1	7%	-	-	1	2%
Other/unknown	95	11%	251	21%	9	19%	2	14%	-	-	11	24%
Subtotal	864	100%	1,184	100%	48	100%	14	100%	1	100%	45	100%
Female adult or adolescent												
Injection drug use	36	24%	30	7%	3	25%	-	-	-	-	1	10%
Heterosexual contact	90	60%	310	75%	9	75%	1	100%	-	-	7	70%
Other/unknown	25	17%	73	18%	-	-	-	-	-	-	2	20%
Subtotal	151	100%	413	100%	12	100%	1	100%	-	-	10	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	50%	14	78%	-	-	-	-	-	-	-	-
Other/unknown	3	50%	4	22%	-	-	-	-	-	-	-	-
Subtotal	6	100%	18	100%	-	-	-	-	-	-	-	-
Total	1,021		1,615		60		15		1		55	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Hamilton

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	666	65%	690	43%	21	35%	11	73%	1	100%	29	53%
Injection drug use (IDU) only	28	3%	46	3%	8	13%	-	-	-	-	-	-
Heterosexual contact only	101	10%	412	26%	13	22%	2	13%	-	-	8	15%
MSM & IDU	45	4%	34	2%	2	3%	-	-	-	-	1	2%
IDU & Heterosexual contact	27	3%	32	2%	2	3%	-	-	-	-	2	4%
MSM & Heterosexual contact	23	2%	49	3%	4	7%	-	-	-	-	2	4%
MSM & IDU & Heterosexual contact	5	0%	10	1%	1	2%	-	-	-	-	-	-
Perinatal exposure	3	0%	16	1%	-	-	-	-	-	-	-	-
Other/unknown	123	12%	326	20%	9	15%	2	13%	-	-	13	24%
Total	1,021		1,615		60		15		1		55	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Hamilton

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	1,700	79%	1,802	78%	1,916	78%	2,046	78%	2,167	78%
Females	460	21%	499	22%	537	22%	570	22%	600	22%
Age at end of year										
<13	3	<1%	4	<1%	5	<1%	6	<1%	6	<1%
13-14	1	<1%	-	-	1	<1%	1	<1%	-	-
15-19	32	1%	35	2%	30	1%	28	1%	26	1%
20-24	124	6%	150	7%	177	7%	180	7%	156	6%
25-29	152	7%	169	7%	191	8%	239	9%	294	11%
30-34	176	8%	181	8%	187	8%	196	7%	213	8%
35-39	216	10%	225	10%	228	9%	228	9%	254	9%
40-44	348	16%	309	13%	297	12%	276	11%	263	10%
45-49	429	20%	446	19%	454	19%	453	17%	428	15%
50-54	337	16%	379	16%	418	17%	456	17%	480	17%
55-64	276	13%	327	14%	372	15%	441	17%	517	19%
65+	66	3%	76	3%	93	4%	112	4%	130	5%
Race/Ethnicity										
White, not Hispanic	833	39%	879	38%	916	37%	975	37%	1,021	37%
Black/African American, not Hispanic	1,232	57%	1,323	57%	1,426	58%	1,521	58%	1,615	58%
Hispanic/Latino	44	2%	46	2%	50	2%	54	2%	60	2%
Asian/Pacific Islander	11	1%	12	1%	12	<1%	15	1%	15	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	39	2%	40	2%	48	2%	50	2%	55	2%
Race/Sex										
White, not Hispanic Males	709	33%	746	32%	777	32%	827	32%	867	31%
White, not Hispanic Females	124	6%	133	6%	139	6%	148	6%	154	6%
Black/African American, not Hispanic Males	912	42%	974	42%	1,047	43%	1,119	43%	1,192	43%
Black/African American, not Hispanic Females	320	15%	349	15%	379	15%	402	15%	423	15%
Hispanic/Latino Males	37	2%	38	2%	41	2%	44	2%	48	2%
Hispanic/Latina Females	7	<1%	8	<1%	9	<1%	10	<1%	12	<1%
Asian/Pacific Islander Males	10	<1%	11	<1%	11	<1%	14	1%	14	1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	39	2%	40	2%	48	2%	50	2%	55	2%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Hamilton

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	1,175	69%	1,245	69%	1,331	70%	1,424	70%	1,496	69%
Injection drug use (IDU)	63	4%	64	4%	66	3%	70	3%	75	3%
Male-to-male sexual contact and IDU	83	5%	86	5%	88	5%	94	5%	98	5%
Heterosexual contact	105	6%	107	6%	109	6%	112	6%	119	6%
Other/unknown	265	16%	291	16%	312	16%	335	16%	368	17%
Subtotal	1,691	100%	1,793	100%	1,906	100%	2,035	100%	2,156	100%
Female adult or adolescent										
Injection drug use	57	13%	58	12%	60	11%	65	12%	70	12%
Heterosexual contact	320	71%	348	71%	377	72%	401	72%	417	71%
Other/unknown	74	16%	83	17%	88	17%	91	16%	100	17%
Subtotal	451	100%	489	100%	525	100%	557	100%	587	100%
Child (<13 yrs at diagnosis)										
Perinatal	15	83%	16	84%	16	73%	17	71%	17	71%
Other/unknown	3	17%	3	16%	6	27%	7	29%	7	29%
Subtotal	18	100%	19	100%	22	100%	24	100%	24	100%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Hamilton

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	1,115	52%	1,182	51%	1,260	51%	1,351	52%	1,418	51%
Injection drug use (IDU) only	74	3%	75	3%	77	3%	80	3%	82	3%
Heterosexual contact only	425	20%	455	20%	486	20%	513	20%	536	19%
MSM & IDU	69	3%	72	3%	73	3%	78	3%	82	3%
IDU & Heterosexual contact	46	2%	47	2%	49	2%	55	2%	63	2%
MSM & Heterosexual contact	60	3%	63	3%	71	3%	73	3%	78	3%
MSM & IDU & Heterosexual contact	14	1%	14	1%	15	1%	16	1%	16	1%
Perinatal exposure	15	1%	16	1%	17	1%	19	1%	19	1%
Other/unknown	342	16%	377	16%	405	17%	431	16%	473	17%
Total	2,160		2,301		2,453		2,616		2,767	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.