Persons Living with a Diagnosis of HIV and/or AIDS Reported in HIV Prevention Planning Jefferson

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Jefferson

	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
				HIV (not AIDS)		AIDS	
Characteristic	**Rate[a]**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Sex							
Males	106.6	35	60%	16	59%	19	61%
Females	66.0	23	40%	11	41%	12	39%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	2%	1	4%	-	-
25-29	*	3	5%	1	4%	2	6%
30-34	*	1	2%	1	4%	-	-
35-39	*	4	7%	2	7%	2	6%
40-44	265.4	11	19%	4	15%	7	23%
45-49	142.9	6	10%	4	15%	2	6%
50-54	197.3	10	17%	2	7%	8	26%
55-64	139.7	15	26%	9	33%	6	19%
65+	52.9	7	12%	3	11%	4	13%
Race/Ethnicity							
White, not Hispanic	71.6	44	76%	20	74%	24	77%
Black/African American, not Hispanic	297.5	11	19%	5	19%	6	19%
Hispanic/Latino	*	2	3%	1	4%	1	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	1	2%	1	4%	-	-
Race/Sex							
White, not Hispanic Males	94.1	28	48%	12	44%	16	52%
White, not Hispanic Females	50.5	16	28%	8	30%	8	26%
Black/African American, not Hispanic Males	*	4	7%	2	7%	2	6%
Black/African American, not Hispanic Females	379.8	7	12%	3	11%	4	13%
Hispanic/Latino Males	*	2	3%	1	4%	1	3%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	1	2%	1	4%	-	-
Total	**85.7**	**58**		**27**		**31**	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

[a] The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Jefferson

	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
			HIV (not AIDS)		AIDS	
Transmission Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male adult or adolescent						
Male-to-male sexual contact	19	54%	8	50%	11	58%
Injection drug use (IDU)	4	11%	3	19%	1	5%
Male-to-male sexual contact and IDU	2	6%	1	6%	1	5%
Heterosexual contact	4	11%	1	6%	3	16%
Other/unknown	6	17%	3	19%	3	16%
Subtotal	35	100%	16	100%	19	100%
Female adult or adolescent						
Injection drug use	4	18%	3	27%	1	9%
Heterosexual contact	11	50%	3	27%	8	73%
Other/unknown	7	32%	5	45%	2	18%
Subtotal	22	100%	11	100%	11	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
Total	**58**		**27**		**31**	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

[a]Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Jefferson

	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
			HIV (not AIDS)		AIDS	
Exposure Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male-to-male sex (MSM) only	19	33%	8	30%	11	35%
Injection drug use (IDU) only	5	9%	5	19%	-	-
Heterosexual contact only	15	26%	4	15%	11	35%
MSM & IDU	2	3%	1	4%	1	3%
IDU & Heterosexual contact	3	5%	1	4%	2	6%
MSM & Heterosexual contact	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	2%	-	-	1	3%
Other/unknown	13	22%	8	30%	5	16%
Total	**58**		**27**		**31**	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

[a]Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Jefferson

	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/ Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
Transmission Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male adult or adolescent												
Male-to-male sexual contact	16	57%	1	25%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	3	11%	1	25%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	4%	1	25%	-	-	-	-	-	-	-	-
Heterosexual contact	4	14%	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	14%	1	25%	<10	*	-	-	-	-	-	-
Subtotal	28	100%	4	100%	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	3	20%	1	14%	-	-	-	-	-	-	-	-
Heterosexual contact	9	60%	2	29%	-	-	-	-	-	-	-	-
Other/unknown	3	20%	4	57%	-	-	-	-	-	-	-	-
Subtotal	15	100%	7	100%	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-	-	-
Total	**44**		**11**		**<10**		**0**		**0**		**<10**	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

[a]Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Jefferson

	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/ Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
Exposure Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male-to-male sex (MSM) only	16	36%	1	9%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	4	9%	1	9%	-	-	-	-	-	-	-	-
Heterosexual contact only	13	30%	2	18%	-	-	-	-	-	-	-	-
MSM & IDU	1	2%	1	9%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	5%	1	9%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	-	-	-	-	-	-	-	-	-	-
Other/unknown	7	16%	5	45%	<10	*	-	-	-	-	-	-
Total	**44**		**11**		**<10**		**0**		**0**		**<10**	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

[a]Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Jefferson

	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
Characteristic	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Sex										
Males	28	56%	30	58%	33	60%	34	60%	35	60%
Females	22	44%	22	42%	22	40%	23	40%	23	40%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	2	4%	2	4%	1	2%	2	4%	1	2%
25-29	-	-	-	-	2	4%	2	4%	3	5%
30-34	3	6%	2	4%	3	5%	2	4%	1	2%
35-39	9	18%	8	15%	7	13%	7	12%	4	7%
40-44	6	12%	8	15%	7	13%	8	14%	11	19%
45-49	9	18%	6	12%	8	15%	7	12%	6	10%
50-54	10	20%	13	25%	11	20%	9	16%	10	17%
55-64	10	20%	11	21%	11	20%	14	25%	15	26%
65+	1	2%	2	4%	5	9%	6	11%	7	12%
Race/Ethnicity										
White, not Hispanic	39	78%	40	77%	42	76%	44	77%	44	76%
Black/African American, not Hispanic	9	18%	10	19%	11	20%	11	19%	11	19%
Hispanic/Latino	1	2%	1	2%	1	2%	1	2%	2	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	2%	1	2%	1	2%
Race/Sex										
White, not Hispanic Males	24	48%	25	48%	27	49%	28	49%	28	48%
White, not Hispanic Females	15	30%	15	29%	15	27%	16	28%	16	28%
Black/African American, not Hispanic Males	2	4%	3	6%	4	7%	4	7%	4	7%
Black/African American, not Hispanic Females	7	14%	7	13%	7	13%	7	12%	7	12%
Hispanic/Latino Males	1	2%	1	2%	1	2%	1	2%	2	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	2%	1	2%	1	2%
Total	**50**		**52**		**55**		**57**		**58**	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Jefferson

	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
Transmission Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male adult or adolescent										
Male-to-male sexual contact	15	54%	15	50%	18	55%	19	56%	19	54%
Injection drug use (IDU)	4	14%	4	13%	4	12%	4	12%	4	11%
Male-to-male sexual contact and IDU	2	7%	2	7%	2	6%	2	6%	2	6%
Heterosexual contact	4	14%	4	13%	4	12%	4	12%	4	11%
Other/unknown	3	11%	5	17%	5	15%	5	15%	6	17%
Subtotal	28	100%	30	100%	33	100%	34	100%	35	100%
Female adult or adolescent										
Injection drug use	3	14%	3	14%	3	14%	4	18%	4	18%
Heterosexual contact	11	52%	11	52%	11	52%	11	50%	11	50%
Other/unknown	7	33%	7	33%	7	33%	7	32%	7	32%
Subtotal	21	100%	21	100%	21	100%	22	100%	22	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	100%	1	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	1	100%	1	100%	1	100%
Total	**50**		**52**		**55**		**57**		**58**	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

[a]Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Jefferson

	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
Exposure Category[a]	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**	**No.**	**%**
Male-to-male sex (MSM) only	15	30%	15	29%	18	33%	19	33%	19	33%
Injection drug use (IDU) only	5	10%	5	10%	5	9%	5	9%	5	9%
Heterosexual contact only	15	30%	15	29%	15	27%	15	26%	15	26%
MSM & IDU	2	4%	2	4%	2	4%	2	4%	2	3%
IDU & Heterosexual contact	2	4%	2	4%	2	4%	3	5%	3	5%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	1	2%	1	2%	1	2%	1	2%
Other/unknown	10	20%	12	23%	12	22%	12	21%	13	22%
Total	**50**		**52**		**55**		**57**		**58**	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

[a]Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.