

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Lawrence

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Lawrence

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	152.7	46	81%	23	82%	23	79%
Females	34.9	11	19%	5	18%	6	21%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	2%	1	4%	-	-
25-29	*	2	4%	2	7%	-	-
30-34	187.9	7	12%	6	21%	1	3%
35-39	*	4	7%	3	11%	1	3%
40-44	218.4	9	16%	3	11%	6	21%
45-49	209.1	9	16%	3	11%	6	21%
50-54	318.3	14	25%	5	18%	9	31%
55-64	93.9	8	14%	4	14%	4	14%
65+	*	3	5%	1	4%	2	7%
Race/Ethnicity							
White, not Hispanic	82.0	48	84%	25	89%	23	79%
Black/African American, not Hispanic	443.1	6	11%	2	7%	4	14%
Hispanic/Latino	*	1	2%	-	-	1	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	4%	1	4%	1	3%
Race/Sex							
White, not Hispanic Males	133.1	38	67%	20	71%	18	62%
White, not Hispanic Females	33.4	10	18%	5	18%	5	17%
Black/African American, not Hispanic Males	814.1	6	11%	2	7%	4	14%
Black/African American, not Hispanic Females	*	-	-	-	-	-	-
Hispanic/Latino Males	*	1	2%	-	-	1	3%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	4%	1	4%	1	3%
Total	92.5	57		28		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Lawrence

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	31	67%	14	61%	17	74%
Injection drug use (IDU)	3	7%	2	9%	1	4%
Male-to-male sexual contact and IDU	6	13%	3	13%	3	13%
Heterosexual contact	3	7%	2	9%	1	4%
Other/unknown	3	7%	2	9%	1	4%
Subtotal	46	100%	23	100%	23	100%
Female adult or adolescent						
Injection drug use	1	9%	1	20%	-	-
Heterosexual contact	7	64%	2	40%	5	83%
Other/unknown	3	27%	2	40%	1	17%
Subtotal	11	100%	5	100%	6	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	57		28		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Lawrence

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	31	54%	14	50%	17	59%
Injection drug use (IDU) only	3	5%	2	7%	1	3%
Heterosexual contact only	10	18%	4	14%	6	21%
MSM & IDU	4	7%	3	11%	1	3%
IDU & Heterosexual contact	1	2%	1	4%	-	-
MSM & Heterosexual contact	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	2	4%	-	-	2	7%
Perinatal exposure	-	-	-	-	-	-
Other/unknown	6	11%	4	14%	2	7%
Total	57		28		29	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Lawrence

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	26	68%	<10	*	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	2	5%	<10	*	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	11%	<10	*	-	-	-	-	-	-	-	-
Heterosexual contact	3	8%	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	8%	-	-	-	-	-	-	-	-	-	-
Subtotal	38	100%	<10	*	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	1	10%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	6	60%	-	-	-	-	-	-	-	-	<10	*
Other/unknown	3	30%	-	-	-	-	-	-	-	-	-	-
Subtotal	10	100%	-	-	-	-	-	-	-	-	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	48		<10		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Lawrence

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	26	54%	<10	*	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	2	4%	<10	*	-	-	-	-	-	-	-	-
Heterosexual contact only	9	19%	-	-	-	-	-	-	-	-	<10	*
MSM & IDU	3	6%	<10	*	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	2%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	2%	<10	*	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	13%	-	-	-	-	-	-	-	-	-	-
Total	48		<10		<10		0		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Lawrence

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	43	80%	45	80%	46	81%	46	81%	46	81%
Females	11	20%	11	20%	11	19%	11	19%	11	19%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	1	2%	2	4%	1	2%	1	2%	1	2%
25-29	7	13%	7	13%	4	7%	2	4%	2	4%
30-34	3	6%	2	4%	6	11%	7	12%	7	12%
35-39	8	15%	9	16%	7	12%	6	11%	4	7%
40-44	9	17%	7	13%	9	16%	9	16%	9	16%
45-49	12	22%	10	18%	10	18%	10	18%	9	16%
50-54	8	15%	10	18%	11	19%	13	23%	14	25%
55-64	5	9%	7	13%	7	12%	7	12%	8	14%
65+	1	2%	2	4%	2	4%	2	4%	3	5%
Race/Ethnicity										
White, not Hispanic	45	83%	47	84%	48	84%	48	84%	48	84%
Black/African American, not Hispanic	6	11%	6	11%	6	11%	6	11%	6	11%
Hispanic/Latino	1	2%	1	2%	1	2%	1	2%	1	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	4%	2	4%	2	4%	2	4%	2	4%
Race/Sex										
White, not Hispanic Males	35	65%	37	66%	38	67%	38	67%	38	67%
White, not Hispanic Females	10	19%	10	18%	10	18%	10	18%	10	18%
Black/African American, not Hispanic Males	6	11%	6	11%	6	11%	6	11%	6	11%
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino Males	1	2%	1	2%	1	2%	1	2%	1	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	4%	2	4%	2	4%	2	4%	2	4%
Total	54		56		57		57		57	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Lawrence

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	29	67%	30	67%	31	67%	31	67%	31	67%
Injection drug use (IDU)	3	7%	3	7%	3	7%	3	7%	3	7%
Male-to-male sexual contact and IDU	6	14%	6	13%	6	13%	6	13%	6	13%
Heterosexual contact	3	7%	3	7%	3	7%	3	7%	3	7%
Other/unknown	2	5%	3	7%	3	7%	3	7%	3	7%
Subtotal	43	100%	45	100%	46	100%	46	100%	46	100%
Female adult or adolescent										
Injection drug use	1	9%	1	9%	1	9%	1	9%	1	9%
Heterosexual contact	7	64%	7	64%	7	64%	7	64%	7	64%
Other/unknown	3	27%	3	27%	3	27%	3	27%	3	27%
Subtotal	11	100%	11	100%	11	100%	11	100%	11	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	54		56		57		57		57	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Lawrence

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	29	54%	30	54%	31	54%	31	54%	31	54%
Injection drug use (IDU) only	3	6%	3	5%	3	5%	3	5%	3	5%
Heterosexual contact only	10	19%	10	18%	10	18%	10	18%	10	18%
MSM & IDU	4	7%	4	7%	4	7%	4	7%	4	7%
IDU & Heterosexual contact	1	2%	1	2%	1	2%	1	2%	1	2%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	2	4%	2	4%	2	4%	2	4%	2	4%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	9%	6	11%	6	11%	6	11%	6	11%
Total	54		56		57		57		57	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.