

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Licking

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Licking

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	153.0	127	80%	61	86%	66	75%
Females	37.0	32	20%	10	14%	22	25%
Age at end of year							
<13	*	2	1%	-	-	2	2%
13-14	*	-	-	-	-	-	-
15-19	*	4	3%	4	6%	-	-
20-24	84.1	9	6%	8	11%	1	1%
25-29	65.0	6	4%	3	4%	3	3%
30-34	119.6	12	8%	10	14%	2	2%
35-39	122.0	12	8%	5	7%	7	8%
40-44	193.3	22	14%	6	8%	16	18%
45-49	247.8	29	18%	11	15%	18	20%
50-54	175.5	23	14%	5	7%	18	20%
55-64	139.9	32	20%	15	21%	17	19%
65+	31.0	8	5%	4	6%	4	5%
Race/Ethnicity							
White, not Hispanic	81.3	126	79%	55	77%	71	81%
Black/African American, not Hispanic	284.5	18	11%	8	11%	10	11%
Hispanic/Latino	208.0	6	4%	3	4%	3	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	1%	1	1%	-	-
Unknown	*	8	5%	4	6%	4	5%
Race/Sex							
White, not Hispanic Males	138.3	105	66%	48	68%	57	65%
White, not Hispanic Females	26.5	21	13%	7	10%	14	16%
Black/African American, not Hispanic Males	398.0	13	8%	7	10%	6	7%
Black/African American, not Hispanic Females	163.4	5	3%	1	1%	4	5%
Hispanic/Latino Males	*	3	2%	2	3%	1	1%
Hispanic/Latina Females	*	3	2%	1	1%	2	2%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	1	1%	1	1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	8	5%	4	6%	4	5%
Total	93.9	159		71		88	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Licking

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	83	67%	38	66%	45	69%
Injection drug use (IDU)	7	6%	1	2%	6	9%
Male-to-male sexual contact and IDU	6	5%	3	5%	3	5%
Heterosexual contact	7	6%	3	5%	4	6%
Other/unknown	20	16%	13	22%	7	11%
Subtotal	123	100%	58	100%	65	100%
Female adult or adolescent						
Injection drug use	3	10%	1	11%	2	10%
Heterosexual contact	25	83%	6	67%	19	90%
Other/unknown	2	7%	2	22%	-	-
Subtotal	30	100%	9	100%	21	100%
Child (<13 yrs at diagnosis)						
Perinatal	5	83%	3	75%	2	100%
Other/unknown	1	17%	1	25%	-	-
Subtotal	6	100%	4	100%	2	100%
Total	159		71		88	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Licking

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	79	50%	35	49%	44	50%
Injection drug use (IDU) only	6	4%	-	-	6	7%
Heterosexual contact only	32	20%	9	13%	23	26%
MSM & IDU	6	4%	3	4%	3	3%
IDU & Heterosexual contact	4	3%	2	3%	2	2%
MSM & Heterosexual contact	4	3%	3	4%	1	1%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	5	3%	3	4%	2	2%
Other/unknown	23	14%	16	23%	7	8%
Total	159		71		88	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Licking

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	73	71%	5	45%	<10	*	-	-	<10	*	<10	*
Injection drug use (IDU)	4	4%	2	18%	<10	*	-	-	-	-	-	-
Male-to-male sexual contact and IDU	5	5%	1	9%	-	-	-	-	-	-	-	-
Heterosexual contact	4	4%	1	9%	<10	*	-	-	-	-	<10	*
Other/unknown	17	17%	2	18%	-	-	-	-	-	-	<10	*
Subtotal	103	100%	11	100%	<10	*	-	-	<10	*	<10	*
Female adult or adolescent												
Injection drug use	2	10%	-	-	<10	*	-	-	-	-	-	-
Heterosexual contact	18	86%	4	100%	<10	*	-	-	-	-	<10	*
Other/unknown	1	5%	-	-	-	-	-	-	-	-	<10	*
Subtotal	21	100%	4	100%	<10	*	-	-	-	-	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	1	50%	3	100%	<10	*	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	3	100%	<10	*	-	-	-	-	-	-
Total	126		18		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Licking

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	70	56%	4	22%	<10	*	-	-	<10	*	<10	*
Injection drug use (IDU) only	3	2%	1	6%	<10	*	-	-	-	-	-	-
Heterosexual contact only	22	17%	5	28%	<10	*	-	-	-	-	<10	*
MSM & IDU	5	4%	1	6%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	3	2%	1	6%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	3	2%	1	6%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	3	17%	<10	*	-	-	-	-	-	-
Other/unknown	19	15%	2	11%	-	-	-	-	-	-	<10	*
Total	126		18		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Licking

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	105	80%	109	80%	119	80%	122	80%	127	80%
Females	27	20%	27	20%	30	20%	31	20%	32	20%
Age at end of year										
<13	1	1%	1	1%	2	1%	2	1%	2	1%
13-14	2	2%	2	1%	-	-	-	-	-	-
15-19	1	1%	1	1%	3	2%	3	2%	4	3%
20-24	4	3%	4	3%	10	7%	12	8%	9	6%
25-29	7	5%	6	4%	5	3%	3	2%	6	4%
30-34	8	6%	10	7%	10	7%	12	8%	12	8%
35-39	17	13%	11	8%	10	7%	8	5%	12	8%
40-44	22	17%	26	19%	26	17%	23	15%	22	14%
45-49	26	20%	26	19%	25	17%	30	20%	29	18%
50-54	21	16%	22	16%	27	18%	27	18%	23	14%
55-64	18	14%	21	15%	25	17%	26	17%	32	20%
65+	5	4%	6	4%	6	4%	7	5%	8	5%
Race/Ethnicity										
White, not Hispanic	106	80%	110	81%	119	80%	121	79%	126	79%
Black/African American, not Hispanic	15	11%	15	11%	17	11%	18	12%	18	11%
Hispanic/Latino	5	4%	5	4%	5	3%	5	3%	6	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	1	1%	1	1%
Unknown	6	5%	6	4%	8	5%	8	5%	8	5%
Race/Sex										
White, not Hispanic Males	88	67%	92	68%	100	67%	101	66%	105	66%
White, not Hispanic Females	18	14%	18	13%	19	13%	20	13%	21	13%
Black/African American, not Hispanic Males	11	8%	11	8%	12	8%	13	8%	13	8%
Black/African American, not Hispanic Females	4	3%	4	3%	5	3%	5	3%	5	3%
Hispanic/Latino Males	2	2%	2	1%	2	1%	2	1%	3	2%
Hispanic/Latina Females	3	2%	3	2%	3	2%	3	2%	3	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	1	1%	1	1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	6	5%	6	4%	8	5%	8	5%	8	5%
Total	132		136		149		153		159	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Licking

Transmission Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	66	65%	68	65%	77	67%	79	67%	83	67%
Injection drug use (IDU)	7	7%	7	7%	7	6%	7	6%	7	6%
Male-to-male sexual contact and IDU	6	6%	6	6%	6	5%	6	5%	6	5%
Heterosexual contact	5	5%	6	6%	7	6%	7	6%	7	6%
Other/unknown	17	17%	18	17%	18	16%	19	16%	20	16%
Subtotal	101	100%	105	100%	115	100%	118	100%	123	100%
Female adult or adolescent										
Injection drug use	2	8%	2	8%	2	7%	3	10%	3	10%
Heterosexual contact	22	85%	22	85%	24	86%	24	83%	25	83%
Other/unknown	2	8%	2	8%	2	7%	2	7%	2	7%
Subtotal	26	100%	26	100%	28	100%	29	100%	30	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	5	83%	5	83%	5	83%
Other/unknown	1	20%	1	20%	1	17%	1	17%	1	17%
Subtotal	5	100%	5	100%	6	100%	6	100%	6	100%
Total	132		136		149		153		159	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Licking

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	63	48%	65	48%	73	49%	75	49%	79	50%
Injection drug use (IDU) only	6	5%	6	4%	6	4%	6	4%	6	4%
Heterosexual contact only	27	20%	28	21%	31	21%	31	20%	32	20%
MSM & IDU	6	5%	6	4%	6	4%	6	4%	6	4%
IDU & Heterosexual contact	3	2%	3	2%	3	2%	4	3%	4	3%
MSM & Heterosexual contact	3	2%	3	2%	4	3%	4	3%	4	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	4	3%	4	3%	5	3%	5	3%	5	3%
Other/unknown	20	15%	21	15%	21	14%	22	14%	23	14%
Total	132		136		149		153		159	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.