

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Lorain

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Lorain

| Characteristic | 2014 diagnosis of HIV infection | | | Disease Status at Initial Diagnosis | | | | | |
|--|---------------------------------|----------|-----|-------------------------------------|-----|------------------|---|----------|------|
| | Rate ^a | No. | % | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | | | | No. | % | No. | % | No. | % |
| Sex | | | | | | | | | |
| Males | 4.0 | 6 | 75% | 5 | 71% | - | - | 1 | 100% |
| Females | * | 2 | 25% | 2 | 29% | - | - | - | - |
| Age at diagnosis (yr) | | | | | | | | | |
| <13 | * | - | - | - | - | - | - | - | - |
| 13-14 | * | - | - | - | - | - | - | - | - |
| 15-19 | * | 1 | 13% | 1 | 14% | - | - | - | - |
| 20-24 | * | 1 | 13% | 1 | 14% | - | - | - | - |
| 25-29 | * | 2 | 25% | 1 | 14% | - | - | 1 | 100% |
| 30-34 | * | 1 | 13% | 1 | 14% | - | - | - | - |
| 35-39 | * | 2 | 25% | 2 | 29% | - | - | - | - |
| 40-44 | * | 1 | 13% | 1 | 14% | - | - | - | - |
| 45-49 | * | - | - | - | - | - | - | - | - |
| 50-54 | * | - | - | - | - | - | - | - | - |
| 55-64 | * | - | - | - | - | - | - | - | - |
| 65+ | * | - | - | - | - | - | - | - | - |
| Race/Ethnicity | | | | | | | | | |
| White, not Hispanic | * | 2 | 25% | 2 | 29% | - | - | - | - |
| Black/African American, not Hispanic | * | 2 | 25% | 1 | 14% | - | - | 1 | 100% |
| Hispanic/Latino | * | 4 | 50% | 4 | 57% | - | - | - | - |
| Asian/Pacific Islander | * | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native | * | - | - | - | - | - | - | - | - |
| Unknown | * | - | - | - | - | - | - | - | - |
| Race/Sex | | | | | | | | | |
| White, not Hispanic Males | * | 1 | 13% | 1 | 14% | - | - | - | - |
| White, not Hispanic Females | * | 1 | 13% | 1 | 14% | - | - | - | - |
| Black/African American, not Hispanic Males | * | 1 | 13% | - | - | - | - | 1 | 100% |
| Black/African American, not Hispanic Females | * | 1 | 13% | 1 | 14% | - | - | - | - |
| Hispanic/Latino Males | * | 4 | 50% | 4 | 57% | - | - | - | - |
| Hispanic/Latina Females | * | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Males | * | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Females | * | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Males | * | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Females | * | - | - | - | - | - | - | - | - |
| Unknown | * | - | - | - | - | - | - | - | - |
| Total | 2.6 | 8 | | 7 | | - | | 1 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | 2014 diagnosis of HIV infection | | Disease Status at Initial Diagnosis | | | | | |
|--|---------------------------------|------|-------------------------------------|------|------------------|---|----------|------|
| | | | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | |
| Male-to-male sexual contact | 2 | 33% | 2 | 40% | - | - | - | - |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | - | - | - | - | - | - |
| Heterosexual contact | 1 | 17% | - | - | - | - | 1 | 100% |
| Other/unknown | 3 | 50% | 3 | 60% | - | - | - | - |
| Subtotal | 6 | 100% | 5 | 100% | - | - | 1 | 100% |
| Female adult or adolescent | | | | | | | | |
| Injection drug use | - | - | - | - | - | - | - | - |
| Heterosexual contact | 1 | 50% | 1 | 50% | - | - | - | - |
| Other/unknown | 1 | 50% | 1 | 50% | - | - | - | - |
| Subtotal | 2 | 100% | 2 | 100% | - | - | - | - |
| Child (<13 yrs at diagnosis) | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - |
| Total | 8 | | 7 | | 0 | | 1 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | 2014 diagnosis of HIV infection | | Disease Status at Initial Diagnosis | | | | | |
|----------------------------------|---------------------------------|-----|-------------------------------------|-----|------------------|---|----------|------|
| | | | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 2 | 25% | 2 | 29% | - | - | - | - |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 2 | 25% | 1 | 14% | - | - | 1 | 100% |
| MSM & IDU | - | - | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - |
| Other/unknown | 4 | 50% | 4 | 57% | - | - | - | - |
| Total | 8 | | 7 | | - | | 1 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | 2014 Diagnosis of HIV Infection | | | | | | | | | | | |
|--|---------------------------------|------|------------------------|------|-----------------|------|------------------------|---|-------------------------------|---|----------|---|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 1 | 100% | - | - | 1 | 25% | - | - | - | - | - | - |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | - | - | 1 | 100% | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | 3 | 75% | - | - | - | - | - | - |
| Subtotal | 1 | 100% | 1 | 100% | 4 | 100% | - | - | - | - | - | - |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact | 1 | 100% | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | 1 | 100% | - | - | - | - | - | - | - | - |
| Subtotal | 1 | 100% | 1 | 100% | - | - | - | - | - | - | - | - |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - | - | - | - | - |
| Total | 2 | | 2 | | 4 | | 0 | | 0 | | 0 | |

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | 2014 Diagnosis of HIV Infection | | | | | | | | | | | |
|----------------------------------|---------------------------------|-----|------------------------|-----|-----------------|-----|------------------------|---|-------------------------------|---|----------|---|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 1 | 50% | - | - | 1 | 25% | - | - | - | - | - | - |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 1 | 50% | 1 | 50% | - | - | - | - | - | - | - | - |
| MSM & IDU | - | - | - | - | - | - | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | 1 | 50% | 3 | 75% | - | - | - | - | - | - |
| Total | 2 | | 2 | | 4 | | 0 | | 0 | | 0 | |

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Lorain

Diagnosis of HIV Infection by Year

| Characteristic | 2010 | | | 2011 | | | 2012 | | | 2013 | | |
|--|-------------------|-----------|-----|-------------------|-----------|-----|-------------------|-----------|-----|-------------------|-----------|-----|
| | Rate ^a | No. | % |
| Sex | | | | | | | | | | | | |
| Males | 8.7 | 13 | 87% | 8.0 | 12 | 75% | 8.7 | 13 | 76% | 7.3 | 11 | 73% |
| Females | * | 2 | 13% | * | 4 | 25% | * | 4 | 24% | * | 4 | 27% |
| Age at diagnosis (yr) | | | | | | | | | | | | |
| <13 | * | - | - | * | - | - | * | - | - | * | - | - |
| 13-14 | * | - | - | * | 0 | <1% | * | - | - | * | - | - |
| 15-19 | * | - | - | * | 1 | 6% | * | 3 | 18% | * | - | - |
| 20-24 | * | 3 | 20% | * | 2 | 13% | 31.9 | 6 | 35% | * | 2 | 13% |
| 25-29 | * | 3 | 20% | * | 2 | 13% | * | 2 | 12% | * | 1 | 7% |
| 30-34 | * | 3 | 20% | * | 1 | 6% | * | - | - | * | 3 | 20% |
| 35-39 | * | 3 | 20% | * | 2 | 13% | * | 1 | 6% | * | 2 | 13% |
| 40-44 | * | - | - | * | 2 | 13% | * | 1 | 6% | * | 2 | 13% |
| 45-49 | * | 2 | 13% | * | 2 | 13% | * | - | - | * | 4 | 27% |
| 50-54 | * | 1 | 7% | * | 3 | 19% | * | 2 | 12% | * | - | - |
| 55-64 | * | - | - | * | 1 | 6% | * | 2 | 12% | * | 1 | 7% |
| 65+ | * | - | - | * | - | - | * | - | - | * | - | - |
| Race/Ethnicity | | | | | | | | | | | | |
| White, not Hispanic | 4.1 | 10 | 67% | 2.9 | 7 | 44% | 3.7 | 9 | 53% | 3.3 | 8 | 53% |
| Black/African American, not Hispanic | * | 3 | 20% | 25.0 | 6 | 38% | 25.0 | 6 | 35% | * | 3 | 20% |
| Hispanic/Latino | * | 2 | 13% | * | 2 | 13% | * | 2 | 12% | * | 4 | 27% |
| Asian/Pacific Islander | * | - | - | * | - | - | * | - | - | * | - | - |
| American Indian/Alaska Native | * | - | - | * | - | - | * | - | - | * | - | - |
| Unknown | * | - | - | * | 1 | 6% | * | - | - | * | - | - |
| Race/Sex | | | | | | | | | | | | |
| White, not Hispanic Males | 7.6 | 9 | 60% | * | 4 | 25% | 6.8 | 8 | 47% | 5.9 | 7 | 47% |
| White, not Hispanic Females | * | 1 | 7% | * | 3 | 19% | * | 1 | 6% | * | 1 | 7% |
| Black/African American, not Hispanic Males | * | 3 | 20% | 47.1 | 6 | 38% | * | 3 | 18% | * | 1 | 7% |
| Black/African American, not Hispanic Females | * | - | - | * | - | - | * | 3 | 18% | * | 2 | 13% |
| Hispanic/Latino Males | * | 1 | 7% | * | 1 | 6% | * | 2 | 12% | * | 3 | 20% |
| Hispanic/Latina Females | * | 1 | 7% | * | 1 | 6% | * | - | - | * | 1 | 7% |
| Asian/Pacific Islander Males | * | - | - | * | - | - | * | - | - | * | - | - |
| Asian/Pacific Islander Females | * | - | - | * | - | - | * | - | - | * | - | - |
| American Indian/Alaska Native Males | * | - | - | * | - | - | * | - | - | * | - | - |
| American Indian/Alaska Native Females | * | - | - | * | - | - | * | - | - | * | - | - |
| Unknown | * | - | - | * | 1 | 6% | * | - | - | * | - | - |
| Total | 4.9 | 15 | | 5.3 | 16 | | 5.6 | 17 | | 4.9 | 15 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Lorain

Diagnosis of HIV Infection by Year

| Transmission Category ^a | 2010 | | 2011 | | 2012 | | 2013 | |
|--|-----------|------|-----------|------|-----------|------|-----------|------|
| | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | |
| Male-to-male sexual contact | 9 | 69% | 4 | 33% | 11 | 85% | 9 | 82% |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | 1 | 8% | 1 | 8% | - | - | - | - |
| Heterosexual contact | - | - | - | - | - | - | 1 | 9% |
| Other/unknown | 3 | 23% | 7 | 58% | 2 | 15% | 1 | 9% |
| Subtotal | 13 | 100% | 12 | 100% | 13 | 100% | 11 | 100% |
| Female adult or adolescent | | | | | | | | |
| Injection drug use | - | - | 2 | 50% | - | - | - | - |
| Heterosexual contact | 2 | 100% | 1 | 25% | 1 | 25% | 3 | 75% |
| Other/unknown | - | - | 1 | 25% | 3 | 75% | 1 | 25% |
| Subtotal | 2 | 100% | 4 | 100% | 4 | 100% | 4 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - |
| Total | 15 | | 16 | | 17 | | 15 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Lorain

Diagnosis of HIV Infection by Year

| Exposure Category ^a | 2010 | | 2011 | | 2012 | | 2013 | |
|----------------------------------|-----------|-----|-----------|-----|-----------|-----|-----------|-----|
| | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 9 | 60% | 4 | 25% | 11 | 65% | 9 | 60% |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 2 | 13% | 1 | 6% | 1 | 6% | 4 | 27% |
| MSM & IDU | - | - | 1 | 6% | - | - | - | - |
| IDU & Heterosexual contact | - | - | 2 | 13% | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | 1 | 7% | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - |
| Other/unknown | 3 | 20% | 8 | 50% | 5 | 29% | 2 | 13% |
| Total | 15 | | 16 | | 17 | | 15 | |

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Lorain

| Characteristic | Year of HIV (not AIDS) Diagnosis | | | | | | | | | |
|--|----------------------------------|-----|-----------|-----|----------|-----|-----------|-----|----------|-----|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Sex | | | | | | | | | | |
| Males | 7 | 78% | 8 | 80% | 6 | 86% | 7 | 64% | 5 | 71% |
| Females | 2 | 22% | 2 | 20% | 1 | 14% | 4 | 36% | 2 | 29% |
| Age at HIV diagnosis (yr) | | | | | | | | | | |
| <13 | - | - | - | - | - | - | - | - | - | - |
| 13-14 | - | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | 1 | 10% | 2 | 29% | - | - | 1 | 14% |
| 20-24 | 2 | 22% | 2 | 20% | 3 | 43% | 2 | 18% | 1 | 14% |
| 25-29 | 2 | 22% | - | - | 1 | 14% | 1 | 9% | 1 | 14% |
| 30-34 | 1 | 11% | 1 | 10% | - | - | 3 | 27% | 1 | 14% |
| 35-39 | 1 | 11% | 2 | 20% | - | - | 1 | 9% | 2 | 29% |
| 40-44 | - | - | 1 | 10% | - | - | 1 | 9% | 1 | 14% |
| 45-49 | 2 | 22% | 1 | 10% | - | - | 3 | 27% | - | - |
| 50-54 | 1 | 11% | 1 | 10% | 1 | 14% | - | - | - | - |
| 55-64 | - | - | 1 | 10% | - | - | - | - | - | - |
| 65+ | - | - | - | - | - | - | - | - | - | - |
| Race/Ethnicity | | | | | | | | | | |
| White, not Hispanic | 6 | 67% | 5 | 50% | 2 | 29% | 6 | 55% | 2 | 29% |
| Black/African American, not Hispanic | 1 | 11% | 4 | 40% | 3 | 43% | 3 | 27% | 1 | 14% |
| Hispanic/Latino | 2 | 22% | - | - | 2 | 29% | 2 | 18% | 4 | 57% |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | 1 | 10% | - | - | - | - | - | - |
| Race/Sex | | | | | | | | | | |
| White, not Hispanic Males | 5 | 56% | 3 | 30% | 2 | 29% | 5 | 45% | 1 | 14% |
| White, not Hispanic Females | 1 | 11% | 2 | 20% | - | - | 1 | 9% | 1 | 14% |
| Black/African American, not Hispanic Males | 1 | 11% | 4 | 40% | 2 | 29% | 1 | 9% | - | - |
| Black/African American, not Hispanic Females | - | - | - | - | 1 | 14% | 2 | 18% | 1 | 14% |
| Hispanic/Latino Males | 1 | 11% | - | - | 2 | 29% | 1 | 9% | 4 | 57% |
| Hispanic/Latina Females | 1 | 11% | - | - | - | - | 1 | 9% | - | - |
| Asian/Pacific Islander Males | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Females | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Males | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Females | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | 1 | 10% | - | - | - | - | - | - |
| Total | 9 | | 10 | | 7 | | 11 | | 7 | |

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | Year of HIV (not AIDS) Diagnosis | | | | | | | | | |
|--|----------------------------------|------|-----------|------|----------|------|-----------|------|----------|------|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | |
| Male-to-male sexual contact | 4 | 57% | 3 | 38% | 6 | 100% | 6 | 86% | 2 | 40% |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | - | - | 1 | 13% | - | - | - | - | - | - |
| Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | 3 | 43% | 4 | 50% | - | - | 1 | 14% | 3 | 60% |
| Subtotal | 7 | 100% | 8 | 100% | 6 | 100% | 7 | 100% | 5 | 100% |
| Female adult or adolescent | | | | | | | | | | |
| Injection drug use | - | - | 1 | 50% | - | - | - | - | - | - |
| Heterosexual contact | 2 | 100% | 1 | 50% | 1 | 100% | 3 | 75% | 1 | 50% |
| Other/unknown | - | - | - | - | - | - | 1 | 25% | 1 | 50% |
| Subtotal | 2 | 100% | 2 | 100% | 1 | 100% | 4 | 100% | 2 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | - | - | - | - | - | - |
| Total | 9 | | 10 | | 7 | | 11 | | 7 | |

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | Year of HIV (not AIDS) Diagnosis | | | | | | | | | |
|----------------------------------|----------------------------------|-----|-----------|-----|----------|-----|-----------|-----|----------|-----|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 4 | 44% | 3 | 30% | 6 | 86% | 6 | 55% | 2 | 29% |
| Injection drug use (IDU) only | - | - | - | - | - | - | - | - | - | - |
| Heterosexual contact only | 2 | 22% | 1 | 10% | 1 | 14% | 3 | 27% | 1 | 14% |
| MSM & IDU | - | - | 1 | 10% | - | - | - | - | - | - |
| IDU & Heterosexual contact | - | - | 1 | 10% | - | - | - | - | - | - |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | - | - | - | - | - | - | - | - | - | - |
| Perinatal exposure | - | - | - | - | - | - | - | - | - | - |
| Other/unknown | 3 | 33% | 4 | 40% | - | - | 2 | 18% | 4 | 57% |
| Total | 9 | | 10 | | 7 | | 11 | | 7 | |

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Lorain

| Characteristic | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|--|------------------------|-----|-----------|-----|----------|-----|----------|-----|----------|-----|-----------------|-----|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Sex | | | | | | | | | | | | |
| Males | 5 | 83% | 7 | 70% | 4 | 50% | 5 | 83% | 3 | 75% | 216 | 76% |
| Females | 1 | 17% | 3 | 30% | 4 | 50% | 1 | 17% | 1 | 25% | 70 | 24% |
| Age at AIDS diagnosis (yr) | | | | | | | | | | | | |
| <13 | - | - | - | - | - | - | - | - | - | - | 5 | 2% |
| 13-14 | - | - | - | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | - | - | 2 | 25% | - | - | - | - | 8 | 3% |
| 20-24 | 2 | 33% | 3 | 30% | 2 | 25% | - | - | 1 | 25% | 31 | 11% |
| 25-29 | - | - | 3 | 30% | - | - | 1 | 17% | 1 | 25% | 52 | 18% |
| 30-34 | 2 | 33% | - | - | - | - | - | - | - | - | 44 | 15% |
| 35-39 | 1 | 17% | 1 | 10% | 2 | 25% | 1 | 17% | - | - | 54 | 19% |
| 40-44 | 1 | 17% | 1 | 10% | - | - | 1 | 17% | 1 | 25% | 34 | 12% |
| 45-49 | - | - | - | - | - | - | 1 | 17% | 1 | 25% | 34 | 12% |
| 50-54 | - | - | 2 | 20% | 1 | 13% | - | - | - | - | 11 | 4% |
| 55-64 | - | - | - | - | 1 | 13% | 2 | 33% | - | - | 12 | 4% |
| 65+ | - | - | - | - | - | - | - | - | - | - | 1 | <1% |
| Race/Ethnicity | | | | | | | | | | | | |
| White, not Hispanic | 3 | 50% | 4 | 40% | 5 | 63% | 5 | 83% | 1 | 25% | 155 | 54% |
| Black/African American, not Hispanic | 2 | 33% | 4 | 40% | 2 | 25% | - | - | 2 | 50% | 63 | 22% |
| Hispanic/Latino | 1 | 17% | 2 | 20% | 1 | 13% | 1 | 17% | 1 | 25% | 63 | 22% |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | - | 3 | 1% |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - | 2 | 1% |
| Race/Sex | | | | | | | | | | | | |
| White, not Hispanic Males | 2 | 33% | 3 | 30% | 3 | 38% | 4 | 67% | 1 | 25% | 128 | 45% |
| White, not Hispanic Females | 1 | 17% | 1 | 10% | 2 | 25% | 1 | 17% | - | - | 27 | 9% |
| Black/African American, not Hispanic Males | 2 | 33% | 3 | 30% | 1 | 13% | - | - | 1 | 25% | 48 | 17% |
| Black/African American, not Hispanic Females | - | - | 1 | 10% | 1 | 13% | - | - | 1 | 25% | 15 | 5% |
| Hispanic/Latino Males | 1 | 17% | 1 | 10% | - | - | 1 | 17% | 1 | 25% | 38 | 13% |
| Hispanic/Latina Females | - | - | 1 | 10% | 1 | 13% | - | - | - | - | 25 | 9% |
| Asian/Pacific Islander Males | - | - | - | - | - | - | - | - | - | - | 2 | 1% |
| Asian/Pacific Islander Females | - | - | - | - | - | - | - | - | - | - | 1 | <1% |
| American Indian/Alaska Native Males | - | - | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Females | - | - | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - | 2 | 1% |
| Total | 6 | | 10 | | 8 | | 6 | | 4 | | 286 | |

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|--|------------------------|------|-----------|------|----------|------|----------|------|----------|------|-----------------|------|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 4 | 80% | 2 | 29% | 3 | 75% | 3 | 60% | 2 | 67% | 129 | 61% |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - | 25 | 12% |
| Male-to-male sexual contact and IDU | - | - | 1 | 14% | - | - | - | - | - | - | 12 | 6% |
| Heterosexual contact | 1 | 20% | - | - | - | - | 1 | 20% | 1 | 33% | 14 | 7% |
| Other/unknown | - | - | 4 | 57% | 1 | 25% | 1 | 20% | - | - | 33 | 15% |
| Subtotal | 5 | 100% | 7 | 100% | 4 | 100% | 5 | 100% | 3 | 100% | 213 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | 1 | 25% | 1 | 100% | - | - | 14 | 21% |
| Heterosexual contact | 1 | 100% | 1 | 33% | 1 | 25% | - | - | - | - | 41 | 60% |
| Other/unknown | - | - | 2 | 67% | 2 | 50% | - | - | 1 | 100% | 13 | 19% |
| Subtotal | 1 | 100% | 3 | 100% | 4 | 100% | 1 | 100% | 1 | 100% | 68 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | - | - | - | - | - | - | - | - | - | - | 3 | 60% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | 2 | 40% |
| Subtotal | - | - | - | - | - | - | - | - | - | - | 5 | 100% |
| Total | 6 | | 10 | | 8 | | 6 | | 4 | | 286 | |

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|----------------------------------|------------------------|-----|-----------|-----|----------|-----|----------|-----|----------|-----|-----------------|-----|
| | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male-to-male sex (MSM) only | 4 | 67% | 2 | 20% | 3 | 38% | 3 | 50% | 2 | 50% | 124 | 43% |
| Injection drug use (IDU) only | - | - | - | - | 1 | 13% | - | - | - | - | 25 | 9% |
| Heterosexual contact only | 2 | 33% | 1 | 10% | 1 | 13% | 1 | 17% | 1 | 25% | 55 | 19% |
| MSM & IDU | - | - | - | - | - | - | - | - | - | - | 10 | 3% |
| IDU & Heterosexual contact | - | - | - | - | - | - | 1 | 17% | - | - | 14 | 5% |
| MSM & Heterosexual contact | - | - | - | - | - | - | - | - | - | - | 5 | 2% |
| MSM & IDU & Heterosexual contact | - | - | 1 | 10% | - | - | - | - | - | - | 2 | 1% |
| Perinatal exposure | - | - | - | - | - | - | - | - | - | - | 3 | 1% |
| Other/unknown | - | - | 6 | 60% | 3 | 38% | 1 | 17% | 1 | 25% | 48 | 17% |
| Total | 6 | | 10 | | 8 | | 6 | | 4 | | 286 | |

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Lorain

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Lorain

| Characteristic | Living with a diagnosis of HIV infection in 2014 | | | Current Disease Status | | | |
|--|--|------------|-----|------------------------|-----|------------|-----|
| | Rate ^a | No. | % | HIV (not AIDS) | | AIDS | |
| | | | | No. | % | No. | % |
| Sex | | | | | | | |
| Males | 170.1 | 255 | 74% | 118 | 72% | 137 | 75% |
| Females | 59.0 | 91 | 26% | 46 | 28% | 45 | 25% |
| Age at end of year | | | | | | | |
| <13 | * | 1 | <1% | 1 | 1% | - | - |
| 13-14 | * | 1 | <1% | - | - | 1 | 1% |
| 15-19 | * | 2 | 1% | 1 | 1% | 1 | 1% |
| 20-24 | 42.5 | 8 | 2% | 8 | 5% | - | - |
| 25-29 | 149.1 | 24 | 7% | 15 | 9% | 9 | 5% |
| 30-34 | 127.6 | 22 | 6% | 14 | 9% | 8 | 4% |
| 35-39 | 165.4 | 30 | 9% | 15 | 9% | 15 | 8% |
| 40-44 | 260.9 | 52 | 15% | 25 | 15% | 27 | 15% |
| 45-49 | 300.9 | 63 | 18% | 30 | 18% | 33 | 18% |
| 50-54 | 276.3 | 64 | 18% | 22 | 13% | 42 | 23% |
| 55-64 | 132.8 | 57 | 16% | 19 | 12% | 38 | 21% |
| 65+ | 44.3 | 22 | 6% | 14 | 9% | 8 | 4% |
| Race/Ethnicity | | | | | | | |
| White, not Hispanic | 68.5 | 165 | 48% | 87 | 53% | 78 | 43% |
| Black/African American, not Hispanic | 346.2 | 83 | 24% | 30 | 18% | 53 | 29% |
| Hispanic/Latino | 323.5 | 91 | 26% | 44 | 27% | 47 | 26% |
| Asian/Pacific Islander | * | 2 | 1% | 1 | 1% | 1 | 1% |
| American Indian/Alaska Native | * | - | - | - | - | - | - |
| Unknown | * | 5 | 1% | 2 | 1% | 3 | 2% |
| Race/Sex | | | | | | | |
| White, not Hispanic Males | 109.5 | 129 | 37% | 66 | 40% | 63 | 35% |
| White, not Hispanic Females | 29.2 | 36 | 10% | 21 | 13% | 15 | 8% |
| Black/African American, not Hispanic Males | 478.6 | 61 | 18% | 21 | 13% | 40 | 22% |
| Black/African American, not Hispanic Females | 195.9 | 22 | 6% | 9 | 5% | 13 | 7% |
| Hispanic/Latino Males | 423.9 | 60 | 17% | 29 | 18% | 31 | 17% |
| Hispanic/Latina Females | 221.8 | 31 | 9% | 15 | 9% | 16 | 9% |
| Asian/Pacific Islander Males | * | 2 | 1% | 1 | 1% | 1 | 1% |
| Asian/Pacific Islander Females | * | - | - | - | - | - | - |
| American Indian/Alaska Native Males | * | - | - | - | - | - | - |
| American Indian/Alaska Native Females | * | - | - | - | - | - | - |
| Unknown | * | 5 | 1% | 2 | 1% | 3 | 2% |
| Total | 113.7 | 346 | | 164 | | 182 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | Living with a diagnosis of HIV infection in 2014 | | Current Disease Status | | | |
|--|--|------|------------------------|------|------------|------|
| | No. | % | HIV (not AIDS) | | AIDS | |
| | | | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 150 | 59% | 73 | 62% | 77 | 57% |
| Injection drug use (IDU) | 22 | 9% | 11 | 9% | 11 | 8% |
| Male-to-male sexual contact and IDU | 11 | 4% | 1 | 1% | 10 | 7% |
| Heterosexual contact | 20 | 8% | 4 | 3% | 16 | 12% |
| Other/unknown | 50 | 20% | 28 | 24% | 22 | 16% |
| Subtotal | 253 | 100% | 117 | 100% | 136 | 100% |
| Female adult or adolescent | | | | | | |
| Injection drug use | 10 | 11% | 6 | 14% | 4 | 9% |
| Heterosexual contact | 59 | 67% | 29 | 66% | 30 | 68% |
| Other/unknown | 19 | 22% | 9 | 20% | 10 | 23% |
| Subtotal | 88 | 100% | 44 | 100% | 44 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | 5 | 100% | 3 | 100% | 2 | 100% |
| Other/unknown | - | - | - | - | - | - |
| Subtotal | 5 | 100% | 3 | 100% | 2 | 100% |
| Total | 346 | | 164 | | 182 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | Living with a diagnosis of HIV infection in 2014 | | Current Disease Status | | | |
|----------------------------------|--|-----|------------------------|-----|------------|-----|
| | No. | % | HIV (not AIDS) | | AIDS | |
| | | | No. | % | No. | % |
| Male-to-male sex (MSM) only | 143 | 41% | 71 | 43% | 72 | 40% |
| Injection drug use (IDU) only | 20 | 6% | 11 | 7% | 9 | 5% |
| Heterosexual contact only | 79 | 23% | 33 | 20% | 46 | 25% |
| MSM & IDU | 7 | 2% | 1 | 1% | 6 | 3% |
| IDU & Heterosexual contact | 12 | 3% | 6 | 4% | 6 | 3% |
| MSM & Heterosexual contact | 7 | 2% | 2 | 1% | 5 | 3% |
| MSM & IDU & Heterosexual contact | 4 | 1% | - | - | 4 | 2% |
| Perinatal exposure | 6 | 2% | 3 | 2% | 3 | 2% |
| Other/unknown | 68 | 20% | 37 | 23% | 31 | 17% |
| Total | 346 | | 164 | | 182 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Lorain

| Transmission Category ^a | Living with a Diagnosis of HIV Infection in 2014 | | | | | | | | | | | |
|--|--|------|------------------------|------|-----------------|------|------------------------|------|-------------------------------|---|----------|------|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 86 | 67% | 33 | 56% | 29 | 48% | 1 | 50% | - | - | 1 | 33% |
| Injection drug use (IDU) | 5 | 4% | 7 | 12% | 10 | 17% | - | - | - | - | - | - |
| Male-to-male sexual contact and IDU | 6 | 5% | 1 | 2% | 4 | 7% | - | - | - | - | - | - |
| Heterosexual contact | 6 | 5% | 6 | 10% | 7 | 12% | - | - | - | - | 1 | 33% |
| Other/unknown | 26 | 20% | 12 | 20% | 10 | 17% | 1 | 50% | - | - | 1 | 33% |
| Subtotal | 129 | 100% | 59 | 100% | 60 | 100% | 2 | 100% | - | - | 3 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | 6 | 18% | - | - | 4 | 13% | - | - | - | - | - | - |
| Heterosexual contact | 23 | 68% | 16 | 73% | 19 | 61% | - | - | - | - | 1 | 100% |
| Other/unknown | 5 | 15% | 6 | 27% | 8 | 26% | - | - | - | - | - | - |
| Subtotal | 34 | 100% | 22 | 100% | 31 | 100% | - | - | - | - | 1 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | 2 | 100% | 2 | 100% | - | - | - | - | - | - | 1 | 100% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 2 | 100% | 2 | 100% | - | - | - | - | - | - | 1 | 100% |
| Total | 165 | | 83 | | 91 | | 2 | | 0 | | 5 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Lorain

| Exposure Category ^a | Living with a Diagnosis of HIV Infection in 2014 | | | | | | | | | | | |
|----------------------------------|--|-----|------------------------|-----|-----------------|-----|------------------------|-----|-------------------------------|---|----------|-----|
| | White, not Hispanic | | Black/African American | | Hispanic/Latino | | Asian/Pacific Islander | | American Indian/Alaska Native | | Unknown | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sex (MSM) only | 82 | 50% | 31 | 37% | 28 | 31% | 1 | 50% | - | - | 1 | 20% |
| Injection drug use (IDU) only | 6 | 4% | 4 | 5% | 10 | 11% | - | - | - | - | - | - |
| Heterosexual contact only | 29 | 18% | 22 | 27% | 26 | 29% | - | - | - | - | 2 | 40% |
| MSM & IDU | 4 | 2% | 1 | 1% | 2 | 2% | - | - | - | - | - | - |
| IDU & Heterosexual contact | 5 | 3% | 3 | 4% | 4 | 4% | - | - | - | - | - | - |
| MSM & Heterosexual contact | 4 | 2% | 2 | 2% | 1 | 1% | - | - | - | - | - | - |
| MSM & IDU & Heterosexual contact | 2 | 1% | - | - | 2 | 2% | - | - | - | - | - | - |
| Perinatal exposure | 2 | 1% | 3 | 4% | - | - | - | - | - | - | 1 | 20% |
| Other/unknown | 31 | 19% | 17 | 20% | 18 | 20% | 1 | 50% | - | - | 1 | 20% |
| Total | 165 | | 83 | | 91 | | 2 | | 0 | | 5 | |

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Lorain

Living with a Diagnosis of HIV Infection

| Characteristic | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|--|------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|
| | No. | % |
| Sex | | | | | | | | | | |
| Males | 218 | 74% | 225 | 74% | 240 | 74% | 251 | 74% | 255 | 74% |
| Females | 78 | 26% | 81 | 26% | 85 | 26% | 89 | 26% | 91 | 26% |
| Age at end of year | | | | | | | | | | |
| <13 | 3 | 1% | 2 | 1% | 2 | 1% | 2 | 1% | 1 | <1% |
| 13-14 | 1 | <1% | 2 | 1% | 1 | <1% | - | - | 1 | <1% |
| 15-19 | - | - | - | - | 1 | <1% | 2 | 1% | 2 | 1% |
| 20-24 | 12 | 4% | 13 | 4% | 15 | 5% | 12 | 4% | 8 | 2% |
| 25-29 | 18 | 6% | 17 | 6% | 21 | 6% | 20 | 6% | 24 | 7% |
| 30-34 | 22 | 7% | 23 | 8% | 23 | 7% | 24 | 7% | 22 | 6% |
| 35-39 | 42 | 14% | 35 | 11% | 30 | 9% | 33 | 10% | 30 | 9% |
| 40-44 | 51 | 17% | 60 | 20% | 53 | 16% | 51 | 15% | 52 | 15% |
| 45-49 | 68 | 23% | 55 | 18% | 62 | 19% | 65 | 19% | 63 | 18% |
| 50-54 | 37 | 13% | 52 | 17% | 62 | 19% | 67 | 20% | 64 | 18% |
| 55-64 | 32 | 11% | 33 | 11% | 37 | 11% | 44 | 13% | 57 | 16% |
| 65+ | 10 | 3% | 14 | 5% | 18 | 6% | 20 | 6% | 22 | 6% |
| Race/Ethnicity | | | | | | | | | | |
| White, not Hispanic | 140 | 47% | 145 | 47% | 155 | 48% | 164 | 48% | 165 | 48% |
| Black/African American, not Hispanic | 69 | 23% | 72 | 24% | 79 | 24% | 81 | 24% | 83 | 24% |
| Hispanic/Latino | 80 | 27% | 82 | 27% | 84 | 26% | 88 | 26% | 91 | 26% |
| Asian/Pacific Islander | 2 | 1% | 2 | 1% | 2 | 1% | 2 | 1% | 2 | 1% |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - |
| Unknown | 5 | 2% | 5 | 2% | 5 | 2% | 5 | 1% | 5 | 1% |
| Race/Sex | | | | | | | | | | |
| White, not Hispanic Males | 109 | 37% | 112 | 37% | 121 | 37% | 129 | 38% | 129 | 37% |
| White, not Hispanic Females | 31 | 10% | 33 | 11% | 34 | 10% | 35 | 10% | 36 | 10% |
| Black/African American, not Hispanic Males | 53 | 18% | 56 | 18% | 60 | 18% | 60 | 18% | 61 | 18% |
| Black/African American, not Hispanic Females | 16 | 5% | 16 | 5% | 19 | 6% | 21 | 6% | 22 | 6% |
| Hispanic/Latino Males | 51 | 17% | 52 | 17% | 54 | 17% | 57 | 17% | 60 | 17% |
| Hispanic/Latina Females | 29 | 10% | 30 | 10% | 30 | 9% | 31 | 9% | 31 | 9% |
| Asian/Pacific Islander Males | 2 | 1% | 2 | 1% | 2 | 1% | 2 | 1% | 2 | 1% |
| Asian/Pacific Islander Females | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Males | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Females | - | - | - | - | - | - | - | - | - | - |
| Unknown | 5 | 2% | 5 | 2% | 5 | 2% | 5 | 1% | 5 | 1% |
| Total | 296 | | 306 | | 325 | | 340 | | 346 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Lorain

Living with a Diagnosis of HIV Infection

| Transmission Category ^a | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|--|------------|------|------------|------|------------|------|------------|------|------------|------|
| | No. | % |
| Male adult or adolescent | | | | | | | | | | |
| Male-to-male sexual contact | 130 | 60% | 132 | 59% | 142 | 60% | 149 | 60% | 150 | 59% |
| Injection drug use (IDU) | 21 | 10% | 21 | 9% | 21 | 9% | 22 | 9% | 22 | 9% |
| Male-to-male sexual contact and IDU | 10 | 5% | 10 | 4% | 11 | 5% | 11 | 4% | 11 | 4% |
| Heterosexual contact | 17 | 8% | 17 | 8% | 18 | 8% | 19 | 8% | 20 | 8% |
| Other/unknown | 38 | 18% | 43 | 19% | 46 | 19% | 48 | 19% | 50 | 20% |
| Subtotal | 216 | 100% | 223 | 100% | 238 | 100% | 249 | 100% | 253 | 100% |
| Female adult or adolescent | | | | | | | | | | |
| Injection drug use | 9 | 12% | 10 | 13% | 10 | 12% | 10 | 12% | 10 | 11% |
| Heterosexual contact | 53 | 71% | 54 | 69% | 55 | 67% | 58 | 67% | 59 | 67% |
| Other/unknown | 13 | 17% | 14 | 18% | 17 | 21% | 18 | 21% | 19 | 22% |
| Subtotal | 75 | 100% | 78 | 100% | 82 | 100% | 86 | 100% | 88 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | |
| Perinatal | 5 | 100% | 5 | 100% | 5 | 100% | 5 | 100% | 5 | 100% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - |
| Subtotal | 5 | 100% | 5 | 100% | 5 | 100% | 5 | 100% | 5 | 100% |
| Total | 296 | | 306 | | 325 | | 340 | | 346 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Lorain

Living with a Diagnosis of HIV Infection

| Exposure Category ^a | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | |
|----------------------------------|------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|
| | No. | % |
| Male-to-male sex (MSM) only | 123 | 42% | 125 | 41% | 135 | 42% | 142 | 42% | 143 | 41% |
| Injection drug use (IDU) only | 19 | 6% | 19 | 6% | 19 | 6% | 20 | 6% | 20 | 6% |
| Heterosexual contact only | 70 | 24% | 71 | 23% | 73 | 22% | 77 | 23% | 79 | 23% |
| MSM & IDU | 7 | 2% | 7 | 2% | 7 | 2% | 7 | 2% | 7 | 2% |
| IDU & Heterosexual contact | 11 | 4% | 12 | 4% | 12 | 4% | 12 | 4% | 12 | 3% |
| MSM & Heterosexual contact | 7 | 2% | 7 | 2% | 7 | 2% | 7 | 2% | 7 | 2% |
| MSM & IDU & Heterosexual contact | 3 | 1% | 3 | 1% | 4 | 1% | 4 | 1% | 4 | 1% |
| Perinatal exposure | 6 | 2% | 6 | 2% | 6 | 2% | 6 | 2% | 6 | 2% |
| Other/unknown | 50 | 17% | 56 | 18% | 62 | 19% | 65 | 19% | 68 | 20% |
| Total | 296 | | 306 | | 325 | | 340 | | 346 | |

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.