

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Madison

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Madison**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	171.2	41	91%	16	89%	25	93%
Females	*	4	9%	2	11%	2	7%
<b>Age at end of year</b>							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	2	4%	2	11%	-	-
25-29	*	3	7%	1	6%	2	7%
30-34	*	4	9%	3	17%	1	4%
35-39	162.1	5	11%	2	11%	3	11%
40-44	205.9	7	16%	4	22%	3	11%
45-49	237.2	8	18%	-	-	8	30%
50-54	*	4	9%	-	-	4	15%
55-64	123.2	7	16%	3	17%	4	15%
65+	82.0	5	11%	3	17%	2	7%
<b>Race/Ethnicity</b>							
White, not Hispanic	69.1	27	60%	14	78%	13	48%
Black/African American, not Hispanic	488.3	14	31%	4	22%	10	37%
Hispanic/Latino	*	2	4%	-	-	2	7%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	4%	-	-	2	7%
<b>Race/Sex</b>							
White, not Hispanic Males	112.8	23	51%	12	67%	11	41%
White, not Hispanic Females	*	4	9%	2	11%	2	7%
Black/African American, not Hispanic Males	555.6	14	31%	4	22%	10	37%
Black/African American, not Hispanic Females	*	-	-	-	-	-	-
Hispanic/Latino Males	*	2	4%	-	-	2	7%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	4%	-	-	2	7%
<b>Total</b>	<b>102.5</b>	<b>45</b>		<b>18</b>		<b>27</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Madison**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	16	39%	5	31%	11	44%
Injection drug use (IDU)	3	7%	1	6%	2	8%
Male-to-male sexual contact and IDU	2	5%	-	-	2	8%
Heterosexual contact	6	15%	2	13%	4	16%
Other/unknown	14	34%	8	50%	6	24%
Subtotal	41	100%	16	100%	25	100%
<b>Female adult or adolescent</b>						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	3	75%	1	50%	2	100%
Other/unknown	1	25%	1	50%	-	-
Subtotal	4	100%	2	100%	2	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>45</b>		<b>18</b>		<b>27</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Madison**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	15	33%	5	28%	10	37%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	9	20%	3	17%	6	22%
MSM & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	3	7%	1	6%	2	7%
MSM & Heterosexual contact	1	2%	-	-	1	4%
MSM & IDU & Heterosexual contact	2	4%	-	-	2	7%
Perinatal exposure	-	-	-	-	-	-
Other/unknown	15	33%	9	50%	6	22%
<b>Total</b>	<b>45</b>		<b>18</b>		<b>27</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Madison**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	13	57%	2	14%	-	-	-	-	-	-	<10	*
Injection drug use (IDU)	1	4%	-	-	<10	*	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	2	14%	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	6	43%	-	-	-	-	-	-	-	-
Other/unknown	9	39%	4	29%	-	-	-	-	-	-	<10	*
Subtotal	23	100%	14	100%	<10	*	-	-	-	-	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	75%	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	25%	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	-	-	-	-	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>27</b>		<b>14</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Madison**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	13	48%	2	14%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	11%	6	43%	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	4%	-	-	<10	*	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	<10	*
MSM & IDU & Heterosexual contact	-	-	2	14%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	10	37%	4	29%	-	-	-	-	-	-	<10	*
<b>Total</b>	<b>27</b>		<b>14</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Madison**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	38	90%	39	91%	41	91%	41	91%	41	91%
Females	4	10%	4	9%	4	9%	4	9%	4	9%
<b>Age at end of year</b>										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	2	5%	1	2%	1	2%	-	-	-	-
20-24	2	5%	3	7%	3	7%	4	9%	2	4%
25-29	5	12%	3	7%	1	2%	1	2%	3	7%
30-34	4	10%	7	16%	7	16%	7	16%	4	9%
35-39	4	10%	4	9%	5	11%	2	4%	5	11%
40-44	8	19%	6	14%	8	18%	8	18%	7	16%
45-49	4	10%	5	12%	6	13%	7	16%	8	18%
50-54	5	12%	5	12%	4	9%	5	11%	4	9%
55-64	4	10%	5	12%	6	13%	6	13%	7	16%
65+	4	10%	4	9%	4	9%	5	11%	5	11%
<b>Race/Ethnicity</b>										
White, not Hispanic	25	60%	26	60%	27	60%	27	60%	27	60%
Black/African American, not Hispanic	14	33%	14	33%	14	31%	14	31%	14	31%
Hispanic/Latino	2	5%	2	5%	2	4%	2	4%	2	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	2	4%	2	4%	2	4%
<b>Race/Sex</b>										
White, not Hispanic Males	21	50%	22	51%	23	51%	23	51%	23	51%
White, not Hispanic Females	4	10%	4	9%	4	9%	4	9%	4	9%
Black/African American, not Hispanic Males	14	33%	14	33%	14	31%	14	31%	14	31%
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino Males	2	5%	2	5%	2	4%	2	4%	2	4%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	2	4%	2	4%	2	4%
<b>Total</b>	<b>42</b>		<b>43</b>		<b>45</b>		<b>45</b>		<b>45</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Madison

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	14	37%	15	38%	16	39%	16	39%	16	39%
Injection drug use (IDU)	3	8%	3	8%	3	7%	3	7%	3	7%
Male-to-male sexual contact and IDU	2	5%	2	5%	2	5%	2	5%	2	5%
Heterosexual contact	6	16%	6	15%	6	15%	6	15%	6	15%
Other/unknown	13	34%	13	33%	14	34%	14	34%	14	34%
Subtotal	38	100%	39	100%	41	100%	41	100%	41	100%
<b>Female adult or adolescent</b>										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	75%	3	75%	3	75%	3	75%	3	75%
Other/unknown	1	25%	1	25%	1	25%	1	25%	1	25%
Subtotal	4	100%	4	100%	4	100%	4	100%	4	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>42</b>		<b>43</b>		<b>45</b>		<b>45</b>		<b>45</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Madison

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	14	33%	15	35%	15	33%	15	33%	15	33%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	9	21%	9	21%	9	20%	9	20%	9	20%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	3	7%	3	7%	3	7%	3	7%	3	7%
MSM & Heterosexual contact	-	-	-	-	1	2%	1	2%	1	2%
MSM & IDU & Heterosexual contact	2	5%	2	5%	2	4%	2	4%	2	4%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	14	33%	14	33%	15	33%	15	33%	15	33%
<b>Total</b>	<b>42</b>		<b>43</b>		<b>45</b>		<b>45</b>		<b>45</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.