

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Montgomery

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Montgomery

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	17.9	46	78%	36	80%	8	73%	2	67%
Females	4.7	13	22%	9	20%	3	27%	1	33%
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	3	5%	3	7%	-	-	-	-
20-24	48.6	18	31%	16	36%	2	18%	-	-
25-29	25.2	9	15%	8	18%	1	9%	-	-
30-34	*	2	3%	1	2%	1	9%	-	-
35-39	36.9	11	19%	7	16%	2	18%	2	67%
40-44	22.1	7	12%	5	11%	1	9%	1	33%
45-49	*	3	5%	1	2%	2	18%	-	-
50-54	*	2	3%	2	4%	-	-	-	-
55-64	*	1	2%	-	-	1	9%	-	-
65+	*	3	5%	2	4%	1	9%	-	-
Race/Ethnicity									
White, not Hispanic	5.5	21	36%	16	36%	3	27%	2	67%
Black/African American, not Hispanic	32.4	36	61%	27	60%	8	73%	1	33%
Hispanic/Latino	*	2	3%	2	4%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Race/Sex									
White, not Hispanic Males	10.7	20	34%	15	33%	3	27%	2	67%
White, not Hispanic Females	*	1	2%	1	2%	-	-	-	-
Black/African American, not Hispanic Males	48.4	25	42%	20	44%	5	45%	-	-
Black/African American, not Hispanic Females	18.5	11	19%	7	16%	3	27%	1	33%
Hispanic/Latino Males	*	1	2%	1	2%	-	-	-	-
Hispanic/Latina Females	*	1	2%	1	2%	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Total	11.1	59		45		11		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	39	85%	31	86%	7	88%	1	50%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	2%	1	3%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	6	13%	4	11%	1	13%	1	50%
Subtotal	46	100%	36	100%	8	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	6	46%	5	56%	1	33%	-	-
Other/unknown	7	54%	4	44%	2	67%	1	100%
Subtotal	13	100%	9	100%	3	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	59		45		11		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	39	66%	31	69%	7	64%	1	33%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	6	10%	5	11%	1	9%	-	-
MSM & IDU	1	2%	1	2%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	13	22%	8	18%	3	27%	2	67%
Total	59		45		11		3	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	16	80%	22	88%	1	100%	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	15%	3	12%	-	-	-	-	-	-	-	-
Subtotal	20	100%	25	100%	1	100%	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	100%	5	45%	-	-	-	-	-	-	-	-
Other/unknown	-	-	6	55%	1	100%	-	-	-	-	-	-
Subtotal	1	100%	11	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	21		36		2		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	76%	22	61%	1	50%	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	5%	5	14%	-	-	-	-	-	-	-	-
MSM & IDU	1	5%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	14%	9	25%	1	50%	-	-	-	-	-	-
Total	21		36		2		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Montgomery

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	24.6	63	84%	23.0	59	77%	19.1	49	78%	19.9	51	78%
Females	4.3	12	16%	6.5	18	23%	5.1	14	22%	5.1	14	22%
Age at diagnosis (yr)												
<13	*	2	3%	*	-	-	*	-	-	*	1	2%
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	29.5	10	13%	*	4	5%	*	2	3%	14.8	5	8%
20-24	51.3	19	25%	59.4	22	29%	40.5	15	24%	35.1	13	20%
25-29	36.5	13	17%	25.2	9	12%	44.9	16	25%	33.7	12	18%
30-34	14.7	5	7%	29.5	10	13%	*	3	5%	17.7	6	9%
35-39	*	4	5%	33.5	10	13%	33.5	10	16%	33.5	10	15%
40-44	22.1	7	9%	*	4	5%	*	2	3%	15.8	5	8%
45-49	17.9	6	8%	23.9	8	10%	20.9	7	11%	*	4	6%
50-54	13.1	5	7%	18.3	7	9%	15.7	6	10%	13.1	5	8%
55-64	*	1	1%	*	2	3%	*	2	3%	*	3	5%
65+	*	3	4%	*	1	1%	*	-	-	*	1	2%
Race/Ethnicity												
White, not Hispanic	5.7	22	29%	6.5	25	32%	7.0	27	43%	3.9	15	23%
Black/African American, not Hispanic	45.9	51	68%	43.2	48	62%	29.7	33	52%	39.6	44	68%
Hispanic/Latino	*	-	-	*	3	4%	*	2	3%	*	4	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	3%	*	1	1%	*	1	2%	*	2	3%
Race/Sex												
White, not Hispanic Males	11.8	22	29%	10.7	20	26%	11.8	22	35%	7.0	13	20%
White, not Hispanic Females	*	-	-	2.5	5	6%	2.5	5	8%	*	2	3%
Black/African American, not Hispanic Males	77.4	40	53%	67.7	35	45%	46.5	24	38%	67.7	35	54%
Black/African American, not Hispanic Females	18.5	11	15%	21.8	13	17%	15.1	9	14%	15.1	9	14%
Hispanic/Latino Males	*	-	-	*	3	4%	*	2	3%	*	2	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	2	3%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	3%	*	1	1%	*	1	2%	*	2	3%
Total	14.1	75		14.4	77		11.8	63		12.2	65	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Montgomery

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	52	85%	41	69%	40	82%	36	72%
Injection drug use (IDU)	1	2%	1	2%	-	-	-	-
Male-to-male sexual contact and IDU	1	2%	-	-	1	2%	-	-
Heterosexual contact	2	3%	6	10%	3	6%	3	6%
Other/unknown	5	8%	11	19%	5	10%	11	22%
Subtotal	61	100%	59	100%	49	100%	50	100%
Female adult or adolescent								
Injection drug use	-	-	3	17%	-	-	1	7%
Heterosexual contact	10	83%	11	61%	9	64%	5	36%
Other/unknown	2	17%	4	22%	5	36%	8	57%
Subtotal	12	100%	18	100%	14	100%	14	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	50%	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	1	100%
Subtotal	2	100%	-	-	-	-	1	100%
Total	75		77		63		65	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Montgomery

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	52	69%	38	49%	38	60%	33	51%
Injection drug use (IDU) only	1	1%	1	1%	-	-	-	-
Heterosexual contact only	12	16%	17	22%	12	19%	8	12%
MSM & IDU	-	-	-	-	1	2%	-	-
IDU & Heterosexual contact	-	-	3	4%	-	-	1	2%
MSM & Heterosexual contact	-	-	3	4%	2	3%	3	5%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-
Perinatal exposure	1	1%	-	-	-	-	-	-
Other/unknown	8	11%	15	19%	10	16%	20	31%
Total	75		77		63		65	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Montgomery

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	42	84%	38	76%	28	72%	37	74%	36	80%
Females	8	16%	12	24%	11	28%	13	26%	9	20%
Age at HIV diagnosis (yr)										
<13	2	4%	-	-	-	-	1	2%	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	8	16%	3	6%	2	5%	4	8%	3	7%
20-24	17	34%	15	30%	11	28%	11	22%	16	36%
25-29	9	18%	5	10%	11	28%	11	22%	8	18%
30-34	2	4%	9	18%	2	5%	2	4%	1	2%
35-39	2	4%	5	10%	7	18%	8	16%	7	16%
40-44	4	8%	4	8%	-	-	5	10%	5	11%
45-49	1	2%	3	6%	1	3%	3	6%	1	2%
50-54	3	6%	3	6%	3	8%	3	6%	2	4%
55-64	-	-	2	4%	2	5%	1	2%	-	-
65+	2	4%	1	2%	-	-	1	2%	2	4%
Race/Ethnicity										
White, not Hispanic	12	24%	18	36%	14	36%	13	26%	16	36%
Black/African American, not Hispanic	37	74%	29	58%	23	59%	32	64%	27	60%
Hispanic/Latino	-	-	2	4%	1	3%	4	8%	2	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	3%	1	2%	-	-
Race/Sex										
White, not Hispanic Males	12	24%	14	28%	11	28%	11	22%	15	33%
White, not Hispanic Females	-	-	4	8%	3	8%	2	4%	1	2%
Black/African American, not Hispanic Males	30	60%	21	42%	15	38%	23	46%	20	44%
Black/African American, not Hispanic Females	7	14%	8	16%	8	21%	9	18%	7	16%
Hispanic/Latino Males	-	-	2	4%	1	3%	2	4%	1	2%
Hispanic/Latina Females	-	-	-	-	-	-	2	4%	1	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	3%	1	2%	-	-
Total	50		50		39		50		45	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	34	85%	26	68%	23	82%	27	75%	31	86%
Injection drug use (IDU)	-	-	1	3%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	-	-	-	-	-	-	1	3%
Heterosexual contact	1	3%	3	8%	2	7%	2	6%	-	-
Other/unknown	4	10%	8	21%	3	11%	7	19%	4	11%
Subtotal	40	100%	38	100%	28	100%	36	100%	36	100%
Female adult or adolescent										
Injection drug use	-	-	3	25%	-	-	1	8%	-	-
Heterosexual contact	6	75%	6	50%	6	55%	4	31%	5	56%
Other/unknown	2	25%	3	25%	5	45%	8	62%	4	44%
Subtotal	8	100%	12	100%	11	100%	13	100%	9	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	50%	-	-	-	-	-	-	-	-
Other/unknown	1	50%	-	-	-	-	1	100%	-	-
Subtotal	2	100%	-	-	-	-	1	100%	-	-
Total	50		50		39		50		45	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	34	68%	23	46%	22	56%	24	48%	31	69%
Injection drug use (IDU) only	-	-	1	2%	-	-	-	-	-	-
Heterosexual contact only	7	14%	9	18%	8	21%	6	12%	5	11%
MSM & IDU	-	-	-	-	-	-	-	-	1	2%
IDU & Heterosexual contact	-	-	3	6%	-	-	1	2%	-	-
MSM & Heterosexual contact	-	-	3	6%	1	3%	3	6%	-	-
MSM & IDU & Heterosexual contact	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	-	-	-	-	-	-	-	-
Other/unknown	7	14%	11	22%	8	21%	16	32%	8	18%
Total	50		50		39		50		45	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Montgomery

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	25	78%	26	84%	33	92%	24	83%	24	80%	1,089	84%
Females	7	22%	5	16%	3	8%	5	17%	6	20%	203	16%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	3%	-	-	17	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	<1%
15-19	3	9%	3	10%	2	6%	1	3%	3	10%	34	3%
20-24	3	9%	5	16%	4	11%	4	14%	5	17%	110	9%
25-29	3	9%	5	16%	7	19%	6	21%	4	13%	221	17%
30-34	8	25%	6	19%	3	8%	3	10%	3	10%	252	20%
35-39	2	6%	4	13%	7	19%	5	17%	6	20%	243	19%
40-44	4	13%	-	-	3	8%	3	10%	3	10%	169	13%
45-49	5	16%	5	16%	7	19%	1	3%	2	7%	126	10%
50-54	2	6%	3	10%	3	8%	4	14%	1	3%	60	5%
55-64	1	3%	-	-	-	-	1	3%	2	7%	40	3%
65+	1	3%	-	-	-	-	-	-	1	3%	18	1%
Race/Ethnicity												
White, not Hispanic	13	41%	14	45%	16	44%	7	24%	10	33%	683	53%
Black/African American, not Hispanic	18	56%	15	48%	18	50%	21	72%	18	60%	580	45%
Hispanic/Latino	-	-	-	-	2	6%	-	-	-	-	12	1%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	1	3%	2	6%	-	-	1	3%	2	7%	13	1%
Race/Sex												
White, not Hispanic Males	13	41%	12	39%	15	42%	6	21%	10	33%	622	48%
White, not Hispanic Females	-	-	2	6%	1	3%	1	3%	-	-	61	5%
Black/African American, not Hispanic Males	11	34%	13	42%	16	44%	17	59%	14	47%	443	34%
Black/African American, not Hispanic Females	7	22%	2	6%	2	6%	4	14%	4	13%	137	11%
Hispanic/Latino Males	-	-	-	-	2	6%	-	-	-	-	11	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	3%	2	6%	-	-	1	3%	2	7%	13	1%
Total	32		31		36		29		30		1,292	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	21	84%	18	69%	26	79%	19	79%	17	71%	789	73%
Injection drug use (IDU)	1	4%	-	-	-	-	-	-	1	4%	47	4%
Male-to-male sexual contact and IDU	1	4%	1	4%	1	3%	-	-	-	-	74	7%
Heterosexual contact	1	4%	3	12%	3	9%	1	4%	1	4%	62	6%
Other/unknown	1	4%	4	15%	3	9%	4	17%	5	21%	107	10%
Subtotal	25	100%	26	100%	33	100%	24	100%	24	100%	1,079	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	1	25%	-	-	37	19%
Heterosexual contact	7	100%	5	100%	3	100%	2	50%	3	50%	142	72%
Other/unknown	-	-	-	-	-	-	1	25%	3	50%	17	9%
Subtotal	7	100%	5	100%	3	100%	4	100%	6	100%	196	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	1	100%	-	-	12	71%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	29%
Subtotal	-	-	-	-	-	-	1	100%	-	-	17	100%
Total	32		31		36		29		30		1,292	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	21	66%	18	58%	25	69%	18	62%	17	57%	756	59%
Injection drug use (IDU) only	1	3%	-	-	-	-	1	3%	1	3%	52	4%
Heterosexual contact only	8	25%	8	26%	6	17%	3	10%	4	13%	204	16%
MSM & IDU	1	3%	1	3%	1	3%	-	-	-	-	63	5%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	31	2%
MSM & Heterosexual contact	-	-	-	-	1	3%	1	3%	-	-	33	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	11	1%
Perinatal exposure	-	-	-	-	-	-	1	3%	-	-	12	1%
Other/unknown	1	3%	4	13%	3	8%	5	17%	8	27%	130	10%
Total	32		31		36		29		30		1,292	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Montgomery

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Montgomery

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	402.9	1,033	79%	471	76%	562	82%
Females	98.3	272	21%	151	24%	121	18%
Age at end of year							
<13	*	2	<1%	2	<1%	-	-
13-14	*	1	<1%	1	<1%	-	-
15-19	32.5	11	1%	8	1%	3	<1%
20-24	170.2	63	5%	48	8%	15	2%
25-29	314.1	112	9%	76	12%	36	5%
30-34	300.6	102	8%	56	9%	46	7%
35-39	425.9	127	10%	66	11%	61	9%
40-44	433.3	137	10%	69	11%	68	10%
45-49	575.9	193	15%	78	13%	115	17%
50-54	621.8	238	18%	92	15%	146	21%
55-64	356.9	255	20%	102	16%	153	22%
65+	72.2	64	5%	24	4%	40	6%
Race/Ethnicity							
White, not Hispanic	143.4	550	42%	251	40%	299	44%
Black/African American, not Hispanic	610.6	679	52%	337	54%	342	50%
Hispanic/Latino	267.9	37	3%	16	3%	21	3%
Asian/Pacific Islander	80.9	9	1%	4	1%	5	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	30	2%	14	2%	16	2%
Race/Sex							
White, not Hispanic Males	261.5	487	37%	217	35%	270	40%
White, not Hispanic Females	31.9	63	5%	34	5%	29	4%
Black/African American, not Hispanic Males	936.8	484	37%	227	36%	257	38%
Black/African American, not Hispanic Females	327.6	195	15%	110	18%	85	12%
Hispanic/Latino Males	456.0	32	2%	13	2%	19	3%
Hispanic/Latina Females	73.6	5	<1%	3	<1%	2	<1%
Asian/Pacific Islander Males	174.5	9	1%	4	1%	5	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	30	2%	14	2%	16	2%
Total	244.8	1,305		622		683	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	760	74%	336	72%	424	76%
Injection drug use (IDU)	26	3%	13	3%	13	2%
Male-to-male sexual contact and IDU	44	4%	13	3%	31	6%
Heterosexual contact	54	5%	17	4%	37	7%
Other/unknown	143	14%	89	19%	54	10%
Subtotal	1,027	100%	468	100%	559	100%
Female adult or adolescent						
Injection drug use	25	10%	14	10%	11	9%
Heterosexual contact	186	71%	95	65%	91	78%
Other/unknown	52	20%	38	26%	14	12%
Subtotal	263	100%	147	100%	116	100%
Child (<13 yrs at diagnosis)						
Perinatal	12	80%	6	86%	6	75%
Other/unknown	3	20%	1	14%	2	25%
Subtotal	15	100%	7	100%	8	100%
Total	1,305		622		683	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	726	56%	324	52%	402	59%
Injection drug use (IDU) only	25	2%	13	2%	12	2%
Heterosexual contact only	240	18%	112	18%	128	19%
MSM & IDU	39	3%	11	2%	28	4%
IDU & Heterosexual contact	26	2%	14	2%	12	2%
MSM & Heterosexual contact	34	3%	12	2%	22	3%
MSM & IDU & Heterosexual contact	5	<1%	2	<1%	3	<1%
Perinatal exposure	13	1%	7	1%	6	1%
Other/unknown	197	15%	127	20%	70	10%
Total	1,305		622		683	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Montgomery

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	396	82%	321	67%	26	81%	4	44%	-	-	13	62%
Injection drug use (IDU)	6	1%	19	4%	1	3%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	22	5%	19	4%	2	6%	-	-	-	-	1	5%
Heterosexual contact	9	2%	43	9%	1	3%	1	11%	-	-	-	-
Other/unknown	51	11%	79	16%	2	6%	4	44%	-	-	7	33%
Subtotal	484	100%	481	100%	32	100%	9	100%	-	-	21	100%
Female adult or adolescent												
Injection drug use	12	20%	13	7%	-	-	-	-	-	-	-	-
Heterosexual contact	39	64%	137	73%	3	60%	-	-	-	-	7	78%
Other/unknown	10	16%	38	20%	2	40%	-	-	-	-	2	22%
Subtotal	61	100%	188	100%	5	100%	-	-	-	-	9	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	60%	9	90%	-	-	-	-	-	-	-	-
Other/unknown	2	40%	1	10%	-	-	-	-	-	-	-	-
Subtotal	5	100%	10	100%	-	-	-	-	-	-	-	-
Total	550		679		37		9		0		30	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	388	71%	299	44%	24	65%	3	33%	-	-	12	40%
Injection drug use (IDU) only	7	1%	17	3%	1	3%	-	-	-	-	-	-
Heterosexual contact only	48	9%	180	27%	4	11%	1	11%	-	-	7	23%
MSM & IDU	20	4%	17	3%	1	3%	-	-	-	-	1	3%
IDU & Heterosexual contact	11	2%	15	2%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	8	1%	22	3%	2	5%	1	11%	-	-	1	3%
MSM & IDU & Heterosexual contact	2	0%	2	0%	1	3%	-	-	-	-	-	-
Perinatal exposure	3	1%	10	1%	-	-	-	-	-	-	-	-
Other/unknown	63	11%	117	17%	4	11%	4	44%	-	-	9	30%
Total	550		679		37		9		0		30	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Montgomery

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	827	79%	884	79%	932	79%	992	79%	1,033	79%
Females	221	21%	235	21%	246	21%	259	21%	272	21%
Age at end of year										
<13	5	<1%	4	<1%	3	<1%	3	<1%	2	<1%
13-14	1	<1%	1	<1%	2	<1%	1	<1%	1	<1%
15-19	17	2%	15	1%	11	1%	13	1%	11	1%
20-24	39	4%	60	5%	65	6%	70	6%	63	5%
25-29	71	7%	78	7%	88	7%	102	8%	112	9%
30-34	93	9%	90	8%	94	8%	96	8%	102	8%
35-39	95	9%	90	8%	104	9%	122	10%	127	10%
40-44	164	16%	159	14%	143	12%	133	11%	137	10%
45-49	229	22%	234	21%	221	19%	203	16%	193	15%
50-54	155	15%	180	16%	197	17%	231	18%	238	18%
55-64	155	15%	176	16%	208	18%	228	18%	255	20%
65+	24	2%	32	3%	42	4%	49	4%	64	5%
Race/Ethnicity										
White, not Hispanic	462	44%	486	43%	508	43%	529	42%	550	42%
Black/African American, not Hispanic	527	50%	571	51%	603	51%	648	52%	679	52%
Hispanic/Latino	26	2%	28	3%	31	3%	35	3%	37	3%
Asian/Pacific Islander	8	1%	8	1%	8	1%	9	1%	9	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	25	2%	26	2%	28	2%	30	2%	30	2%
Race/Sex										
White, not Hispanic Males	411	39%	431	39%	449	38%	468	37%	487	37%
White, not Hispanic Females	51	5%	55	5%	59	5%	61	5%	63	5%
Black/African American, not Hispanic Males	367	35%	401	36%	426	36%	463	37%	484	37%
Black/African American, not Hispanic Females	160	15%	170	15%	177	15%	185	15%	195	15%
Hispanic/Latino Males	24	2%	26	2%	29	2%	31	2%	32	2%
Hispanic/Latina Females	2	<1%	2	<1%	2	<1%	4	<1%	5	<1%
Asian/Pacific Islander Males	8	1%	8	1%	8	1%	9	1%	9	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	25	2%	26	2%	28	2%	30	2%	30	2%
Total	1,048		1,119		1,178		1,251		1,305	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Montgomery

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	598	73%	643	73%	678	73%	724	73%	760	74%
Injection drug use (IDU)	26	3%	26	3%	26	3%	26	3%	26	3%
Male-to-male sexual contact and IDU	43	5%	43	5%	44	5%	44	4%	44	4%
Heterosexual contact	42	5%	47	5%	51	6%	54	5%	54	5%
Other/unknown	112	14%	119	14%	127	14%	138	14%	143	14%
Subtotal	821	100%	878	100%	926	100%	986	100%	1,027	100%
Female adult or adolescent										
Injection drug use	22	10%	24	11%	24	10%	25	10%	25	10%
Heterosexual contact	158	75%	168	74%	176	74%	181	72%	186	71%
Other/unknown	32	15%	34	15%	37	16%	44	18%	52	20%
Subtotal	212	100%	226	100%	237	100%	250	100%	263	100%
Child (<13 yrs at diagnosis)										
Perinatal	12	80%	12	80%	12	80%	12	80%	12	80%
Other/unknown	3	20%	3	20%	3	20%	3	20%	3	20%
Subtotal	15	100%	15	100%	15	100%	15	100%	15	100%
Total	1,048		1,119		1,178		1,251		1,305	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Montgomery

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	571	54%	613	55%	647	55%	690	55%	726	56%
Injection drug use (IDU) only	24	2%	25	2%	25	2%	25	2%	25	2%
Heterosexual contact only	200	19%	215	19%	227	19%	235	19%	240	18%
MSM & IDU	38	4%	38	3%	39	3%	39	3%	39	3%
IDU & Heterosexual contact	24	2%	25	2%	25	2%	26	2%	26	2%
MSM & Heterosexual contact	27	3%	30	3%	31	3%	34	3%	34	3%
MSM & IDU & Heterosexual contact	5	<1%	5	<1%	5	<1%	5	<1%	5	<1%
Perinatal exposure	13	1%	13	1%	13	1%	13	1%	13	1%
Other/unknown	146	14%	155	14%	166	14%	184	15%	197	15%
Total	1,048		1,119		1,178		1,251		1,305	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.