

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Ottawa

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	63.8	13	87%	<10	*	<10	*
Females	*	2	13%	<10	*	<10	*
Age at end of year							
<13	*	1	7%	<10	*	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	7%	<10	*	-	-
25-29	*	1	7%	<10	*	-	-
30-34	*	-	-	-	-	-	-
35-39	*	-	-	-	-	-	-
40-44	*	2	13%	<10	*	-	-
45-49	*	2	13%	<10	*	<10	*
50-54	*	3	20%	<10	*	<10	*
55-64	70.8	5	33%	<10	*	<10	*
65+	*	-	-	-	-	-	-
Race/Ethnicity							
White, not Hispanic	23.6	9	60%	<10	*	<10	*
Black/African American, not Hispanic	*	3	20%	<10	*	<10	*
Hispanic/Latino	*	1	7%	-	-	<10	*
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	2	13%	<10	*	-	-
Race/Sex							
White, not Hispanic Males	42.5	8	53%	<10	*	<10	*
White, not Hispanic Females	*	1	7%	<10	*	-	-
Black/African American, not Hispanic Males	*	3	20%	<10	*	<10	*
Black/African American, not Hispanic Females	*	-	-	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-
Hispanic/Latina Females	*	1	7%	-	-	<10	*
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	13%	<10	*	-	-
Total	36.4	15		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Ottawa

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	9	75%	<10	*	<10	*
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	8%	-	-	<10	*
Heterosexual contact	2	17%	<10	*	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	12	100%	<10	*	<10	*
Female adult or adolescent						
Injection drug use	1	50%	<10	*	-	-
Heterosexual contact	1	50%	-	-	<10	*
Other/unknown	-	-	-	-	-	-
Subtotal	2	100%	<10	*	<10	*
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	<10	*	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	<10	*	-	-
Total	15		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Ottawa

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	8	53%	<10	*	<10	*
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	3	20%	<10	*	<10	*
MSM & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	1	7%	<10	*	-	-
MSM & Heterosexual contact	1	7%	<10	*	-	-
MSM & IDU & Heterosexual contact	1	7%	-	-	<10	*
Perinatal exposure	1	7%	<10	*	-	-
Other/unknown	-	-	-	-	-	-
Total	15		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Ottawa

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	9	82%	11	85%	12	86%	13	87%	13	87%
Females	2	18%	2	15%	2	14%	2	13%	2	13%
Age at end of year										
<13	-	-	-	-	-	-	1	7%	1	7%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	-	-	1	8%	1	7%	1	7%	1	7%
25-29	-	-	-	-	1	7%	1	7%	1	7%
30-34	-	-	-	-	-	-	-	-	-	-
35-39	1	9%	1	8%	1	7%	1	7%	-	-
40-44	1	9%	1	8%	1	7%	1	7%	2	13%
45-49	3	27%	3	23%	2	14%	2	13%	2	13%
50-54	5	45%	3	23%	4	29%	3	20%	3	20%
55-64	1	9%	4	31%	4	29%	5	33%	5	33%
65+	-	-	-	-	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	7	64%	8	62%	9	64%	9	60%	9	60%
Black/African American, not Hispanic	2	18%	2	15%	2	14%	3	20%	3	20%
Hispanic/Latino	1	9%	1	8%	1	7%	1	7%	1	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	9%	2	15%	2	14%	2	13%	2	13%
Race/Sex										
White, not Hispanic Males	6	55%	7	54%	8	57%	8	53%	8	53%
White, not Hispanic Females	1	9%	1	8%	1	7%	1	7%	1	7%
Black/African American, not Hispanic Males	2	18%	2	15%	2	14%	3	20%	3	20%
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-
Hispanic/Latina Females	1	9%	1	8%	1	7%	1	7%	1	7%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	9%	2	15%	2	14%	2	13%	2	13%
Total	11		13		14		15		15	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Ottawa

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	6	67%	8	73%	9	75%	9	75%	9	75%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	11%	1	9%	1	8%	1	8%	1	8%
Heterosexual contact	2	22%	2	18%	2	17%	2	17%	2	17%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	9	100%	11	100%	12	100%	12	100%	12	100%
Female adult or adolescent										
Injection drug use	1	50%	1	50%	1	50%	1	50%	1	50%
Heterosexual contact	1	50%	1	50%	1	50%	1	50%	1	50%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	2	100%	2	100%	2	100%	2	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	1	100%
Total	11		13		14		15		15	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Ottawa

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	5	45%	7	54%	8	57%	8	53%	8	53%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	3	27%	3	23%	3	21%	3	20%	3	20%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	9%	1	8%	1	7%	1	7%	1	7%
MSM & Heterosexual contact	1	9%	1	8%	1	7%	1	7%	1	7%
MSM & IDU & Heterosexual contact	1	9%	1	8%	1	7%	1	7%	1	7%
Perinatal exposure	-	-	-	-	-	-	1	7%	1	7%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Total	11		13		14		15		15	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.