

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Perry

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Perry

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	128.5	23	79%	11	85%	12	75%
Females	33.5	6	21%	2	15%	4	25%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	3%	1	8%	-	-
25-29	*	1	3%	1	8%	-	-
30-34	*	1	3%	-	-	1	6%
35-39	*	3	10%	3	23%	-	-
40-44	298.4	7	24%	4	31%	3	19%
45-49	276.6	7	24%	1	8%	6	38%
50-54	*	4	14%	-	-	4	25%
55-64	*	4	14%	3	23%	1	6%
65+	*	1	3%	-	-	1	6%
Race/Ethnicity							
White, not Hispanic	72.0	25	86%	12	92%	13	81%
Black/African American, not Hispanic	*	-	-	-	-	-	-
Hispanic/Latino	*	1	3%	-	-	1	6%
Asian/Pacific Islander	*	2	7%	1	8%	1	6%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	1	3%	-	-	1	6%
Race/Sex							
White, not Hispanic Males	109.4	19	66%	10	77%	9	56%
White, not Hispanic Females	34.6	6	21%	2	15%	4	25%
Black/African American, not Hispanic Males	*	-	-	-	-	-	-
Black/African American, not Hispanic Females	*	-	-	-	-	-	-
Hispanic/Latino Males	*	1	3%	-	-	1	6%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	7%	1	8%	1	6%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	1	3%	-	-	1	6%
Total	81.0	29		13		16	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Perry

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	12	52%	6	55%	6	50%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	4%	1	9%	-	-
Heterosexual contact	4	17%	2	18%	2	17%
Other/unknown	6	26%	2	18%	4	33%
Subtotal	23	100%	11	100%	12	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	6	100%	2	100%	4	100%
Other/unknown	-	-	-	-	-	-
Subtotal	6	100%	2	100%	4	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	29		13		16	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Perry

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	12	41%	6	46%	6	38%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	10	34%	4	31%	6	38%
MSM & IDU	1	3%	1	8%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	6	21%	2	15%	4	25%
Total	29		13		16	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Perry

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	10	53%	-	-	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	<10	*	-	-	-	-
Heterosexual contact	4	21%	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	26%	-	-	-	-	<10	*	-	-	-	-
Subtotal	19	100%	-	-	<10	*	<10	*	-	-	<10	*
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	6	100%	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	6	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	25		0		<10		<10		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Perry

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	10	40%	-	-	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	10	40%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	<10	*	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	20%	-	-	-	-	<10	*	-	-	-	-
Total	25		0		<10		<10		0		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Perry

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	17	74%	18	75%	20	77%	21	78%	23	79%
Females	6	26%	6	25%	6	23%	6	22%	6	21%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	-	-	1	4%	1	4%	1	4%	1	3%
25-29	-	-	-	-	-	-	-	-	1	3%
30-34	1	4%	-	-	1	4%	1	4%	1	3%
35-39	6	26%	6	25%	4	15%	4	15%	3	10%
40-44	7	30%	6	25%	6	23%	6	22%	7	24%
45-49	4	17%	5	21%	8	31%	8	30%	7	24%
50-54	-	-	1	4%	1	4%	2	7%	4	14%
55-64	5	22%	5	21%	5	19%	5	19%	4	14%
65+	-	-	-	-	-	-	-	-	1	3%
Race/Ethnicity										
White, not Hispanic	19	83%	20	83%	22	85%	23	85%	25	86%
Black/African American, not Hispanic	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino	1	4%	1	4%	1	4%	1	4%	1	3%
Asian/Pacific Islander	2	9%	2	8%	2	8%	2	7%	2	7%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	1	4%	1	4%	1	4%	1	3%
Race/Sex										
White, not Hispanic Males	13	57%	14	58%	16	62%	17	63%	19	66%
White, not Hispanic Females	6	26%	6	25%	6	23%	6	22%	6	21%
Black/African American, not Hispanic Males	-	-	-	-	-	-	-	-	-	-
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	-	-
Hispanic/Latino Males	1	4%	1	4%	1	4%	1	4%	1	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	2	9%	2	8%	2	8%	2	7%	2	7%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	1	4%	1	4%	1	4%	1	3%
Total	23		24		26		27		29	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Perry

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	8	47%	9	50%	10	50%	11	52%	12	52%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	6%	1	6%	1	5%	1	5%	1	4%
Heterosexual contact	4	24%	4	22%	4	20%	4	19%	4	17%
Other/unknown	4	24%	4	22%	5	25%	5	24%	6	26%
Subtotal	17	100%	18	100%	20	100%	21	100%	23	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	6	100%	6	100%	6	100%	6	100%	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	6	100%	6	100%	6	100%	6	100%	6	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	23		24		26		27		29	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Perry

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	8	35%	9	38%	10	38%	11	41%	12	41%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	10	43%	10	42%	10	38%	10	37%	10	34%
MSM & IDU	1	4%	1	4%	1	4%	1	4%	1	3%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	4	17%	4	17%	5	19%	5	19%	6	21%
Total	23		24		26		27		29	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.