

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Richland

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Richland

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	130.8	81	75%	39	78%	42	72%
Females	45.0	27	25%	11	22%	16	28%
Age at end of year							
<13	*	1	1%	1	2%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	4	4%	1	2%	3	5%
25-29	124.4	9	8%	6	12%	3	5%
30-34	96.4	7	6%	3	6%	4	7%
35-39	237.2	17	16%	8	16%	9	16%
40-44	215.7	16	15%	9	18%	7	12%
45-49	230.1	18	17%	6	12%	12	21%
50-54	182.0	16	15%	9	18%	7	12%
55-64	99.7	17	16%	6	12%	11	19%
65+	*	3	3%	1	2%	2	3%
Race/Ethnicity							
White, not Hispanic	49.5	52	48%	27	54%	25	43%
Black/African American, not Hispanic	405.6	46	43%	20	40%	26	45%
Hispanic/Latino	377.0	8	7%	3	6%	5	9%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	1%	-	-	1	2%
Unknown	*	1	1%	-	-	1	2%
Race/Sex							
White, not Hispanic Males	80.9	42	39%	22	44%	20	34%
White, not Hispanic Females	18.8	10	9%	5	10%	5	9%
Black/African American, not Hispanic Males	432.8	31	29%	14	28%	17	29%
Black/African American, not Hispanic Females	359.0	15	14%	6	12%	9	16%
Hispanic/Latino Males	605.0	7	6%	3	6%	4	7%
Hispanic/Latina Females	*	1	1%	-	-	1	2%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	1%	-	-	1	2%
Unknown	*	1	1%	-	-	1	2%
Total	88.6	108		50		58	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Richland

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	46	57%	19	49%	27	64%
Injection drug use (IDU)	3	4%	-	-	3	7%
Male-to-male sexual contact and IDU	4	5%	2	5%	2	5%
Heterosexual contact	9	11%	5	13%	4	10%
Other/unknown	19	23%	13	33%	6	14%
Subtotal	81	100%	39	100%	42	100%
Female adult or adolescent						
Injection drug use	2	8%	-	-	2	13%
Heterosexual contact	17	68%	7	70%	10	67%
Other/unknown	6	24%	3	30%	3	20%
Subtotal	25	100%	10	100%	15	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	2	100%	1	100%	1	100%
Total	108		50		58	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Richland

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	42	39%	17	34%	25	43%
Injection drug use (IDU) only	2	2%	-	-	2	3%
Heterosexual contact only	26	24%	12	24%	14	24%
MSM & IDU	3	3%	1	2%	2	3%
IDU & Heterosexual contact	3	3%	-	-	3	5%
MSM & Heterosexual contact	4	4%	2	4%	2	3%
MSM & IDU & Heterosexual contact	1	1%	1	2%	-	-
Perinatal exposure	2	2%	1	2%	1	2%
Other/unknown	25	23%	16	32%	9	16%
Total	108		50		58	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Richland

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	25	60%	17	55%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	3	7%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	5%	2	6%	-	-	-	-	-	-	-	-
Heterosexual contact	3	7%	5	16%	<10	*	-	-	-	-	-	-
Other/unknown	9	21%	7	23%	<10	*	-	-	-	-	-	-
Subtotal	42	100%	31	100%	<10	*	-	-	-	-	<10	*
Female adult or adolescent												
Injection drug use	1	11%	1	7%	-	-	-	-	-	-	-	-
Heterosexual contact	4	44%	12	86%	-	-	-	-	<10	*	-	-
Other/unknown	4	44%	1	7%	<10	*	-	-	-	-	-	-
Subtotal	9	100%	14	100%	<10	*	-	-	<10	*	-	-
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-	-	-	-	-
Total	52		46		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Richland

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	23	44%	15	33%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	1	2%	1	2%	-	-	-	-	-	-	-	-
Heterosexual contact only	7	13%	17	37%	<10	*	-	-	<10	*	-	-
MSM & IDU	2	4%	1	2%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	3	6%	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	2	4%	2	4%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	1	2%	1	2%	-	-	-	-	-	-	-	-
Other/unknown	13	25%	8	17%	<10	*	-	-	-	-	-	-
Total	52		46		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Richland

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	64	72%	68	73%	74	75%	81	75%	81	75%
Females	25	28%	25	27%	25	25%	27	25%	27	25%
Age at end of year										
<13	-	-	-	-	-	-	1	1%	1	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	1%	1	1%	1	1%	1	1%	-	-
20-24	4	4%	4	4%	5	5%	3	3%	4	4%
25-29	4	4%	6	6%	7	7%	9	8%	9	8%
30-34	15	17%	11	12%	9	9%	9	8%	7	6%
35-39	12	13%	15	16%	15	15%	17	16%	17	16%
40-44	18	20%	19	20%	20	20%	21	19%	16	15%
45-49	13	15%	11	12%	13	13%	12	11%	18	17%
50-54	12	13%	13	14%	12	12%	16	15%	16	15%
55-64	9	10%	12	13%	16	16%	16	15%	17	16%
65+	1	1%	1	1%	1	1%	3	3%	3	3%
Race/Ethnicity										
White, not Hispanic	43	48%	45	48%	48	48%	52	48%	52	48%
Black/African American, not Hispanic	39	44%	41	44%	42	42%	46	43%	46	43%
Hispanic/Latino	5	6%	5	5%	7	7%	8	7%	8	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	1	1%	1	1%	1	1%	1	1%	1	1%
Race/Sex										
White, not Hispanic Males	34	38%	36	39%	39	39%	42	39%	42	39%
White, not Hispanic Females	9	10%	9	10%	9	9%	10	9%	10	9%
Black/African American, not Hispanic Males	25	28%	27	29%	28	28%	31	29%	31	29%
Black/African American, not Hispanic Females	14	16%	14	15%	14	14%	15	14%	15	14%
Hispanic/Latino Males	4	4%	4	4%	6	6%	7	6%	7	6%
Hispanic/Latina Females	1	1%	1	1%	1	1%	1	1%	1	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	1	1%	1	1%	1	1%	1	1%	1	1%
Total	89		93		99		108		108	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Richland

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	36	56%	37	54%	42	57%	46	57%	46	57%
Injection drug use (IDU)	3	5%	3	4%	3	4%	3	4%	3	4%
Male-to-male sexual contact and IDU	4	6%	4	6%	4	5%	4	5%	4	5%
Heterosexual contact	7	11%	7	10%	7	9%	9	11%	9	11%
Other/unknown	14	22%	17	25%	18	24%	19	23%	19	23%
Subtotal	64	100%	68	100%	74	100%	81	100%	81	100%
Female adult or adolescent										
Injection drug use	2	8%	2	8%	2	8%	2	8%	2	8%
Heterosexual contact	16	67%	16	67%	16	67%	17	68%	17	68%
Other/unknown	6	25%	6	25%	6	25%	6	24%	6	24%
Subtotal	24	100%	24	100%	24	100%	25	100%	25	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	100%	1	100%	2	100%	2	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	1	100%	2	100%	2	100%
Total	89		93		99		108		108	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Richland

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	33	37%	34	37%	38	38%	42	39%	42	39%
Injection drug use (IDU) only	2	2%	2	2%	2	2%	2	2%	2	2%
Heterosexual contact only	23	26%	23	25%	23	23%	26	24%	26	24%
MSM & IDU	3	3%	3	3%	3	3%	3	3%	3	3%
IDU & Heterosexual contact	3	3%	3	3%	3	3%	3	3%	3	3%
MSM & Heterosexual contact	3	3%	3	3%	4	4%	4	4%	4	4%
MSM & IDU & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
Perinatal exposure	1	1%	1	1%	1	1%	2	2%	2	2%
Other/unknown	20	22%	23	25%	24	24%	25	23%	25	23%
Total	89		93		99		108		108	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.