

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Stark

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Stark

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	8.8	16	80%	12	80%	3	75%	1	100%
Females	*	4	20%	3	20%	1	25%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	20.7	5	25%	3	20%	1	25%	1	100%
25-29	32.0	7	35%	5	33%	2	50%	-	-
30-34	*	2	10%	2	13%	-	-	-	-
35-39	*	-	-	-	-	-	-	-	-
40-44	*	2	10%	2	13%	-	-	-	-
45-49	*	2	10%	1	7%	1	25%	-	-
50-54	*	-	-	-	-	-	-	-	-
55-64	*	2	10%	2	13%	-	-	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	2.8	9	45%	7	47%	1	25%	1	100%
Black/African American, not Hispanic	27.8	8	40%	5	33%	3	75%	-	-
Hispanic/Latino	*	1	5%	1	7%	-	-	-	-
Asian/Pacific Islander	*	1	5%	1	7%	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	1	5%	1	7%	-	-	-	-
Race/Sex									
White, not Hispanic Males	5.0	8	40%	6	40%	1	25%	1	100%
White, not Hispanic Females	*	1	5%	1	7%	-	-	-	-
Black/African American, not Hispanic Males	42.5	6	30%	4	27%	2	50%	-	-
Black/African American, not Hispanic Females	*	2	10%	1	7%	1	25%	-	-
Hispanic/Latino Males	*	1	5%	1	7%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	5%	1	7%	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	1	5%	1	7%	-	-	-	-
Total	5.3	20		15		4		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Stark

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	11	69%	7	58%	3	100%	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	6%	1	8%	-	-	-	-
Other/unknown	4	25%	4	33%	-	-	-	-
Subtotal	16	100%	12	100%	3	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	4	100%	3	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	4	100%	3	100%	1	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	20		15		4		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Stark

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	11	55%	7	47%	3	75%	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	5	25%	4	27%	1	25%	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	4	20%	4	27%	-	-	-	-
Total	20		15		4		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Stark

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	75%	5	83%	-	-	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	13%	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	13%	1	17%	1	100%	-	-	-	-	1	100%
Subtotal	8	100%	6	100%	1	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	100%	2	100%	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	2	100%	-	-	1	100%	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	9		8		1		1		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Stark

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	6	67%	5	63%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	2	22%	2	25%	-	-	1	100%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	11%	1	13%	1	100%	-	-	-	-	1	100%
Total	9		8		1		1		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Stark

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	13.7	25	89%	13.7	25	86%	6.6	12	92%	6.0	11	92%
Females	*	3	11%	*	4	14%	*	1	8%	*	1	8%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	*	4	14%	*	2	7%	*	1	8%	*	-	-
20-24	*	3	11%	33.1	8	28%	*	-	-	*	4	33%
25-29	*	2	7%	22.9	5	17%	*	3	23%	*	3	25%
30-34	*	3	11%	*	2	7%	*	2	15%	*	2	17%
35-39	24.2	5	18%	*	2	7%	*	-	-	*	-	-
40-44	*	4	14%	*	1	3%	*	4	31%	*	-	-
45-49	*	1	4%	*	2	7%	*	3	23%	*	2	17%
50-54	*	4	14%	*	3	10%	*	-	-	*	-	-
55-64	*	2	7%	*	3	10%	*	-	-	*	1	8%
65+	*	-	-	*	1	3%	*	-	-	*	-	-
Race/Ethnicity												
White, not Hispanic	5.5	18	64%	5.5	18	62%	3.1	10	77%	2.8	9	75%
Black/African American, not Hispanic	27.8	8	29%	31.2	9	31%	*	3	23%	*	3	25%
Hispanic/Latino	*	1	4%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	2	7%	*	-	-	*	-	-
Race/Sex												
White, not Hispanic Males	10.1	16	57%	10.1	16	55%	5.7	9	69%	5.0	8	67%
White, not Hispanic Females	*	2	7%	*	2	7%	*	1	8%	*	1	8%
Black/African American, not Hispanic Males	49.5	7	25%	49.5	7	24%	*	3	23%	*	3	25%
Black/African American, not Hispanic Females	*	1	4%	*	2	7%	*	-	-	*	-	-
Hispanic/Latino Males	*	1	4%	*	-	-	*	-	-	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	2	7%	*	-	-	*	-	-
Total	7.5	28		7.7	29		3.5	13		3.2	12	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Stark

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	19	76%	20	80%	10	83%	9	82%
Injection drug use (IDU)	-	-	1	4%	-	-	1	9%
Male-to-male sexual contact and IDU	2	8%	-	-	1	8%	-	-
Heterosexual contact	3	12%	2	8%	-	-	-	-
Other/unknown	1	4%	2	8%	1	8%	1	9%
Subtotal	25	100%	25	100%	12	100%	11	100%
Female adult or adolescent								
Injection drug use	-	-	1	25%	-	-	-	-
Heterosexual contact	3	100%	3	75%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	4	100%	1	100%	1	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	28		29		13		12	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Stark

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	18	64%	18	62%	10	77%	9	75%
Injection drug use (IDU) only	-	-	1	3%	-	-	1	8%
Heterosexual contact only	6	21%	5	17%	1	8%	1	8%
MSM & IDU	1	4%	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	3%	-	-	-	-
MSM & Heterosexual contact	1	4%	2	7%	-	-	-	-
MSM & IDU & Heterosexual contact	1	4%	-	-	1	8%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	4%	2	7%	1	8%	1	8%
Total	28		29		13		12	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Stark

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	17	89%	18	90%	11	92%	8	89%	12	80%
Females	2	11%	2	10%	1	8%	1	11%	3	20%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	16%	2	10%	1	8%	-	-	-	-
20-24	2	11%	8	40%	-	-	4	44%	3	20%
25-29	1	5%	3	15%	3	25%	3	33%	5	33%
30-34	3	16%	2	10%	2	17%	1	11%	2	13%
35-39	3	16%	1	5%	-	-	-	-	-	-
40-44	1	5%	1	5%	4	33%	-	-	2	13%
45-49	1	5%	-	-	2	17%	1	11%	1	7%
50-54	4	21%	2	10%	-	-	-	-	-	-
55-64	1	5%	1	5%	-	-	-	-	2	13%
65+	-	-	-	-	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	14	74%	16	80%	9	75%	6	67%	7	47%
Black/African American, not Hispanic	3	16%	3	15%	3	25%	3	33%	5	33%
Hispanic/Latino	1	5%	-	-	-	-	-	-	1	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	7%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	1	5%	-	-	-	-	1	7%
Race/Sex										
White, not Hispanic Males	12	63%	14	70%	8	67%	5	56%	6	40%
White, not Hispanic Females	2	11%	2	10%	1	8%	1	11%	1	7%
Black/African American, not Hispanic Males	3	16%	3	15%	3	25%	3	33%	4	27%
Black/African American, not Hispanic Females	-	-	-	-	-	-	-	-	1	7%
Hispanic/Latino Males	1	5%	-	-	-	-	-	-	1	7%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	7%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	5%	1	5%	-	-	-	-	1	7%
Total	19		20		12		9		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Stark

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	12	71%	15	83%	10	91%	7	88%	7	58%
Injection drug use (IDU)	-	-	-	-	-	-	1	13%	-	-
Male-to-male sexual contact and IDU	2	12%	-	-	-	-	-	-	-	-
Heterosexual contact	2	12%	1	6%	-	-	-	-	1	8%
Other/unknown	1	6%	2	11%	1	9%	-	-	4	33%
Subtotal	17	100%	18	100%	11	100%	8	100%	12	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	2	100%	1	100%	1	100%	3	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	2	100%	1	100%	1	100%	3	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	19		20		12		9		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Stark

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	11	58%	14	70%	10	83%	7	78%	7	47%
Injection drug use (IDU) only	-	-	-	-	-	-	1	11%	-	-
Heterosexual contact only	4	21%	3	15%	1	8%	1	11%	4	27%
MSM & IDU	1	5%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	5%	1	5%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	5%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	5%	2	10%	1	8%	-	-	4	27%
Total	19		20		12		9		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Stark

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	7	64%	11	79%	3	100%	5	100%	7	88%	310	84%
Females	4	36%	3	21%	-	-	-	-	1	13%	57	16%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	2%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	1	13%	6	2%
20-24	1	9%	1	7%	-	-	1	20%	3	38%	40	11%
25-29	2	18%	3	21%	-	-	-	-	2	25%	55	15%
30-34	2	18%	1	7%	-	-	2	40%	-	-	71	19%
35-39	1	9%	1	7%	2	67%	-	-	-	-	64	17%
40-44	4	36%	1	7%	-	-	-	-	1	13%	49	13%
45-49	-	-	3	21%	1	33%	1	20%	1	13%	34	9%
50-54	-	-	1	7%	-	-	-	-	-	-	20	5%
55-64	1	9%	2	14%	-	-	1	20%	-	-	17	5%
65+	-	-	1	7%	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	4	36%	4	29%	2	67%	3	60%	4	50%	256	70%
Black/African American, not Hispanic	6	55%	9	64%	1	33%	2	40%	4	50%	101	28%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	6	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	9%	1	7%	-	-	-	-	-	-	4	1%
Race/Sex												
White, not Hispanic Males	4	36%	4	29%	2	67%	3	60%	4	50%	227	62%
White, not Hispanic Females	-	-	-	-	-	-	-	-	-	-	29	8%
Black/African American, not Hispanic Males	2	18%	6	43%	1	33%	2	40%	3	38%	73	20%
Black/African American, not Hispanic Females	4	36%	3	21%	-	-	-	-	1	13%	28	8%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	6	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	9%	1	7%	-	-	-	-	-	-	4	1%
Total	11		14		3		5		8		367	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Stark

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	6	86%	7	64%	2	67%	4	80%	7	100%	224	73%
Injection drug use (IDU)	-	-	1	9%	-	-	-	-	-	-	13	4%
Male-to-male sexual contact and IDU	-	-	-	-	1	33%	-	-	-	-	13	4%
Heterosexual contact	-	-	2	18%	-	-	-	-	-	-	16	5%
Other/unknown	1	14%	1	9%	-	-	1	20%	-	-	42	14%
Subtotal	7	100%	11	100%	3	100%	5	100%	7	100%	308	100%
Female adult or adolescent												
Injection drug use	-	-	1	33%	-	-	-	-	-	-	13	25%
Heterosexual contact	4	100%	2	67%	-	-	-	-	1	100%	35	66%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	9%
Subtotal	4	100%	3	100%	-	-	-	-	1	100%	53	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	11		14		3		5		8		367	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Stark

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	6	55%	6	43%	2	67%	4	80%	7	88%	217	59%
Injection drug use (IDU) only	-	-	1	7%	-	-	-	-	-	-	15	4%
Heterosexual contact only	4	36%	4	29%	-	-	-	-	1	13%	51	14%
MSM & IDU	-	-	-	-	-	-	-	-	-	-	7	2%
IDU & Heterosexual contact	-	-	1	7%	-	-	-	-	-	-	11	3%
MSM & Heterosexual contact	-	-	1	7%	-	-	-	-	-	-	7	2%
MSM & IDU & Heterosexual contact	-	-	-	-	1	33%	-	-	-	-	6	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	6	2%
Other/unknown	1	9%	1	7%	-	-	1	20%	-	-	47	13%
Total	11		14		3		5		8		367	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Stark

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Stark

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	180.4	329	81%	183	81%	146	81%
Females	40.3	78	19%	43	19%	35	19%
Age at end of year							
<13	*	2	<1%	2	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	2	<1%	1	<1%	1	1%
20-24	78.5	19	5%	15	7%	4	2%
25-29	146.4	32	8%	27	12%	5	3%
30-34	152.0	33	8%	24	11%	9	5%
35-39	126.0	26	6%	14	6%	12	7%
40-44	214.0	49	12%	27	12%	22	12%
45-49	261.7	64	16%	33	15%	31	17%
50-54	250.3	69	17%	28	12%	41	23%
55-64	155.3	84	21%	41	18%	43	24%
65+	40.6	27	7%	14	6%	13	7%
Race/Ethnicity							
White, not Hispanic	87.1	285	70%	159	70%	126	70%
Black/African American, not Hispanic	357.4	103	25%	55	24%	48	27%
Hispanic/Latino	125.5	9	2%	5	2%	4	2%
Asian/Pacific Islander	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	9	2%	6	3%	3	2%
Race/Sex							
White, not Hispanic Males	150.8	239	59%	130	58%	109	60%
White, not Hispanic Females	27.3	46	11%	29	13%	17	9%
Black/African American, not Hispanic Males	516.7	73	18%	43	19%	30	17%
Black/African American, not Hispanic Females	204.2	30	7%	12	5%	18	10%
Hispanic/Latino Males	236.5	9	2%	5	2%	4	2%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	9	2%	6	3%	3	2%
Total	108.3	407		226		181	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Stark

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	241	73%	129	71%	112	77%
Injection drug use (IDU)	5	2%	-	-	5	3%
Male-to-male sexual contact and IDU	13	4%	6	3%	7	5%
Heterosexual contact	15	5%	9	5%	6	4%
Other/unknown	54	16%	38	21%	16	11%
Subtotal	328	100%	182	100%	146	100%
Female adult or adolescent						
Injection drug use	5	7%	3	8%	2	6%
Heterosexual contact	53	73%	25	63%	28	85%
Other/unknown	15	21%	12	30%	3	9%
Subtotal	73	100%	40	100%	33	100%
Child (<13 yrs at diagnosis)						
Perinatal	6	100%	4	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	6	100%	4	100%	2	100%
Total	407		226		181	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Stark

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	230	57%	125	55%	105	58%
Injection drug use (IDU) only	7	2%	2	1%	5	3%
Heterosexual contact only	68	17%	34	15%	34	19%
MSM & IDU	9	2%	5	2%	4	2%
IDU & Heterosexual contact	3	1%	1	<1%	2	1%
MSM & Heterosexual contact	11	3%	4	2%	7	4%
MSM & IDU & Heterosexual contact	4	1%	1	<1%	3	2%
Perinatal exposure	6	1%	4	2%	2	1%
Other/unknown	69	17%	50	22%	19	10%
Total	407		226		181	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Stark

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	181	76%	54	74%	3	33%	-	-	-	-	3	38%
Injection drug use (IDU)	2	1%	2	3%	1	11%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	11	5%	1	1%	1	11%	-	-	-	-	-	-
Heterosexual contact	5	2%	5	7%	1	11%	-	-	-	-	4	50%
Other/unknown	39	16%	11	15%	3	33%	-	-	-	-	1	13%
Subtotal	238	100%	73	100%	9	100%	-	-	-	-	8	100%
Female adult or adolescent												
Injection drug use	4	9%	1	4%	-	-	-	-	-	-	-	-
Heterosexual contact	28	65%	23	82%	-	-	1	100%	-	-	1	100%
Other/unknown	11	26%	4	14%	-	-	-	-	-	-	-	-
Subtotal	43	100%	28	100%	-	-	1	100%	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	4	100%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	2	100%	-	-	-	-	-	-	-	-
Total	285		103		9		1		0		9	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Stark

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	174	61%	50	49%	3	33%	-	-	-	-	3	33%
Injection drug use (IDU) only	5	2%	1	1%	1	11%	-	-	-	-	-	-
Heterosexual contact only	33	12%	28	27%	1	11%	1	100%	-	-	5	56%
MSM & IDU	8	3%	-	-	1	11%	-	-	-	-	-	-
IDU & Heterosexual contact	1	0%	2	2%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	7	2%	4	4%	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	3	1%	1	1%	-	-	-	-	-	-	-	-
Perinatal exposure	4	1%	2	2%	-	-	-	-	-	-	-	-
Other/unknown	50	18%	15	15%	3	33%	-	-	-	-	1	11%
Total	285		103		9		1		0		9	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Stark

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	262	80%	284	80%	302	80%	314	81%	329	81%
Females	65	20%	69	20%	74	20%	74	19%	78	19%
Age at end of year										
<13	1	<1%	1	<1%	2	1%	2	1%	2	<1%
13-14	-	-	-	-	1	<1%	1	<1%	-	-
15-19	4	1%	5	1%	4	1%	2	1%	2	<1%
20-24	16	5%	20	6%	18	5%	19	5%	19	5%
25-29	17	5%	22	6%	27	7%	30	8%	32	8%
30-34	18	6%	20	6%	22	6%	23	6%	33	8%
35-39	31	9%	24	7%	25	7%	28	7%	26	6%
40-44	67	20%	69	20%	66	18%	61	16%	49	12%
45-49	54	17%	57	16%	58	15%	58	15%	64	16%
50-54	57	17%	62	18%	71	19%	71	18%	69	17%
55-64	56	17%	61	17%	67	18%	72	19%	84	21%
65+	6	2%	12	3%	15	4%	21	5%	27	7%
Race/Ethnicity										
White, not Hispanic	233	71%	250	71%	269	72%	276	71%	285	70%
Black/African American, not Hispanic	78	24%	86	24%	90	24%	95	24%	103	25%
Hispanic/Latino	8	2%	8	2%	8	2%	8	2%	9	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	8	2%	9	3%	9	2%	9	2%	9	2%
Race/Sex										
White, not Hispanic Males	194	59%	209	59%	224	60%	231	60%	239	59%
White, not Hispanic Females	39	12%	41	12%	45	12%	45	12%	46	11%
Black/African American, not Hispanic Males	53	16%	59	17%	62	16%	67	17%	73	18%
Black/African American, not Hispanic Females	25	8%	27	8%	28	7%	28	7%	30	7%
Hispanic/Latino Males	8	2%	8	2%	8	2%	8	2%	9	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	8	2%	9	3%	9	2%	9	2%	9	2%
Total	327		353		376		388		407	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Stark

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	187	71%	205	72%	218	72%	230	73%	241	73%
Injection drug use (IDU)	5	2%	5	2%	5	2%	5	2%	5	2%
Male-to-male sexual contact and IDU	11	4%	11	4%	13	4%	13	4%	13	4%
Heterosexual contact	12	5%	14	5%	14	5%	14	4%	15	5%
Other/unknown	47	18%	49	17%	51	17%	51	16%	54	16%
Subtotal	262	100%	284	100%	301	100%	313	100%	328	100%
Female adult or adolescent										
Injection drug use	4	7%	5	8%	5	7%	5	7%	5	7%
Heterosexual contact	43	70%	46	71%	49	71%	49	71%	53	73%
Other/unknown	14	23%	14	22%	15	22%	15	22%	15	21%
Subtotal	61	100%	65	100%	69	100%	69	100%	73	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	100%	4	100%	6	100%	6	100%	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	4	100%	6	100%	6	100%	6	100%
Total	327		353		376		388		407	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Stark

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	177	54%	194	55%	207	55%	219	56%	230	57%
Injection drug use (IDU) only	6	2%	7	2%	7	2%	7	2%	7	2%
Heterosexual contact only	55	17%	60	17%	63	17%	63	16%	68	17%
MSM & IDU	8	2%	8	2%	9	2%	9	2%	9	2%
IDU & Heterosexual contact	3	1%	3	1%	3	1%	3	1%	3	1%
MSM & Heterosexual contact	10	3%	11	3%	11	3%	11	3%	11	3%
MSM & IDU & Heterosexual contact	3	1%	3	1%	4	1%	4	1%	4	1%
Perinatal exposure	4	1%	4	1%	6	2%	6	2%	6	1%
Other/unknown	61	19%	63	18%	66	18%	66	17%	69	17%
Total	327		353		376		388		407	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.