

Diagnoses of HIV and/or AIDS
Reported in HIV Prevention Planning
Summit

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by selected characteristics, HIV Prevention Planning Summit

Characteristic	2014 diagnosis of HIV infection			Disease Status at Initial Diagnosis						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	12.6	33	87%	29	85%	4	100%	-	-	
Females	1.8	5	13%	5	15%	-	-	-	-	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	*	3	8%	3	9%	-	-	-	-	
20-24	28.5	10	26%	10	29%	-	-	-	-	
25-29	16.8	6	16%	6	18%	-	-	-	-	
30-34	*	2	5%	1	3%	1	25%	-	-	
35-39	*	2	5%	2	6%	-	-	-	-	
40-44	*	4	11%	3	9%	1	25%	-	-	
45-49	*	3	8%	3	9%	-	-	-	-	
50-54	*	4	11%	3	9%	1	25%	-	-	
55-64	*	4	11%	3	9%	1	25%	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity										
White, not Hispanic	2.6	11	29%	8	24%	3	75%	-	-	
Black/African American, not Hispanic	29.2	23	61%	22	65%	1	25%	-	-	
Hispanic/Latino	*	3	8%	3	9%	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Unknown	*	1	3%	1	3%	-	-	-	-	
Race/Sex										
White, not Hispanic Males	5.3	11	29%	8	24%	3	75%	-	-	
White, not Hispanic Females	*	-	-	-	-	-	-	-	-	
Black/African American, not Hispanic Males	54.7	20	53%	19	56%	1	25%	-	-	
Black/African American, not Hispanic Females	*	3	8%	3	9%	-	-	-	-	
Hispanic/Latino Males	*	1	3%	1	3%	-	-	-	-	
Hispanic/Latina Females	*	2	5%	2	6%	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Unknown	*	1	3%	1	3%	-	-	-	-	
Total	7.0	38		34		4				

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by transmission category, HIV Prevention Planning Summit

Transmission Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	25	76%	22	76%	3	75%	-	-
Injection drug use (IDU)	1	3%	1	3%	-	-	-	-
Male-to-male sexual contact and IDU	1	3%	1	3%	-	-	-	-
Heterosexual contact	1	3%	1	3%	-	-	-	-
Other/unknown	5	15%	4	14%	1	25%	-	-
Subtotal	33	100%	29	100%	4	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	75%	3	75%	-	-	-	-
Other/unknown	1	25%	1	25%	-	-	-	-
Subtotal	4	100%	4	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	38		34		4		0	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by disease status at initial diagnosis and by exposure category, HIV Prevention Planning Summit

Exposure Category ^a	2014 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	25	66%	22	65%	3	75%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	11%	4	12%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	3%	1	3%	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	3%	1	3%	-	-	-	-
Perinatal exposure	1	3%	1	3%	-	-	-	-
Other/unknown	6	16%	5	15%	1	25%	-	-
Total	38		34		4		-	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and transmission category, HIV Prevention Planning Summit

Transmission Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	8	73%	15	75%	1	100%	-	-	-	-	1	100%
Injection drug use (IDU)	-	-	1	5%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	9%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	1	5%	-	-	-	-	-	-	-	-
Other/unknown	2	18%	3	15%	-	-	-	-	-	-	-	-
Subtotal	11	100%	20	100%	1	100%	-	-	-	-	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	2	67%	1	100%	-	-	-	-	-	-
Other/unknown	-	-	1	33%	-	-	-	-	-	-	-	-
Subtotal	-	-	3	100%	1	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-
Total	11		23		3		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by race/ethnicity and exposure category, HIV Prevention Planning Summit

Exposure Category ^a	2014 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	8	73%	15	65%	1	33%	-	-	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	3	13%	1	33%	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	4%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	1	9%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	1	33%	-	-	-	-	-	-
Other/unknown	2	18%	4	17%	-	-	-	-	-	-	-	-
Total	11		23		3		0		0		1	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by selected characteristics, HIV Prevention Planning Summit

Diagnosis of HIV Infection by Year

Characteristic	2010			2011			2012			2013		
	Rate ^a	No.	%									
Sex												
Males	7.6	20	77%	12.9	34	79%	11.8	31	78%	13.3	35	92%
Females	2.1	6	23%	3.2	9	21%	3.2	9	23%	*	3	8%
Age at diagnosis (yr)												
<13	*	-	-	*	2	5%	*	-	-	*	-	-
13-14	*	-	-	*	0	<1%	*	-	-	*	-	-
15-19	*	-	-	*	3	7%	17.2	6	15%	*	3	8%
20-24	*	3	12%	22.8	8	19%	14.2	5	13%	25.6	9	24%
25-29	*	2	8%	19.7	7	16%	16.8	6	15%	25.3	9	24%
30-34	24.1	8	31%	*	3	7%	27.2	9	23%	*	2	5%
35-39	*	1	4%	19.4	6	14%	*	1	3%	*	3	8%
40-44	*	-	-	*	4	9%	*	3	8%	*	4	11%
45-49	13.8	5	19%	*	1	2%	*	2	5%	*	-	-
50-54	*	3	12%	17.2	7	16%	*	4	10%	*	2	5%
55-64	*	1	4%	*	2	5%	*	2	5%	7.7	6	16%
65+	*	3	12%	*	-	-	*	2	5%	*	-	-
Race/Ethnicity												
White, not Hispanic	3.1	13	50%	4.7	20	47%	4.7	20	50%	3.5	15	39%
Black/African American, not Hispanic	12.7	10	38%	25.4	20	47%	22.9	18	45%	25.4	20	53%
Hispanic/Latino	*	-	-	*	1	2%	*	1	3%	*	1	3%
Asian/Pacific Islander	*	2	8%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	2	5%	*	1	3%	*	2	5%
Race/Sex												
White, not Hispanic Males	4.8	10	38%	7.2	15	35%	6.8	14	35%	6.8	14	37%
White, not Hispanic Females	*	3	12%	2.3	5	12%	2.8	6	15%	*	1	3%
Black/African American, not Hispanic Males	21.9	8	31%	43.8	16	37%	41.0	15	38%	49.2	18	47%
Black/African American, not Hispanic Females	*	2	8%	*	4	9%	*	3	8%	*	2	5%
Hispanic/Latino Males	*	-	-	*	1	2%	*	1	3%	*	1	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	4%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	1	4%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	4%	*	2	5%	*	1	3%	*	2	5%
Total	4.8	26		7.9	43		7.4	40		7.0	38	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by transmission category, HIV Prevention Planning Summit

Diagnosis of HIV Infection by Year

Transmission Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	13	65%	20	63%	21	68%	24	69%
Injection drug use (IDU)	1	5%	1	3%	-	-	2	6%
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-
Heterosexual contact	1	5%	1	3%	1	3%	1	3%
Other/unknown	4	20%	10	31%	9	29%	8	23%
Subtotal	20	100%	32	100%	31	100%	35	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	6	100%	6	67%	6	67%	1	33%
Other/unknown	-	-	3	33%	3	33%	2	67%
Subtotal	6	100%	9	100%	9	100%	3	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	2	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	-	-	-	-
Total	26		43		40		38	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Diagnoses of HIV infection as of June 30, 2015, by year of diagnosis and by exposure category, HIV Prevention Planning Summit

Diagnosis of HIV Infection by Year

Exposure Category ^a	2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	46%	18	42%	20	50%	24	63%
Injection drug use (IDU) only	1	4%	-	-	-	-	2	5%
Heterosexual contact only	7	27%	7	16%	7	18%	2	5%
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	2%	-	-	-	-
MSM & Heterosexual contact	1	4%	2	5%	1	3%	-	-
MSM & IDU & Heterosexual contact	1	4%	-	-	-	-	-	-
Perinatal exposure	-	-	2	5%	-	-	-	-
Other/unknown	4	15%	13	30%	12	30%	10	26%
Total	26		43		40		38	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by selected characteristics, HIV Prevention Planning Summit

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	12	71%	30	83%	26	81%	29	91%	29	85%
Females	5	29%	6	17%	6	19%	3	9%	5	15%
Age at HIV diagnosis (yr)										
<13	-	-	2	6%	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	3	8%	6	19%	3	9%	3	9%
20-24	2	12%	7	19%	5	16%	8	25%	10	29%
25-29	1	6%	5	14%	4	13%	9	28%	6	18%
30-34	4	24%	3	8%	8	25%	2	6%	1	3%
35-39	1	6%	6	17%	1	3%	2	6%	2	6%
40-44	-	-	3	8%	2	6%	2	6%	3	9%
45-49	4	24%	1	3%	2	6%	-	-	3	9%
50-54	2	12%	5	14%	1	3%	1	3%	3	9%
55-64	1	6%	1	3%	1	3%	5	16%	3	9%
65+	2	12%	-	-	2	6%	-	-	-	-
Race/Ethnicity										
White, not Hispanic	9	53%	16	44%	17	53%	10	31%	8	24%
Black/African American, not Hispanic	7	41%	18	50%	13	41%	19	59%	22	65%
Hispanic/Latino	-	-	1	3%	1	3%	1	3%	3	9%
Asian/Pacific Islander	1	6%	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	3%	1	3%	2	6%	1	3%
Race/Sex										
White, not Hispanic Males	7	41%	12	33%	12	38%	9	28%	8	24%
White, not Hispanic Females	2	12%	4	11%	5	16%	1	3%	-	-
Black/African American, not Hispanic Males	5	29%	16	44%	12	38%	17	53%	19	56%
Black/African American, not Hispanic Females	2	12%	2	6%	1	3%	2	6%	3	9%
Hispanic/Latino Males	-	-	1	3%	1	3%	1	3%	1	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	2	6%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	1	6%	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	1	3%	1	3%	2	6%	1	3%
Total	17		36		32		32		34	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by transmission category, HIV Prevention Planning Summit

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	8	67%	16	57%	18	69%	22	76%	22	76%
Injection drug use (IDU)	-	-	1	4%	-	-	-	-	1	3%
Male-to-male sexual contact and IDU	1	8%	-	-	-	-	-	-	1	3%
Heterosexual contact	-	-	1	4%	1	4%	1	3%	1	3%
Other/unknown	3	25%	10	36%	7	27%	6	21%	4	14%
Subtotal	12	100%	28	100%	26	100%	29	100%	29	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	5	100%	3	50%	5	83%	1	33%	3	75%
Other/unknown	-	-	3	50%	1	17%	2	67%	1	25%
Subtotal	5	100%	6	100%	6	100%	3	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	100%
Subtotal	-	-	2	100%	-	-	-	-	1	100%
Total	17		36		32		32		34	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported HIV (not AIDS) diagnoses as of June 30, 2015, by year of HIV diagnosis and by exposure category, HIV Prevention Planning Summit

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	7	41%	15	42%	17	53%	22	69%	22	65%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	5	29%	4	11%	6	19%	2	6%	4	12%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	1	3%	-	-	-	-	1	3%
MSM & Heterosexual contact	1	6%	1	3%	1	3%	-	-	-	-
MSM & IDU & Heterosexual contact	1	6%	-	-	-	-	-	-	1	3%
Perinatal exposure	-	-	2	6%	-	-	-	-	1	3%
Other/unknown	3	18%	13	36%	8	25%	8	25%	5	15%
Total	17		36		32		32		34	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Summit

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	9	75%	11	85%	8	73%	9	75%	9	82%	660	86%
Females	3	25%	2	15%	3	27%	3	25%	2	18%	106	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	1	<1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	1	8%	1	8%	-	-	1	8%	-	-	17	2%
20-24	1	8%	3	23%	1	9%	1	8%	1	9%	65	8%
25-29	2	17%	3	23%	2	18%	2	17%	2	18%	117	15%
30-34	2	17%	3	23%	2	18%	-	-	2	18%	168	22%
35-39	1	8%	-	-	-	-	3	25%	-	-	141	18%
40-44	2	17%	2	15%	1	9%	2	17%	2	18%	111	14%
45-49	1	8%	-	-	-	-	1	8%	1	9%	69	9%
50-54	1	8%	1	8%	4	36%	-	-	2	18%	44	6%
55-64	-	-	-	-	1	9%	2	17%	1	9%	27	4%
65+	1	8%	-	-	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	7	58%	7	54%	3	27%	7	58%	6	55%	453	59%
Black/African American, not Hispanic	3	25%	5	38%	8	73%	5	42%	5	45%	291	38%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	12	2%
Asian/Pacific Islander	1	8%	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	8%	1	8%	-	-	-	-	-	-	7	1%
Race/Sex												
White, not Hispanic Males	5	42%	7	54%	2	18%	7	58%	5	45%	412	54%
White, not Hispanic Females	2	17%	-	-	1	9%	-	-	1	9%	41	5%
Black/African American, not Hispanic Males	2	17%	3	23%	6	55%	2	17%	4	36%	230	30%
Black/African American, not Hispanic Females	1	8%	2	15%	2	18%	3	25%	1	9%	61	8%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	10	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	1	8%	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	8%	1	8%	-	-	-	-	-	-	7	1%
Total	12		13		11		12		11		766	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Summit

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	6	67%	9	82%	4	50%	5	56%	5	56%	474	72%
Injection drug use (IDU)	1	11%	1	9%	-	-	1	11%	-	-	42	6%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	11%	-	-	39	6%
Heterosexual contact	1	11%	1	9%	1	13%	-	-	1	11%	32	5%
Other/unknown	1	11%	-	-	3	38%	2	22%	3	33%	72	11%
Subtotal	9	100%	11	100%	8	100%	9	100%	9	100%	659	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	14	13%
Heterosexual contact	3	100%	2	100%	1	33%	2	67%	2	100%	75	71%
Other/unknown	-	-	-	-	2	67%	1	33%	-	-	17	16%
Subtotal	3	100%	2	100%	3	100%	3	100%	2	100%	106	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	1	100%
Total	12		13		11		12		11		766	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported cases of AIDS as of June 30, 2015, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Summit

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	6	50%	8	62%	4	36%	5	42%	5	45%	460	60%
Injection drug use (IDU) only	1	8%	-	-	-	-	1	8%	-	-	31	4%
Heterosexual contact only	4	33%	3	23%	2	18%	2	17%	3	27%	107	14%
MSM & IDU	-	-	-	-	-	-	1	8%	-	-	36	5%
IDU & Heterosexual contact	-	-	1	8%	-	-	-	-	-	-	25	3%
MSM & Heterosexual contact	-	-	1	8%	-	-	-	-	-	-	14	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	<1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	1	<1%
Other/unknown	1	8%	-	-	5	45%	3	25%	3	27%	89	12%
Total	12		13		11		12		11		766	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Summit

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Summit

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	254.0	667	81%	363	81%	304	82%
Females	54.8	153	19%	87	19%	66	18%
Age at end of year							
<13	*	3	<1%	3	1%	-	-
13-14	*	2	<1%	2	<1%	-	-
15-19	20.0	7	1%	7	2%	-	-
20-24	102.6	36	4%	32	7%	4	1%
25-29	171.3	61	7%	46	10%	15	4%
30-34	181.0	60	7%	43	10%	17	5%
35-39	213.0	66	8%	47	10%	19	5%
40-44	231.6	78	10%	41	9%	37	10%
45-49	355.4	129	16%	59	13%	70	19%
50-54	371.4	151	18%	68	15%	83	22%
55-64	224.4	174	21%	80	18%	94	25%
65+	60.4	53	6%	22	5%	31	8%
Race/Ethnicity							
White, not Hispanic	93.4	397	48%	203	45%	194	52%
Black/African American, not Hispanic	474.8	374	46%	216	48%	158	43%
Hispanic/Latino	288.7	30	4%	18	4%	12	3%
Asian/Pacific Islander	39.7	6	1%	4	1%	2	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	13	2%	9	2%	4	1%
Race/Sex							
White, not Hispanic Males	169.3	351	43%	178	40%	173	47%
White, not Hispanic Females	21.1	46	6%	25	6%	21	6%
Black/African American, not Hispanic Males	757.6	277	34%	161	36%	116	31%
Black/African American, not Hispanic Females	229.8	97	12%	55	12%	42	11%
Hispanic/Latino Males	487.1	25	3%	14	3%	11	3%
Hispanic/Latina Females	95.1	5	1%	4	1%	1	<1%
Asian/Pacific Islander Males	*	3	<1%	3	1%	-	-
Asian/Pacific Islander Females	*	3	<1%	1	<1%	2	1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	13	2%	9	2%	4	1%
Total	151.3	820		450		370	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Summit

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	453	69%	239	67%	214	71%
Injection drug use (IDU)	23	4%	6	2%	17	6%
Male-to-male sexual contact and IDU	30	5%	10	3%	20	7%
Heterosexual contact	29	4%	12	3%	17	6%
Other/unknown	122	19%	88	25%	34	11%
Subtotal	657	100%	355	100%	302	100%
Female adult or adolescent						
Injection drug use	14	9%	6	7%	8	12%
Heterosexual contact	99	66%	50	59%	49	74%
Other/unknown	38	25%	29	34%	9	14%
Subtotal	151	100%	85	100%	66	100%
Child (<13 yrs at diagnosis)						
Perinatal	12	100%	10	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	12	100%	10	100%	2	100%
Total	820		450		370	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Summit

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	438	53%	233	52%	205	55%
Injection drug use (IDU) only	21	3%	8	2%	13	4%
Heterosexual contact only	128	16%	62	14%	66	18%
MSM & IDU	22	3%	5	1%	17	5%
IDU & Heterosexual contact	16	2%	4	1%	12	3%
MSM & Heterosexual contact	15	2%	6	1%	9	2%
MSM & IDU & Heterosexual contact	8	1%	5	1%	3	1%
Perinatal exposure	12	1%	10	2%	2	1%
Other/unknown	160	20%	117	26%	43	12%
Total	820		450		370	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Summit

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	259	74%	168	62%	19	83%	2	67%	-	-	5	45%
Injection drug use (IDU)	10	3%	11	4%	1	4%	-	-	-	-	1	9%
Male-to-male sexual contact and IDU	18	5%	12	4%	-	-	-	-	-	-	-	-
Heterosexual contact	11	3%	15	6%	1	4%	1	33%	-	-	1	9%
Other/unknown	52	15%	64	24%	2	9%	-	-	-	-	4	36%
Subtotal	350	100%	270	100%	23	100%	3	100%	-	-	11	100%
Female adult or adolescent												
Injection drug use	2	4%	10	11%	1	20%	-	-	-	-	1	50%
Heterosexual contact	31	67%	62	65%	4	80%	2	67%	-	-	-	-
Other/unknown	13	28%	23	24%	-	-	1	33%	-	-	1	50%
Subtotal	46	100%	95	100%	5	100%	3	100%	-	-	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	9	100%	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	9	100%	2	100%	-	-	-	-	-	-
Total	397		374		30		6		0		13	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Summit

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	253	64%	160	43%	19	63%	1	17%	-	-	5	38%
Injection drug use (IDU) only	9	2%	11	3%	1	3%	-	-	-	-	-	-
Heterosexual contact only	42	11%	77	21%	5	17%	3	50%	-	-	1	8%
MSM & IDU	13	3%	9	2%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	3	1%	10	3%	1	3%	-	-	-	-	2	15%
MSM & Heterosexual contact	6	2%	8	2%	-	-	1	17%	-	-	-	-
MSM & IDU & Heterosexual contact	5	1%	3	1%	-	-	-	-	-	-	-	-
Perinatal exposure	1	0%	9	2%	2	7%	-	-	-	-	-	-
Other/unknown	65	16%	87	23%	2	7%	1	17%	-	-	5	38%
Total	397		374		30		6		0		13	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Summit

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	543	81%	572	80%	601	80%	635	81%	667	81%
Females	130	19%	139	20%	147	20%	150	19%	153	19%
Age at end of year										
<13	2	<1%	4	1%	5	1%	5	1%	3	<1%
13-14	2	<1%	1	<1%	-	-	-	-	2	<1%
15-19	5	1%	6	1%	9	1%	6	1%	7	1%
20-24	27	4%	29	4%	26	3%	33	4%	36	4%
25-29	38	6%	49	7%	56	7%	63	8%	61	7%
30-34	55	8%	51	7%	53	7%	56	7%	60	7%
35-39	56	8%	61	9%	61	8%	62	8%	66	8%
40-44	112	17%	101	14%	95	13%	84	11%	78	10%
45-49	144	21%	146	21%	144	19%	136	17%	129	16%
50-54	104	15%	114	16%	135	18%	143	18%	151	18%
55-64	109	16%	121	17%	128	17%	154	20%	174	21%
65+	19	3%	28	4%	36	5%	43	5%	53	6%
Race/Ethnicity										
White, not Hispanic	343	51%	358	50%	376	50%	389	50%	397	48%
Black/African American, not Hispanic	295	44%	314	44%	330	44%	351	45%	374	46%
Hispanic/Latino	23	3%	24	3%	26	3%	27	3%	30	4%
Asian/Pacific Islander	5	1%	6	1%	6	1%	6	1%	6	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	7	1%	9	1%	10	1%	12	2%	13	2%
Race/Sex										
White, not Hispanic Males	308	46%	319	45%	331	44%	343	44%	351	43%
White, not Hispanic Females	35	5%	39	5%	45	6%	46	6%	46	6%
Black/African American, not Hispanic Males	208	31%	223	31%	237	32%	256	33%	277	34%
Black/African American, not Hispanic Females	87	13%	91	13%	93	12%	95	12%	97	12%
Hispanic/Latino Males	19	3%	20	3%	22	3%	23	3%	25	3%
Hispanic/Latina Females	4	1%	4	1%	4	1%	4	1%	5	1%
Asian/Pacific Islander Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	2	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	7	1%	9	1%	10	1%	12	2%	13	2%
Total	673		711		748		785		820	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Summit

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	369	69%	386	69%	405	69%	428	68%	453	69%
Injection drug use (IDU)	19	4%	20	4%	20	3%	22	4%	23	4%
Male-to-male sexual contact and IDU	28	5%	29	5%	29	5%	29	5%	30	5%
Heterosexual contact	25	5%	26	5%	27	5%	28	4%	29	4%
Other/unknown	95	18%	102	18%	110	19%	118	19%	122	19%
Subtotal	536	100%	563	100%	591	100%	625	100%	657	100%
Female adult or adolescent										
Injection drug use	14	11%	14	10%	14	10%	14	9%	14	9%
Heterosexual contact	85	66%	91	66%	96	66%	97	66%	99	66%
Other/unknown	29	23%	32	23%	35	24%	37	25%	38	25%
Subtotal	128	100%	137	100%	145	100%	148	100%	151	100%
Child (<13 yrs at diagnosis)										
Perinatal	9	100%	11	100%	12	100%	12	100%	12	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	9	100%	11	100%	12	100%	12	100%	12	100%
Total	673		711		748		785		820	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Summit

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	355	53%	371	52%	390	52%	413	53%	438	53%
Injection drug use (IDU) only	20	3%	20	3%	20	3%	21	3%	21	3%
Heterosexual contact only	110	16%	117	16%	123	16%	125	16%	128	16%
MSM & IDU	21	3%	22	3%	22	3%	22	3%	22	3%
IDU & Heterosexual contact	13	2%	14	2%	14	2%	15	2%	16	2%
MSM & Heterosexual contact	14	2%	15	2%	15	2%	15	2%	15	2%
MSM & IDU & Heterosexual contact	7	1%	7	1%	7	1%	7	1%	8	1%
Perinatal exposure	9	1%	11	2%	12	2%	12	2%	12	1%
Other/unknown	124	18%	134	19%	145	19%	155	20%	160	20%
Total	673		711		748		785		820	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.