

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Trumbull

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Trumbull

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	155.7	156	78%	78	72%	78	86%
Females	41.9	44	22%	31	28%	13	14%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	48.7	6	3%	4	4%	2	2%
25-29	140.1	16	8%	10	9%	6	7%
30-34	126.8	14	7%	13	12%	1	1%
35-39	154.0	17	9%	11	10%	6	7%
40-44	170.1	21	11%	11	10%	10	11%
45-49	268.8	36	18%	18	17%	18	20%
50-54	183.0	28	14%	14	13%	14	15%
55-64	153.6	48	24%	20	18%	28	31%
65+	35.3	14	7%	8	7%	6	7%
Race/Ethnicity							
White, not Hispanic	64.0	115	58%	65	60%	50	55%
Black/African American, not Hispanic	390.1	67	34%	35	32%	32	35%
Hispanic/Latino	207.3	7	4%	2	2%	5	5%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	1%	-	-	1	1%
Unknown	*	10	5%	7	6%	3	3%
Race/Sex							
White, not Hispanic Males	108.3	95	48%	50	46%	45	49%
White, not Hispanic Females	21.8	20	10%	15	14%	5	5%
Black/African American, not Hispanic Males	545.7	46	23%	20	18%	26	29%
Black/African American, not Hispanic Females	240.2	21	11%	15	14%	6	7%
Hispanic/Latino Males	417.4	7	4%	2	2%	5	5%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	1	1%	-	-	1	1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	10	5%	7	6%	3	3%
Total	97.5	200		109		91	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Trumbull

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	104	68%	47	61%	57	74%
Injection drug use (IDU)	10	6%	4	5%	6	8%
Male-to-male sexual contact and IDU	8	5%	4	5%	4	5%
Heterosexual contact	12	8%	8	10%	4	5%
Other/unknown	20	13%	14	18%	6	8%
Subtotal	154	100%	77	100%	77	100%
Female adult or adolescent						
Injection drug use	5	12%	3	10%	2	17%
Heterosexual contact	24	56%	17	55%	7	58%
Other/unknown	14	33%	11	35%	3	25%
Subtotal	43	100%	31	100%	12	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	67%	-	-	2	100%
Other/unknown	1	33%	1	100%	-	-
Subtotal	3	100%	1	100%	2	100%
Total	200		109		91	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Trumbull

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	97	49%	44	40%	53	58%
Injection drug use (IDU) only	12	6%	6	6%	6	7%
Heterosexual contact only	36	18%	25	23%	11	12%
MSM & IDU	6	3%	2	2%	4	4%
IDU & Heterosexual contact	3	2%	1	1%	2	2%
MSM & Heterosexual contact	7	4%	3	3%	4	4%
MSM & IDU & Heterosexual contact	2	1%	2	2%	-	-
Perinatal exposure	2	1%	-	-	2	2%
Other/unknown	35	18%	26	24%	9	10%
Total	200		109		91	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Trumbull

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	64	67%	29	64%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	4	4%	5	11%	-	-	-	-	<10	*	-	-
Male-to-male sexual contact and IDU	6	6%	2	4%	-	-	-	-	-	-	-	-
Heterosexual contact	8	8%	4	9%	-	-	-	-	-	-	-	-
Other/unknown	13	14%	5	11%	<10	*	-	-	-	-	<10	*
Subtotal	95	100%	45	100%	<10	*	-	-	<10	*	<10	*
Female adult or adolescent												
Injection drug use	1	5%	4	19%	-	-	-	-	-	-	-	-
Heterosexual contact	15	75%	9	43%	-	-	-	-	-	-	-	-
Other/unknown	4	20%	8	38%	-	-	-	-	-	-	<10	*
Subtotal	20	100%	21	100%	-	-	-	-	-	-	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	<10	*	-	-	-	-	<10	*
Other/unknown	-	-	1	100%	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	<10	*	-	-	-	-	<10	*
Total	115		67		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Trumbull

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	61	53%	27	40%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	5	4%	6	9%	-	-	-	-	<10	*	-	-
Heterosexual contact only	23	20%	13	19%	-	-	-	-	-	-	-	-
MSM & IDU	4	3%	2	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	3	4%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	3	3%	2	3%	-	-	-	-	-	-	<10	*
MSM & IDU & Heterosexual contact	2	2%	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	<10	*	-	-	-	-	<10	*
Other/unknown	17	15%	14	21%	<10	*	-	-	-	-	<10	*
Total	115		67		<10		0		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Trumbull

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	136	80%	144	79%	145	79%	150	78%	156	78%
Females	35	20%	38	21%	39	21%	43	22%	44	22%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	2%	1	1%	-	-	-	-	-	-
20-24	7	4%	10	5%	7	4%	8	4%	6	3%
25-29	12	7%	13	7%	16	9%	15	8%	16	8%
30-34	13	8%	13	7%	11	6%	13	7%	14	7%
35-39	19	11%	22	12%	22	12%	18	9%	17	9%
40-44	27	16%	24	13%	23	13%	22	11%	21	11%
45-49	27	16%	29	16%	29	16%	35	18%	36	18%
50-54	26	15%	28	15%	30	16%	27	14%	28	14%
55-64	31	18%	36	20%	40	22%	44	23%	48	24%
65+	6	4%	6	3%	6	3%	11	6%	14	7%
Race/Ethnicity										
White, not Hispanic	96	56%	102	56%	103	56%	108	56%	115	58%
Black/African American, not Hispanic	60	35%	64	35%	64	35%	67	35%	67	34%
Hispanic/Latino	6	4%	6	3%	7	4%	7	4%	7	4%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	8	5%	9	5%	9	5%	10	5%	10	5%
Race/Sex										
White, not Hispanic Males	81	47%	86	47%	86	47%	89	46%	95	48%
White, not Hispanic Females	15	9%	16	9%	17	9%	19	10%	20	10%
Black/African American, not Hispanic Males	43	25%	45	25%	45	24%	46	24%	46	23%
Black/African American, not Hispanic Females	17	10%	19	10%	19	10%	21	11%	21	11%
Hispanic/Latino Males	6	4%	6	3%	7	4%	7	4%	7	4%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	1%	1	1%	1	1%	1	1%	1	1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	8	5%	9	5%	9	5%	10	5%	10	5%
Total	171		182		184		193		200	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Trumbull

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	89	66%	97	68%	98	69%	100	68%	104	68%
Injection drug use (IDU)	9	7%	9	6%	9	6%	9	6%	10	6%
Male-to-male sexual contact and IDU	8	6%	8	6%	8	6%	8	5%	8	5%
Heterosexual contact	10	7%	10	7%	10	7%	11	7%	12	8%
Other/unknown	18	13%	18	13%	18	13%	20	14%	20	13%
Subtotal	134	100%	142	100%	143	100%	148	100%	154	100%
Female adult or adolescent										
Injection drug use	4	12%	5	14%	5	13%	5	12%	5	12%
Heterosexual contact	18	53%	18	49%	19	50%	23	55%	24	56%
Other/unknown	12	35%	14	38%	14	37%	14	33%	14	33%
Subtotal	34	100%	37	100%	38	100%	42	100%	43	100%
Child (<13 yrs at diagnosis)										
Perinatal	2	67%	2	67%	2	67%	2	67%	2	67%
Other/unknown	1	33%	1	33%	1	33%	1	33%	1	33%
Subtotal	3	100%	3	100%	3	100%	3	100%	3	100%
Total	171		182		184		193		200	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Trumbull

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	83	49%	90	49%	91	49%	93	48%	97	49%
Injection drug use (IDU) only	10	6%	11	6%	11	6%	11	6%	12	6%
Heterosexual contact only	28	16%	28	15%	29	16%	34	18%	36	18%
MSM & IDU	6	4%	6	3%	6	3%	6	3%	6	3%
IDU & Heterosexual contact	3	2%	3	2%	3	2%	3	2%	3	2%
MSM & Heterosexual contact	6	4%	7	4%	7	4%	7	4%	7	4%
MSM & IDU & Heterosexual contact	2	1%	2	1%	2	1%	2	1%	2	1%
Perinatal exposure	2	1%	2	1%	2	1%	2	1%	2	1%
Other/unknown	31	18%	33	18%	33	18%	35	18%	35	18%
Total	171		182		184		193		200	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.