

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Union

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Union**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	82.2	21	41%	8	31%	13	52%
Females	106.3	30	59%	18	69%	12	48%
<b>Age at end of year</b>							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	2	4%	2	8%	-	-
25-29	*	2	4%	2	8%	-	-
30-34	166.5	6	12%	4	15%	2	8%
35-39	*	4	8%	1	4%	3	12%
40-44	160.0	7	14%	4	15%	3	12%
45-49	265.0	11	22%	3	12%	8	32%
50-54	217.0	9	18%	4	15%	5	20%
55-64	128.3	8	16%	6	23%	2	8%
65+	*	2	4%	-	-	2	8%
<b>Race/Ethnicity</b>							
White, not Hispanic	79.6	39	76%	19	73%	20	80%
Black/African American, not Hispanic	568.2	8	16%	5	19%	3	12%
Hispanic/Latino	*	3	6%	1	4%	2	8%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	1	2%	1	4%	-	-
<b>Race/Sex</b>							
White, not Hispanic Males	76.4	18	35%	7	27%	11	44%
White, not Hispanic Females	82.5	21	41%	12	46%	9	36%
Black/African American, not Hispanic Males	*	1	2%	-	-	1	4%
Black/African American, not Hispanic Females	636.9	7	14%	5	19%	2	8%
Hispanic/Latino Males	*	1	2%	-	-	1	4%
Hispanic/Latina Females	*	2	4%	1	4%	1	4%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	1	2%	1	4%	-	-
<b>Total</b>	<b>94.8</b>	<b>51</b>		<b>26</b>		<b>25</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Union**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	14	67%	6	75%	8	62%
Injection drug use (IDU)	2	10%	-	-	2	15%
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	2	10%	-	-	2	15%
Other/unknown	3	14%	2	25%	1	8%
Subtotal	21	100%	8	100%	13	100%
<b>Female adult or adolescent</b>						
Injection drug use	12	40%	6	33%	6	50%
Heterosexual contact	14	47%	9	50%	5	42%
Other/unknown	4	13%	3	17%	1	8%
Subtotal	30	100%	18	100%	12	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>51</b>		<b>26</b>		<b>25</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Union**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	14	27%	6	23%	8	32%
Injection drug use (IDU) only	6	12%	3	12%	3	12%
Heterosexual contact only	16	31%	9	35%	7	28%
MSM & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	8	16%	3	12%	5	20%
MSM & Heterosexual contact	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	7	14%	5	19%	2	8%
<b>Total</b>	<b>51</b>		<b>26</b>		<b>25</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Union**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	14	78%	-	-	-	-	-	-	-	-	-	-
Injection drug use (IDU)	1	6%	-	-	<10	*	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	6%	<10	*	-	-	-	-	-	-	-	-
Other/unknown	2	11%	-	-	-	-	-	-	-	-	<10	*
Subtotal	18	100%	<10	*	<10	*	-	-	-	-	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	7	33%	<10	*	<10	*	-	-	-	-	-	-
Heterosexual contact	10	48%	<10	*	<10	*	-	-	-	-	-	-
Other/unknown	4	19%	-	-	-	-	-	-	-	-	-	-
Subtotal	21	100%	<10	*	<10	*	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>39</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Union**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	14	36%	-	-	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	3	8%	<10	*	<10	*	-	-	-	-	-	-
Heterosexual contact only	11	28%	<10	*	<10	*	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	5	13%	<10	*	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	15%	-	-	-	-	-	-	-	-	<10	*
<b>Total</b>	<b>39</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>0</b>		<b>0</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Union**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	19	42%	20	43%	21	43%	21	42%	21	41%
Females	26	58%	26	57%	28	57%	29	58%	30	59%
<b>Age at end of year</b>										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	1	2%	1	2%	-	-	-	-
20-24	2	4%	1	2%	-	-	1	2%	2	4%
25-29	4	9%	4	9%	4	8%	4	8%	2	4%
30-34	2	4%	1	2%	4	8%	4	8%	6	12%
35-39	8	18%	8	17%	6	12%	6	12%	4	8%
40-44	10	22%	9	20%	11	22%	7	14%	7	14%
45-49	6	13%	9	20%	8	16%	13	26%	11	22%
50-54	9	20%	7	15%	9	18%	9	18%	9	18%
55-64	2	4%	4	9%	4	8%	4	8%	8	16%
65+	2	4%	2	4%	2	4%	2	4%	2	4%
<b>Race/Ethnicity</b>										
White, not Hispanic	34	76%	35	76%	38	78%	39	78%	39	76%
Black/African American, not Hispanic	8	18%	8	17%	8	16%	8	16%	8	16%
Hispanic/Latino	2	4%	2	4%	2	4%	2	4%	3	6%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	2%	1	2%	1	2%
<b>Race/Sex</b>										
White, not Hispanic Males	16	36%	17	37%	18	37%	18	36%	18	35%
White, not Hispanic Females	18	40%	18	39%	20	41%	21	42%	21	41%
Black/African American, not Hispanic Males	1	2%	1	2%	1	2%	1	2%	1	2%
Black/African American, not Hispanic Females	7	16%	7	15%	7	14%	7	14%	7	14%
Hispanic/Latino Males	1	2%	1	2%	1	2%	1	2%	1	2%
Hispanic/Latina Females	1	2%	1	2%	1	2%	1	2%	2	4%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	2%	1	2%	1	2%	1	2%	1	2%
<b>Total</b>	<b>45</b>		<b>46</b>		<b>49</b>		<b>50</b>		<b>51</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Union

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	12	63%	13	65%	14	67%	14	67%	14	67%
Injection drug use (IDU)	2	11%	2	10%	2	10%	2	10%	2	10%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	2	11%	2	10%	2	10%	2	10%	2	10%
Other/unknown	3	16%	3	15%	3	14%	3	14%	3	14%
Subtotal	19	100%	20	100%	21	100%	21	100%	21	100%
<b>Female adult or adolescent</b>										
Injection drug use	9	35%	9	35%	10	36%	11	38%	12	40%
Heterosexual contact	14	54%	14	54%	14	50%	14	48%	14	47%
Other/unknown	3	12%	3	12%	4	14%	4	14%	4	13%
Subtotal	26	100%	26	100%	28	100%	29	100%	30	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>45</b>		<b>46</b>		<b>49</b>		<b>50</b>		<b>51</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Union

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	12	27%	13	28%	14	29%	14	28%	14	27%
Injection drug use (IDU) only	5	11%	5	11%	5	10%	5	10%	6	12%
Heterosexual contact only	16	36%	16	35%	16	33%	16	32%	16	31%
MSM & IDU	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	6	13%	6	13%	7	14%	8	16%	8	16%
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	6	13%	6	13%	7	14%	7	14%	7	14%
<b>Total</b>	<b>45</b>		<b>46</b>		<b>49</b>		<b>50</b>		<b>51</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.