

Persons Living with a Diagnosis of HIV  
and/or AIDS Reported in HIV  
Prevention Planning Warren

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Warren**

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	105.2	117	82%	55	82%	62	82%
Females	23.5	26	18%	12	18%	14	18%
<b>Age at end of year</b>							
<13	*	3	2%	3	4%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	50.3	6	4%	4	6%	2	3%
25-29	131.7	15	10%	11	16%	4	5%
30-34	111.6	15	10%	12	18%	3	4%
35-39	48.0	7	5%	3	4%	4	5%
40-44	94.4	16	11%	6	9%	10	13%
45-49	162.0	28	20%	12	18%	16	21%
50-54	159.4	28	20%	10	15%	18	24%
55-64	73.7	20	14%	5	7%	15	20%
65+	17.3	5	3%	1	1%	4	5%
<b>Race/Ethnicity</b>							
White, not Hispanic	51.4	100	70%	43	64%	57	75%
Black/African American, not Hispanic	386.5	29	20%	16	24%	13	17%
Hispanic/Latino	158.8	9	6%	5	7%	4	5%
Asian/Pacific Islander	*	2	1%	2	3%	-	-
American Indian/Alaska Native	*	1	1%	-	-	1	1%
Unknown	*	2	1%	1	1%	1	1%
<b>Race/Sex</b>							
White, not Hispanic Males	84.8	82	57%	36	54%	46	61%
White, not Hispanic Females	18.4	18	13%	7	10%	11	14%
Black/African American, not Hispanic Males	450.5	22	15%	12	18%	10	13%
Black/African American, not Hispanic Females	267.1	7	5%	4	6%	3	4%
Hispanic/Latino Males	315.2	9	6%	5	7%	4	5%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-
Asian/Pacific Islander Females	*	1	1%	1	1%	-	-
American Indian/Alaska Native Males	*	1	1%	-	-	1	1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	2	1%	1	1%	1	1%
<b>Total</b>	<b>64.5</b>	<b>143</b>		<b>67</b>		<b>76</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count < 5 due to unstable rates.

<sup>a</sup> The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Warren**

Transmission Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	79	69%	36	68%	43	69%
Injection drug use (IDU)	6	5%	2	4%	4	6%
Male-to-male sexual contact and IDU	1	1%	1	2%	-	-
Heterosexual contact	8	7%	3	6%	5	8%
Other/unknown	21	18%	11	21%	10	16%
Subtotal	115	100%	53	100%	62	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	13%	2	20%	1	7%
Heterosexual contact	16	67%	4	40%	12	86%
Other/unknown	5	21%	4	40%	1	7%
Subtotal	24	100%	10	100%	14	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	2	50%	2	50%	-	-
Other/unknown	2	50%	2	50%	-	-
Subtotal	4	100%	4	100%	-	-
<b>Total</b>	<b>143</b>		<b>67</b>		<b>76</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Warren**

Exposure Category <sup>a</sup>	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	74	52%	34	51%	40	53%
Injection drug use (IDU) only	5	3%	2	3%	3	4%
Heterosexual contact only	24	17%	7	10%	17	22%
MSM & IDU	1	1%	1	1%	-	-
IDU & Heterosexual contact	4	3%	2	3%	2	3%
MSM & Heterosexual contact	5	3%	2	3%	3	4%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	2	1%	2	3%	-	-
Other/unknown	28	20%	17	25%	11	14%
<b>Total</b>	<b>143</b>		<b>67</b>		<b>76</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Warren**

Transmission Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	60	73%	11	52%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU)	3	4%	1	5%	<10	*	-	-	<10	*	-	-
Male-to-male sexual contact and IDU	-	-	1	5%	-	-	-	-	-	-	-	-
Heterosexual contact	2	2%	5	24%	<10	*	-	-	-	-	-	-
Other/unknown	17	21%	3	14%	<10	*	-	-	-	-	-	-
Subtotal	82	100%	21	100%	<10	*	-	-	<10	*	<10	*
<b>Female adult or adolescent</b>												
Injection drug use	3	17%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	12	67%	4	67%	-	-	-	-	-	-	-	-
Other/unknown	3	17%	2	33%	-	-	-	-	-	-	-	-
Subtotal	18	100%	6	100%	-	-	-	-	-	-	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	<10	*	-	-	-	-
Subtotal	-	-	2	100%	-	-	<10	*	-	-	-	-
<b>Total</b>	<b>100</b>		<b>29</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Warren**

Exposure Category <sup>a</sup>	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	56	56%	11	38%	<10	*	-	-	-	-	<10	*
Injection drug use (IDU) only	3	3%	1	3%	-	-	-	-	<10	*	-	-
Heterosexual contact only	14	14%	9	31%	<10	*	-	-	-	-	-	-
MSM & IDU	-	-	1	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	3	3%	-	-	<10	*	-	-	-	-	-	-
MSM & Heterosexual contact	4	4%	-	-	<10	*	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	2	7%	-	-	-	-	-	-	-	-
Other/unknown	20	20%	5	17%	<10	*	<10	*	-	-	-	-
<b>Total</b>	<b>100</b>		<b>29</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>&lt;10</b>		<b>&lt;10</b>	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (\*) indicates percent not calculated for column < 10 cases due to small numbers.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

**Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Warren**

**Living with a Diagnosis of HIV Infection**

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
<b>Sex</b>										
Males	92	83%	101	83%	103	82%	111	82%	117	82%
Females	19	17%	21	17%	22	18%	24	18%	26	18%
<b>Age at end of year</b>										
<13	-	-	-	-	-	-	2	1%	3	2%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	1%	1	1%	1	1%	-	-	-	-
20-24	5	5%	8	7%	8	6%	6	4%	6	4%
25-29	12	11%	13	11%	11	9%	14	10%	15	10%
30-34	7	6%	8	7%	12	10%	13	10%	15	10%
35-39	14	13%	10	8%	10	8%	7	5%	7	5%
40-44	21	19%	21	17%	17	14%	20	15%	16	11%
45-49	22	20%	27	22%	24	19%	24	18%	28	20%
50-54	14	13%	18	15%	23	18%	28	21%	28	20%
55-64	14	13%	14	11%	16	13%	17	13%	20	14%
65+	1	1%	2	2%	3	2%	4	3%	5	3%
<b>Race/Ethnicity</b>										
White, not Hispanic	85	77%	89	73%	91	73%	98	73%	100	70%
Black/African American, not Hispanic	20	18%	26	21%	27	22%	27	20%	29	20%
Hispanic/Latino	4	4%	5	4%	5	4%	6	4%	9	6%
Asian/Pacific Islander	-	-	-	-	-	-	2	1%	2	1%
American Indian/Alaska Native	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	1	1%	1	1%	1	1%	1	1%	2	1%
<b>Race/Sex</b>										
White, not Hispanic Males	69	62%	73	60%	74	59%	80	59%	82	57%
White, not Hispanic Females	16	14%	16	13%	17	14%	18	13%	18	13%
Black/African American, not Hispanic Males	17	15%	21	17%	22	18%	22	16%	22	15%
Black/African American, not Hispanic Females	3	3%	5	4%	5	4%	5	4%	7	5%
Hispanic/Latino Males	4	4%	5	4%	5	4%	6	4%	9	6%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	1	1%	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	1	1%	1	1%
American Indian/Alaska Native Males	1	1%	1	1%	1	1%	1	1%	1	1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	1	1%	1	1%	1	1%	1	1%	2	1%
<b>Total</b>	<b>111</b>		<b>122</b>		<b>125</b>		<b>135</b>		<b>143</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Warren

### Living with a Diagnosis of HIV Infection

Transmission Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	59	65%	67	67%	68	67%	74	68%	79	69%
Injection drug use (IDU)	6	7%	6	6%	6	6%	6	6%	6	5%
Male-to-male sexual contact and IDU	1	1%	1	1%	1	1%	1	1%	1	1%
Heterosexual contact	7	8%	8	8%	8	8%	8	7%	8	7%
Other/unknown	18	20%	18	18%	19	19%	20	18%	21	18%
Subtotal	91	100%	100	100%	102	100%	109	100%	115	100%
<b>Female adult or adolescent</b>										
Injection drug use	1	5%	1	5%	2	9%	3	13%	3	13%
Heterosexual contact	15	79%	16	76%	16	73%	16	70%	16	67%
Other/unknown	3	16%	4	19%	4	18%	4	17%	5	21%
Subtotal	19	100%	21	100%	22	100%	23	100%	24	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	1	100%	1	100%	1	100%	1	33%	2	50%
Other/unknown	-	-	-	-	-	-	2	67%	2	50%
Subtotal	1	100%	1	100%	1	100%	3	100%	4	100%
<b>Total</b>	<b>111</b>		<b>122</b>		<b>125</b>		<b>135</b>		<b>143</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

## Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Warren

### Living with a Diagnosis of HIV Infection

Exposure Category <sup>a</sup>	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	55	50%	63	52%	64	51%	70	52%	74	52%
Injection drug use (IDU) only	4	4%	4	3%	5	4%	5	4%	5	3%
Heterosexual contact only	22	20%	24	20%	24	19%	24	18%	24	17%
MSM & IDU	1	1%	1	1%	1	1%	1	1%	1	1%
IDU & Heterosexual contact	3	3%	3	2%	3	2%	4	3%	4	3%
MSM & Heterosexual contact	4	4%	4	3%	4	3%	4	3%	5	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	1	1%	1	1%	1	1%	2	1%
Other/unknown	21	19%	22	18%	23	18%	26	19%	28	20%
<b>Total</b>	<b>111</b>		<b>122</b>		<b>125</b>		<b>135</b>		<b>143</b>	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

<sup>a</sup>Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.