



Ohio Department of Health
246 North High Street
Columbus, Ohio 43215
www.odh.ohio.gov

Bob Taft, Governor
J. Nick Baird M.D., Director

N E W S

For Immediate Release
August 29, 2006

Office of Public Affairs • (614) 644-8562
Jay Carey, Director

RACCOON RABIES VACCINATION BAITING SLATED FOR TUESDAY

Operations will begin Sept. 5

COLUMBUS – The Ohio Department of Health (ODH) will once again participate in a multi-state effort coordinated by the U.S. Department of Agriculture’s Animal and Plant Health Inspection Service to reduce the spread of raccoon strain rabies.

Rabies is a viral disease that affects mammals and people. It is almost always fatal. Since the mid-1970s, a strain of rabies associated with raccoons has spread rapidly through the eastern United States and first threatened northeastern Ohio in 1997. To control the disease, ODH has been conducting a program to distribute an oral rabies vaccine-laden bait to immunize wild raccoons along the Pennsylvania and West Virginia border.

The northern portion of the Appalachian Ridge Oral Rabies Vaccination (ORV) baiting begins Sept. 5 in Ohio, weather permitting, and will cover 4,682 square miles of the state’s northeastern and eastern border. Aerial distribution should be completed within 10 days; ground baiting may continue through Sept. 22.

As in past years, baiting will take place in Ashtabula, Belmont, Columbiana, Jefferson, Mahoning and Trumbull counties and parts of Carroll, Harrison and Monroe counties. Baiting teams also will operate in Geauga and Lake counties, plus parts of Cuyahoga, Portage and Summit counties due to the 35 raccoon-strain rabies cases reported in Cuyahoga, Geauga and Lake counties in 2005.

“Rabies baiting is an important tool for limiting the spread of raccoon-strain rabies in Ohio,” said ODH Director J. Nick Baird, M.D. “By partnering with other states and local health departments, we will immunize millions of raccoons.”

Ohio’s partners in the multi-state baiting are: Maryland, North Carolina, Pennsylvania, Tennessee, Virginia and West Virginia, in what is known as the Appalachian Ridge ORV program. The seven-state effort will involve distribution of about 5 million baits and cover more than 26,000 square miles. ODH has participated in the program since 1997 and has dropped more than 8.6 million baits in that time.

-MORE-

ODH Rabies Baiting

Page 2 of 2

Two types of baits will be used. Airplanes will drop a small plastic sachet, about the size of a ketchup packet, coated in fishmeal. In urban areas, the vaccine will be inside a hard, brown 2-x-2-inch fishmeal block which will be distributed by vehicles staffed by local health departments and other local agency volunteers. Most of the 808,176 baits will be distributed by air, with the use of specially equipped planes from the Ontario Ministry of Natural Resources and a helicopter from the Ohio Department of Transportation.

Residents should avoid the baits and keep pets confined during the baiting period. Dogs in particular are attracted to the baits and will occasionally eat them. The baits are not harmful to pets. Please keep the following information in mind:

- Know what the baits look like. The coated sachet, which will be distributed by airplanes, is about the size of a ketchup packet. It is white and rolled in a brown fishmeal glaze. In urban areas, where baits will be distributed by vehicle, the sachet will be inside a hard, brown fishmeal block, about 2-x-2-x-1-inch square.
- Instruct children to leave the baits alone.
- Once your area is baited, keep dogs and cats inside or on leashes for up to five days. Most baits disappear within 24 hours; however, it is important raccoons have every opportunity to eat them.
- Do not attempt to take bait away from your pet; you may be bitten.
- Anyone handling baits should wear gloves. If baits are found in areas frequented by pets or children, toss them into deeper cover. Damaged baits can be disposed of in the trash.
- If a person is exposed to the vaccine (red liquid), thoroughly wash any areas of the skin that came into contact with the vaccine with soap and water.
- If someone has been exposed to the vaccine or has questions about the baiting, call the local health department or ODH's information line at 1-888-RABIES-1.

The rabies virus is found in the saliva of affected animals, most often raccoons, skunks and bats, and is spread by a bite or scratch. Bats, raccoons and skunks pose the greatest risk of rabies in Ohio. To protect your family against this still-deadly disease:

- Avoid contact with wild animals and animals you do not know.
- Vaccinate your pets against rabies and keep them current on their shots.
- If bitten, call your doctor. If your pet has contact with a wild animal, call your veterinarian. Rabies exposures should also be reported to your local health department.

-30-

Contact: Ohio Department of Health – Office of Public Affairs (614) 644-8562

Editors: Please note the following from the Ohio Department of Natural Resources:

“According to Ohio Administrative Code 1501:31-15-03, it is illegal to live trap, move and release raccoons. Persons dealing with a nuisance raccoon cannot relocate the animal. There are only two legal options for nuisance raccoons; euthanize or release on the same property where they were live trapped. Uninformed people who are relocating nuisance raccoons may be contributing to rabies crossing the barrier.”