

<b>Centers for Medicare and Medicaid Services (CMS)/The Joint Commission</b>
<b>*Denotes NQF Endorsed</b>
<b><u>Acute Myocardial Infarction*</u></b>
<i>All-or-none measure (AMI-1, AMI-2, AMI-3, AMI-4, AMI-5, AMI-8a)</i>
<i>AMI-1: Aspirin at Arrival</i>
<i>AMI-2: Aspirin at Discharge</i>
<i>AMI-3: ACEI or ARB for LVSD</i>
<i>AMI-4: Smoking Cessation Counseling</i>
<i>AMI-5: Beta Blocker at Discharge</i>
<i>AMI-8a: Primary PCI received within 90 minutes of hospital arrival</i>
<i>Mort-30-AMI: AMI 30-Day Mortality</i>
<b><u>Heart Failure*</u></b>
<i>All-or-none measure (HF-1, HF-2, HF-3, HF-4)</i>
<i>HF-1: Discharge Instructions</i>
<i>HF-2: Evaluation of left ventricular systolic (LVS) Function</i>
<i>HF-3: ACEI or ARB for LVSD</i>
<i>HF-4: Smoking Cessation Counseling</i>
<b><u>Pneumonia*</u></b>
<i>All-or-none measure ( PN-2, PN-3b, PN-4, PN-5c, PN-6, PN-7)</i>
<i>PN-2: Pneumococcal Vaccination</i>
<i>PN-3b: Blood Culture Performed in Emergency Department Prior to Initial Antibiotic Received in Hospital</i>
<i>PN-4: Smoking Cessation Counseling</i>
<i>PN5c: Initial Antibiotic received within 6 hours of hospital arrival</i>
<i>PN-6: Initial Antibiotic Selection for Community-Acquired Pneumonia in Immunocompetent Patients</i>
<i>PN-7: Influenza Vaccination</i>
<b><u>Surgical Care Improvement Project (SCIP)*</u></b>
<i>Appropriateness of care measure ( SCIP-Inf 1, 2, 3)</i>
<i>SCIP-Inf 1a: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Overall Rate</i>
<i>SCIP-Inf 1b: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - CABG</i>
<i>SCIP-Inf 1c: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Other Cardiac Surgery</i>
<i>SCIP-Inf 1d: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Hip Arthroplasty</i>
<i>SCIP-Inf 1e: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Knee Arthroplasty</i>
<i>SCIP-Inf 1f: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Colon Surgery</i>
<i>SCIP-Inf 1g: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Hysterectomy</i>
<i>SCIP-Inf 1h: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision - Vascular Surgery</i>
<i>SCIP-Inf-2a: Prophylactic Antibiotic Selection for Surgical Patients – Overall Rate</i>
<i>SCIP-Inf-2b: Prophylactic Antibiotic Selection for Surgical Patients - CABG</i>
<i>SCIP-Inf-2c: Prophylactic Antibiotic Selection for Surgical Patients - Other Cardiac Surgery</i>
<i>SCIP-Inf-2d: Prophylactic Antibiotic Selection for Surgical Patients - Hip Arthroplasty</i>
<i>SCIP-Inf-2e: Prophylactic Antibiotic Selection for Surgical Patients - Knee Arthroplasty</i>
<i>SCIP-Inf-2f: Prophylactic Antibiotic Selection for Surgical Patients - Colon Surgery</i>
<i>SCIP-Inf-2g: Prophylactic Antibiotic Selection for Surgical Patients - Hysterectomy</i>
<i>SCIP-Inf-2h: Prophylactic Antibiotic Selection for Surgical Patients - Vascular Surgery</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued within 24 Hours after Surgery End Time – Overall Rate</i>

## Appendix A

<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 48 Hours After Surgery End Time – CABG</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 48 Hours After Surgery End Time - Other Cardiac</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time - Hip Arthroplasty</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time - Knee Arthroplasty</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time - Colon Surgery</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time – Hysterectomy</i>
<i>SCIP-Inf-3a: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time - Vascular Surgery</i>
<i>SCIP-Card-2: Surgery Patients on Beta Blocker Therapy Prior to Admission who Received a Beta Blocker during</i>
<i>SCIP-VTE-1: Surgery Patients with Recommended Venous Thromboembolism Prophylaxis Ordered</i>
<i>SCIP-VTE-2: Surgery Patients who Received Appropriate Venous Thromboembolism Prophylaxis Within 24 Hours Prior to Surgery to 24 Hours after Surgery</i>
<b><u>Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS)</u></b>
<i>Communication with Nurses</i>
<i>Communication with Doctors</i>
<i>Responsiveness of Hospital Staff</i>
<i>Pain Management</i>
<i>Communication about Medicines</i>
<i>Cleanliness of Hospital Environment</i>
<i>Quietness of Hospital Environment</i>
<i>Discharge Information</i>
<i>Overall Rating of this Hospital</i>
<i>Willingness to Recommend This Hospital</i>
<b>Joint Commission</b>
<b><u>Stroke Measures</u></b>
<i>DSC/Stroke-01: Deep Vein Thrombosis (DVT) Prophylaxis</i>
<i>DSC/Stroke-02: Discharged on Antithrombotic Therapy</i>
<i>DSC/Stroke-03: Patients with Atrial Fibrillation Receiving Anticoagulation Therapy</i>
<i>DSC/Stroke-04: Thrombolytic Therapy Administered</i>
<i>DSC/Stroke-05: Antithrombotic Therapy by End of Hospital Day Two</i>
<i>DSC/Stroke-06: Discharged on Cholesterol Reducing Medication</i>
<i>DSC/Stroke-07: Dysphagia Screening</i>
<i>DSC/Stroke-08: Stroke Education</i>
<i>DSC/Stroke-09: Smoking Cessation /Advice/Counseling</i>
<i>DSC/Stroke-10: Assessed for Rehabilitation</i>
<b>Agency for Healthcare Research and Quality (AHRQ)</b>
<b><u>Other Heart Measures</u></b>
<i>IQI-6: Percutaneous Coronary Intervention (Angioplasty) (PCTA) volume</i>
<i>IQI-30: Percutaneous Coronary Intervention (Angioplasty) (PCTA) mortality rate</i>
<i>IQI-5: Coronary Artery Bypass Graft (CABG) volume</i>
<i>IQI-12: Coronary Artery Bypass Graft (CABG) mortality rate</i>
<b><u>Patient Safety Measures</u></b>
<i>PSI-1: Complications of Anesthesia</i>
<i>PSI-3: Decubitus Ulcer</i>
<i>PSI-5: Foreign Body Left During Procedure</i>

Appendix A

<b>Infection Measures</b> <b>*Denotes NQF Endorsed</b>
<b><u>Centers for Disease Control and Prevention- National Healthcare Safety Network (NHSN)</u></b>
<i>Surgical Site Infection Event - Coronary artery bypass graft with chest incision only *</i>
<i>Surgical Site Infection Event - Cesarean Section*</i>
<i>Surgical Site Infection Event - Knee Prosthesis, initial surgery only*</i>
<i>Hospital-Acquired Clostridium difficile (C. Diff.)</i>
<i>Hospital-Acquired Methicillin Resistant and Methicillin Susceptible Staphylococcus aureus Bacteremia (MRSA/MSSA Bacteremia) (SAB)</i>
<b><u>Infection Control Information</u></b>
<i>Hand-washing Program</i>
<i>Infection Control Staffing</i>
<b>Perinatal Measures</b> <b>*Denotes NQF Endorsed</b>
<b><u>California Maternal Quality Care Collaborative</u></b>
<i>Cesarean Rate for Low-Risk First Birth Women (NTSV CS Rate)*</i>
<i>Infants Under 1500g Not Delivered at Appropriate Level of Care*</i>
<b><u>Providence St. Vincent's Hospital/CWISH</u></b>
<i>Appropriate use of Antenatal Steroids*</i>

<b>Pediatric Measures</b> <b>*Denotes NQF Endorsed</b>
<b><u>The Joint Commission - Children's Asthma Care</u></b>
<i>CAC-1a: Systemic Corticosteroids for Inpatient Asthma (age 2-17) overall rate*</i>
<i>CAC-2a: Relievers for Inpatient Asthma (age 2-17) overall rate*</i>